

STATE CHILDHOOD IMMUNIZATION PROGRAM VACCINE ELIGIBILITY CHART

Vaccine	Those eligible for State-supplied Vaccine
DTaP (diphtheria/tetanus/acellular pertussis)	All children under 7 years of age.
DT (diphtheria/tetanus)	All children under 7 years of age with contraindications for pertussis.
Pediarix (diphtheria, tetanus, acellular pertussis, inactivated polio, and hepatitis B)	All VFC eligible children under 7 years of age.* Pediarix is approved for <u>first three doses of DTaP and IPV series only.</u>
Td (tetanus/diphtheria)	All children 7 years through 10 years of age.
Tdap	All children 11 through 14 years of age. May also be given to VFC- Eligible children 15 through 18 years of age.* May also be given to all children 7 through 10 years of age that did not complete a series of a pertussis containing vaccine prior to their 7 th birthday.
Hib (Haemophilus Influenza B)	All children under 5 years of age.
Pentacel (diphtheria, tetanus, acellular pertussis, Hib, and inactivated polio)	All VFC eligible children under 5 years of age. It may only be administered for the first 4 doses of the series. It should not be used as the booster dose.
IPV (inactivated polio vaccine)	All children 2 months through 18 years of age.
Kinrix (DTaP-Polio)	All VFC eligible children 4 through 6 years of age for the 5 th dose of DTaP and the 4 th or 5 th Dose of Polio.*
MMR (measles/mumps/rubella)	All children 12 months through 18 years of age.
Hepatitis B	All children-birth through 18 years of age. If the series is started at age 18 the patient may finish series at age 19.
Hepatitis A	All VFC-Eligible children 12 months through 18 years of age.* The two doses in the series should be administered at least 6 months apart.
PCV13 (Prevnar 13)	All VFC-Eligible children under 6 years of age.* VFC eligible children 60 through 71 months of age with underlying medical conditions that increase their risk for pneumococcal disease or complications. VFC eligible children 14 through 59 months who have received a complete series of PCV7 should receive a single dose of PCV13.
Varicella (chicken pox)	All children 12 to 15 months and 4 to 6 years of age. May also be given to VFC-eligible children 16 months through 18 years of age. *
Meningococcal (MCV4)	All VFC-eligible children 11 through 18 years of age as currently recommended.*
Rotavirus	All VFC-eligible age appropriate children.*
Human Papillomavirus (HPV)	All VFC-eligible children 11 through 18 years of age.*
Influenza	All children 6 months through 18 years of age.

* VFC-Underinsured need to receive their immunization at a FQHC/RHC/CHS facility. Non-VFC children will need to receive "privately purchased" vaccine for the VFC only vaccines. Check the most recent eligibility chart.

Revised 10/17/2011