


Map 1

Percent of Low Birth Weight Infants by County, 2006-2010

U.S. = 8.2% **South Dakota = 6.6%**




Note: "Significantly" refers to statistical significance at the 0.05 level. Although a county's actual percent may be higher or lower than the state percent, the difference may not be statistically significant due to the small number of people in the county. The percent of low birth weight infants is calculated based on the first weight of the newborn obtained after birth. Low birth weight infants are those born alive who weigh less than 2,500 grams (about 5 pounds 9 ounces). The U.S. percent of low birth weight infants is from 2009. See technical notes for more complete explanations.

Source: South Dakota Department of Health, Office of Health Statistics.

Map 2

Percent of Mothers Receiving Prenatal Care in the 1st Trimester by County, 2006-2010

U.S. = 71.0%* **South Dakota = 69.6%**



Note: "Significantly" refers to statistical significance at the 0.05 level. Although a county's actual percent may be higher or lower than the state percent, the difference may not be statistically significant due to the small number of people in the county. The rate is based on pregnant females with live births of one or more infants. The U.S. percent of mothers receiving prenatal care in the 1st trimester is based on 27 states that were using the 2003 standard birth certificate, including South Dakota. See technical notes for more complete explanations.


*Due to lack of comparable data from all 50 states, U.S. and South Dakota cannot be compared. The U.S. percent of first trimester prenatal care is from 2008.

Source: South Dakota Department of Health, Office of Health Statistics.

Map 4

Percent of Births Less Than 37 Weeks Gestation by County, 2006-2010

U.S. = 12.2% South Dakota = 8.8%




Note: "Significantly" refers to statistical significance at the 0.05 level. Although a county's actual percent may be higher or lower than the state percent, the difference may not be statistically significant due to the small number of people in the county. The U.S. percent of births less than 37 weeks gestation is from 2009. See technical notes for more complete explanations.
 Source: South Dakota Department of Health, Office of Health Statistics.

Map 5

Average Age of Mother by Resident County, 2006-2010

U.S. = 27.5* **South Dakota = 26.9**




Note: "Significantly" refers to statistical significance at the 0.05 level. Although a county's actual percent may be higher or lower than the state percent, the difference may not be statistically significant due to the small number of people in the county. The U.S. average age of mothers is from 2009.
 *Due to lack of comparable data from all 50 states, U.S. and South Dakota can not be compared.
 See technical notes for more complete explanations. Source: South Dakota Department of Health, Office of Health Statistics.

Map 7

Death Rate Due to All Causes by County, 2006-2010

U.S. = 758.3 **South Dakota = 708.0**




Note: "Significantly" refers to statistical significance at the 0.05 level. Although a county's actual percent may be higher or lower than the state percent, the difference may not be statistically significant due to the small number of people in the county. The death rate is age-adjusted per 100,000 population. This eliminates age difference between populations, making them easier to compare. The U.S. age-adjusted death rate is from 2008.

See technical notes for more complete explanations. Source: South Dakota Department of Health, Office of Health Statistics.

Map 8

Death Rate Due to Heart Disease by County, 2006-2010

U.S. = 186.5 **South Dakota = 163.9**




Note: "Significantly" refers to statistical significance at the 0.05 level. Although a county's actual percent may be higher or lower than the state percent, the difference may not be statistically significant due to the small number of people in the county. The death rate is age-adjusted per 100,000 population. This eliminates age difference between populations, making them easier to compare. Heart disease is defined as ICD-10 codes I00-I09, I11, I13, and I20-I51. The U.S. age-adjusted Heart Disease death rate is from 2008. See technical notes for more complete explanations. Source: South Dakota Department of Health, Office of Health Statistics.

Map 9

Death Rate Due to Acute Myocardial Infarction by County, 2006-2010

U.S. = 40.7 **South Dakota = 57.1**




Note: "Significantly" refers to statistical significance at the 0.05 level. Although a county's actual percent may be higher or lower than the state percent, the difference may not be statistically significant due to the small number of people in the county. The death rate is age-adjusted per 100,000 population. This eliminates age difference between populations, making them easier to compare. Acute Myocardial Infarction is defined as ICD-10 codes I21-I22. The U.S. age-adjusted Acute Myocardial Infarction death rate is from 2008. See technical notes for more complete explanations. Source: South Dakota Department of Health, Office of Health Statistics.

Map 10

Death Rate Due to Atherosclerotic Heart Disease by County, 2006-2010

U.S. = ** South Dakota = 52.0




Note: "Significantly" refers to statistical significance at the 0.05 level. Although a county's actual percent may be higher or lower than the state percent, the difference may not be statistically significant due to the small number of people in the county. The death rate is age-adjusted per 100,000 population. This eliminates age difference between populations, making them easier to compare. Atherosclerotic Heart Disease is defined as ICD-10 code I25.1. Atherosclerotic Heart Disease I25.1 was not broken out in the national data; therefore, no age-adjusted rate is displayed. See technical notes for more complete explanations. Source: South Dakota Department of Health, Office of Health Statistics.

Map 11

Death Rate Due to Heart Failure by County, 2006-2010

U.S. = 16.9 **South Dakota = 7.4**




Note: "Significantly" refers to statistical significance at the 0.05 level. Although a county's actual percent may be higher or lower than the state percent, the difference may not be statistically significant due to the small number of people in the county. The death rate is age-adjusted per 100,000 population. This eliminates age difference between populations, making them easier to compare. Heart Failure is defined as ICD-10 code I50. The U.S. age-adjusted Heart Failure death rate is from 2008. See technical notes for more complete explanations. Source: South Dakota Department of Health, Office of Health Statistics.

Map 12

Death Rate Due to All Malignant Neoplasms (Cancer) by County, 2006-2010

U.S. = 175.3 **South Dakota = 166.3**




Note: "Significantly" refers to statistical significance at the 0.05 level. Although a county's actual percent may be higher or lower than the state percent, the difference may not be statistically significant due to the small number of people in the county. The death rate is age-adjusted per 100,000 population. This eliminates age difference between populations, making them easier to compare. Malignant Neoplasms (Cancer) are defined as ICD-10 codes C00-C97. The U.S. age-adjusted Malignant Neoplasms (Cancer) death rate is from 2008. See technical notes for more complete explanations. Source: South Dakota Department of Health, Office of Health Statistics.

Map 14

Death Rate Due to Colorectal Cancer by County, 2006-2010

U.S. = 16.4 **South Dakota = 16.4**




Note: "Significantly" refers to statistical significance at the 0.05 level. Although a county's actual percent may be higher or lower than the state percent, the difference may not be statistically significant due to the small number of people in the county. The death rate is age-adjusted per 100,000 population. This eliminates age difference between populations, making them easier to compare. Colorectal Cancer are defined as ICD-10 codes C18-C21. The U.S. age-adjusted Colorectal Cancer death rate is from 2008. See technical notes for more complete explanations. Source: South Dakota Department of Health, Office of Health Statistics.

Map 15

Death Rate Due to Female Breast Cancer by County, 2006-2010

U.S. = 22.5 **South Dakota = 20.1**




Note: "Significantly" refers to statistical significance at the 0.05 level. Although a county's actual percent may be higher or lower than the state percent, the difference may not be statistically significant due to the small number of people in the county. The death rate is age-adjusted per 100,000 population. This eliminates age difference between populations, making them easier to compare. Female Breast Cancer is defined as ICD-10 code C50. The U.S. age-adjusted Female Breast Cancer death rate is from 2008. See technical notes for more complete explanations. Source: South Dakota Department of Health, Office of Health Statistics.

Map 16

Death Rate Due to Prostate Cancer by County, 2006-2010

U.S. = 22.3 South Dakota = 22.7




Note: "Significantly" refers to statistical significance at the 0.05 level. Although a county's actual percent may be higher or lower than the state percent, the difference may not be statistically significant due to the small number of people in the county. The death rate is age-adjusted per 100,000 population. This eliminates age difference between populations, making them easier to compare. Prostate Cancer is defined as ICD-10 code C61. The U.S. age-adjusted Prostate Cancer death rate is from 2008. See technical notes for more complete explanations. Source: South Dakota Department of Health, Office of Health Statistics.

Map 17

Death Rate Due to Pancreatic Cancer by County, 2006-2010

U.S. = 10.9 **South Dakota = 9.9**




Note: "Significantly" refers to statistical significance at the 0.05 level. Although a county's actual percent may be higher or lower than the state percent, the difference may not be statistically significant due to the small number of people in the county. The death rate is age-adjusted per 100,000 population. This eliminates age difference between populations, making them easier to compare. Pancreatic Cancer is defined as ICD-10 code C25. The U.S. age-adjusted Pancreatic Cancer death rate is from 2008. See technical notes for more complete explanations. Source: South Dakota Department of Health, Office of Health Statistics.

Map 20

Death Rate Due to Cerebrovascular Disease by County, 2006-2010

U.S. = 40.7 **South Dakota = 39.3**




Note: "Significantly" refers to statistical significance at the 0.05 level. Although a county's actual percent may be higher or lower than the state percent, the difference may not be statistically significant due to the small number of people in the county. The death rate is age-adjusted per 100,000 population. This eliminates age difference between populations, making them easier to compare. Cerebrovascular disease is defined as ICD-10 code I60-I69. The U.S. age-adjusted Cerebrovascular disease death rate is from 2008. See technical notes for more complete explanations. Source: South Dakota Department of Health, Office of Health Statistics.

Map 22

Death Rate Due to Accidents by County, 2006-2010

U.S. = 38.8 **South Dakota = 43.7**




Note: "Significantly" refers to statistical significance at the 0.05 level. Although a county's actual percent may be higher or lower than the state percent, the difference may not be statistically significant due to the small number of people in the county. The death rate is age-adjusted per 100,000 population. This eliminates age difference between populations, making them easier to compare. Accidents are defined as ICD-10 codes V01-X59, Y85-Y86. The U.S. age-adjusted Accident death rate is from 2008. See technical notes for more complete explanations. Source: South Dakota Department of Health, Office of Health Statistics.

Map 23

Death Rate Due to Motor Vehicle Accidents by County, 2006-2010

U.S. = 12.9
South Dakota = 17.9




Note: "Significantly" refers to statistical significance at the 0.05 level. Although a county's actual percent may be higher or lower than the state percent, the difference may not be statistically significant due to the small number of people in the county. The death rate is age-adjusted per 100,000 population. This eliminates age difference between populations, making them easier to compare. Motor Vehicle Accidents are defined as ICD-10 codes (V02-V04, V09.0, V09.2, V12-V14, V19.0-V19.2, V19.4-V19.6, V20-V79, V80.3-V80.5, V81.0-V81.1, V82.0-V82.1, V83-V86, V87.0-V87.8, V88.0-V88.8, V89.0, and V89.2) The U.S. age-adjusted Motor Vehicle Accident death rate is from 2008. See technical notes for more complete explanations. Source: South Dakota Department of Health, Office of Health Statistics.

Map 24

Death Rate Due to Alzheimer's Disease by County, 2006-2010

U.S. = 24.4 **South Dakota = 33.6**



Note: "Significantly" refers to statistical significance at the 0.05 level. Although a county's actual percent may be higher or lower than the state percent, the difference may not be statistically significant due to the small number of people in the county. The death rate is age-adjusted per 100,000 population. This eliminates age difference between populations, making them easier to compare. Alzheimer's Disease is defined as ICD-10 code G30. The U.S. age-adjusted Alzheimer's Disease death rate is from 2008. See technical notes for more complete explanations. Source: South Dakota Department of Health, Office of Health Statistics.

