





# South Dakota

Demographic Information	Health Status Indicators 2008-2012																																																												
 <p>South Dakota is located in the north central portion of the United States and averages 10.7 persons per square mile.</p> <p style="text-align: center;"><b>2012 Population Information</b></p> <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="text-align: left;">Subject</th> <th style="text-align: right;">Number</th> <th style="text-align: right;">Percent</th> </tr> </thead> <tbody> <tr> <td>Total population</td> <td style="text-align: right;">833,354</td> <td style="text-align: right;">100</td> </tr> <tr> <td>White</td> <td style="text-align: right;">718,352</td> <td style="text-align: right;">86.2</td> </tr> <tr> <td>Black or African American</td> <td style="text-align: right;">13,939</td> <td style="text-align: right;">1.7</td> </tr> <tr> <td>American Indian &amp; Alaska Native</td> <td style="text-align: right;">74,223</td> <td style="text-align: right;">8.9</td> </tr> <tr> <td>Asian</td> <td style="text-align: right;">9,005</td> <td style="text-align: right;">1.1</td> </tr> <tr> <td>Native Hawaiian &amp; Other Pacific Islander</td> <td style="text-align: right;">608</td> <td style="text-align: right;">0.1</td> </tr> <tr> <td>Two or More Races</td> <td style="text-align: right;">17,227</td> <td style="text-align: right;">2.1</td> </tr> <tr> <td>Under 5 years</td> <td style="text-align: right;">59,202</td> <td style="text-align: right;">7.1</td> </tr> <tr> <td>Under 18 years</td> <td style="text-align: right;">204,169</td> <td style="text-align: right;">24.5</td> </tr> <tr> <td>65 years and over</td> <td style="text-align: right;">122,183</td> <td style="text-align: right;">14.7</td> </tr> </tbody> </table> <p>Source: United States Census Bureau, 2012 Population Estimates</p>	Subject	Number	Percent	Total population	833,354	100	White	718,352	86.2	Black or African American	13,939	1.7	American Indian & Alaska Native	74,223	8.9	Asian	9,005	1.1	Native Hawaiian & Other Pacific Islander	608	0.1	Two or More Races	17,227	2.1	Under 5 years	59,202	7.1	Under 18 years	204,169	24.5	65 years and over	122,183	14.7	<p style="text-align: center;"><b>Nativity</b></p> <ul style="list-style-type: none"> <li>● Percent of Low Birth Weight Infants 6.3</li> <li>* Percent of Mothers Receiving Care in 1st Trimester 69.4</li> <li>* Percent of Mothers Who Used Tobacco While Pregnant<sup>1</sup> 17.5</li> <li>Percent of Births Less Than 37 Wks. of Gestation 8.2</li> <li>Average Age of Mother 27.1</li> <li>Teenage Birth Rate<sup>2</sup> 17.4</li> <li>Percent White Births 77.6</li> <li>Percent American Indian Births 16.2</li> <li>Percent Unmarried 38.4</li> <li>Percent WIC births 39.4</li> <li>Percent Breastfeeding at discharge 74.3</li> <li>Percent Payment-Private Insurance 55.5</li> <li>Percent Payment-Medicaid 35.1</li> <li>Percent C-Section 26.0</li> </ul> <p>●Denotes a health status indicator which is significantly lower than the national average. ○Denotes a health status indicator which is significantly higher than the national average. *Due to lack of comparable data from all 50 states, significant differences cannot be calculated. <sup>1</sup>Data for mothers who used tobacco are self-reported. <sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.</p>	<p style="text-align: center;"><b>Mortality<sup>3</sup></b></p> <ul style="list-style-type: none"> <li>● All Causes 707.3</li> <li>● Heart Disease 158.4</li> <li>● Malignant Neoplasms (cancer) 164.5                             <ul style="list-style-type: none"> <li>● Trachea, Bronchus, &amp; Lung 44.8</li> <li>Colon, Rectum, &amp; Anus 15.8</li> <li>Female Breast 20.7</li> <li>Pancreas 10.1</li> </ul> </li> <li>○ Cerebrovascular Disease 39.0</li> <li>○ Chronic Lower Respiratory Diseases 46.3</li> <li>○ Accidents 43.5                             <ul style="list-style-type: none"> <li>○ Motor Vehicle Accidents 15.5</li> </ul> </li> <li>○ Alzheimer's Disease 36.7</li> <li>○ Diabetes Mellitus 23.0</li> <li>Influenza &amp; Pneumonia 15.7</li> <li>○ Intentional Self-Harm (suicide) 16.1</li> <li>○ Chronic Liver Disease and Cirrhosis 10.9</li> <li>○ Infant Mortality 7.40</li> </ul> <p style="text-align: center;"><b>Leading Causes of Death</b></p> <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="text-align: left;">Leading Causes of Death</th> <th style="text-align: right;">Total Deaths</th> </tr> </thead> <tbody> <tr><td>1. Heart Disease</td><td style="text-align: right;">8,333</td></tr> <tr><td>2. Malignant Neoplasms (cancer)</td><td style="text-align: right;">7,997</td></tr> <tr><td>3. Chronic Lower Respiratory Diseases</td><td style="text-align: right;">2,341</td></tr> <tr><td>4. Alzheimer's Disease</td><td style="text-align: right;">2,089</td></tr> <tr><td>5. Cerebrovascular Diseases</td><td style="text-align: right;">2,071</td></tr> <tr><td>6. Accidents</td><td style="text-align: right;">1,935</td></tr> <tr><td>7. Diabetes</td><td style="text-align: right;">1,143</td></tr> <tr><td>8. Influenza and Pneumonia</td><td style="text-align: right;">853</td></tr> <tr><td>9. Suicide</td><td style="text-align: right;">650</td></tr> <tr><td>10. Unspecified Dementia</td><td style="text-align: right;">489</td></tr> <tr><td>Percent of Deaths due to tobacco use</td><td style="text-align: right;">17.5</td></tr> <tr><td>Median age at death</td><td style="text-align: right;">80</td></tr> </tbody> </table> <p>●Denotes a health status indicator which is significantly lower than the national average. ○Denotes a health status indicator which is significantly higher than the national average. <sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.</p> <p>See technical notes for more information. Source: South Dakota Department of Health, Office of Health Statistics</p>	Leading Causes of Death	Total Deaths	1. Heart Disease	8,333	2. Malignant Neoplasms (cancer)	7,997	3. Chronic Lower Respiratory Diseases	2,341	4. Alzheimer's Disease	2,089	5. Cerebrovascular Diseases	2,071	6. Accidents	1,935	7. Diabetes	1,143	8. Influenza and Pneumonia	853	9. Suicide	650	10. Unspecified Dementia	489	Percent of Deaths due to tobacco use	17.5	Median age at death	80
Subject	Number	Percent																																																											
Total population	833,354	100																																																											
White	718,352	86.2																																																											
Black or African American	13,939	1.7																																																											
American Indian & Alaska Native	74,223	8.9																																																											
Asian	9,005	1.1																																																											
Native Hawaiian & Other Pacific Islander	608	0.1																																																											
Two or More Races	17,227	2.1																																																											
Under 5 years	59,202	7.1																																																											
Under 18 years	204,169	24.5																																																											
65 years and over	122,183	14.7																																																											
Leading Causes of Death	Total Deaths																																																												
1. Heart Disease	8,333																																																												
2. Malignant Neoplasms (cancer)	7,997																																																												
3. Chronic Lower Respiratory Diseases	2,341																																																												
4. Alzheimer's Disease	2,089																																																												
5. Cerebrovascular Diseases	2,071																																																												
6. Accidents	1,935																																																												
7. Diabetes	1,143																																																												
8. Influenza and Pneumonia	853																																																												
9. Suicide	650																																																												
10. Unspecified Dementia	489																																																												
Percent of Deaths due to tobacco use	17.5																																																												
Median age at death	80																																																												

# Aurora County

## Demographic Information



Aurora County is located in the south central portion of the state and averages 3.8 persons per square mile.

### 2012 Population Information

Subject	Number	Percent
Total population	2,742	100
White	2,639	96.2
Black or African American	16	0.6
American Indian & Alaska Native	47	1.7
Asian	20	0.7
Native Hawaiian & Other Pacific Islander	0	0.0
Two or More Races	20	0.7
Under 5 years	185	6.7
Under 18 years	728	26.5
65 years and over	549	20.0

Source: United States Census Bureau, 2012 Population Estimates

## Health Status Indicators 2008-2012

### Nativity

Percent of Low Birth Weight Infants	6.5
Percent of Mothers Receiving Care in 1st Trimester	71.1
Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	14.3
Percent of Births Less Than 37 Wks. of Gestation	10.7
○ Average Age of Mother	28.0
Teenage Birth Rate <sup>2</sup>	LNE
Percent White Births	98.2
Percent American Indian Births	LNE
● Percent Unmarried	22.6
Percent WIC births	37.7
Percent Breastfeeding at discharge	78.0
○ Percent Payment-Private Insurance	73.8
● Percent Payment-Medicaid	20.8
Percent C-Section	35.1

●Denotes a health status indicator which is significantly lower than the state average.  
 ○Denotes a health status indicator which is significantly higher than the state average.  
 \*Due to lack of comparable data from all 50 states, significant differences cannot be calculated.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

● All Causes	589.1
Heart Disease	155.9
Malignant Neoplasms (cancer)	153.4
Trachea, Bronchus, & Lung	60.3
Colon, Rectum, & Anus	17.1
Female Breast	40.7
Pancreas	LNE
Cerebrovascular Disease	22.7
Chronic Lower Respiratory Diseases	35.3
Accidents	28.6
Motor Vehicle Accidents	LNE
Alzheimer's Disease	23.5
Diabetes Mellitus	35.8
Influenza & Pneumonia	8.8
Intentional Self-Harm (suicide)	LNE
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	LNE

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	40
2. Malignant Neoplasms (cancer)	34
3. Chronic Lower Respiratory Diseases	9
T4. Alzheimer's Disease	8
T4. Diabetes	8
6. Accidents	7
7. Cerebrovascular Diseases	6
Percent of Deaths due to tobacco use	16.6
Median age at death	83

●Denotes a health status indicator which is significantly lower than the state average.  
 ○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Beadle County

## Demographic Information



Beadle County is located in the center of eastern South Dakota and averages 13.8 persons per square mile.

### 2012 Population Information

Subject	Number	Percent
Total population	17,753	100.0
White	16,064	90.5
Black or African American	268	1.5
American Indian & Alaska Native	226	1.3
Asian	896	5.0
Native Hawaiian & Other Pacific Islander	26	0.1
Two or More Races	273	1.5
Under 5 years	1,387	7.8
Under 18 years	4,356	24.5
65 years and over	3,012	17.0

Source: United States Census Bureau, 2012 Population Estimates

## Health Status Indicators 2008-2012

### Nativity

Percent of Low Birth Weight Infants	7.3
● Percent of Mothers Receiving Care in 1st Trimester	56.5
Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	18.3
Percent of Births Less Than 37 Wks. of Gestation	9.0
● Average Age of Mother	26.3
Teenage Birth Rate <sup>2</sup>	23.2
Percent White Births	86.3
Percent American Indian Births	1.0
Percent Unmarried	38.9
○ Percent WIC births	52.1
Percent Breastfeeding at discharge	71.8
● Percent Payment-Private Insurance	49.4
Percent Payment-Medicaid	38.7
○ Percent C-Section	33.1

●Denotes a health status indicator which is significantly lower than the state average.  
 ○Denotes a health status indicator which is significantly higher than the state average.  
 \*Due to lack of comparable data from all 50 states, significant differences cannot be calculated.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

All Causes	737.7
Heart Disease	165.4
Malignant Neoplasms (cancer)	168.8
Trachea, Bronchus, & Lung	35.1
Colon, Rectum, & Anus	19.2
Female Breast	26.5
Pancreas	10.5
Cerebrovascular Disease	45.2
Chronic Lower Respiratory Diseases	41.4
Accidents	38.5
Motor Vehicle Accidents	14.9
Alzheimer's Disease	36.3
○ Diabetes Mellitus	39.8
Influenza & Pneumonia	20.3
● Intentional Self-Harm (suicide)	7.6
● Chronic Liver Disease and Cirrhosis	3.8
Infant Mortality	13.89

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	235
2. Malignant Neoplasms (cancer)	206
3. Cerebrovascular Diseases	64
4. Alzheimer's Disease	59
5. Chronic Lower Respiratory Diseases	56
6. Diabetes	51
7. Accidents	34
8. Influenza and Pneumonia	30
9. Nephritis, Nephrotic Syndrome, and	14
T10. Certain Conditions Originating in the Perinatal Period	13
T10. Essential (primary) Hypertension and Hypertensive Renal Disease	13
Percent of Deaths due to tobacco use	17.9
Median age at death	82

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Bennett County

## Demographic Information



Bennett County is located on the southern border in western South Dakota and averages 2.9 persons per square mile.

### 2012 Population Information

Subject	Number	Percent
Total population	3,436	100.0
White	1,161	33.8
Black or African American	10	0.3
American Indian & Alaska Native	2,077	60.4
Asian	19	0.6
Native Hawaiian & Other Pacific Islander	2	0.1
Two or More Races	167	4.9
Under 5 years	329	9.6
Under 18 years	1,141	33.2
65 years and over	401	11.7

Source: United States Census Bureau, 2012 Population Estimates

## Health Status Indicators 2008-2012

### Nativity

Percent of Low Birth Weight Infants	4.9
● Percent of Mothers Receiving Care in 1st Trimester	41.5
Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	20.7
Percent of Births Less Than 37 Wks. of Gestation	8.5
● Average Age of Mother	24.7
○ Teenage Birth Rate <sup>2</sup>	42.9
Percent White Births	17.3
Percent American Indian Births	77.5
○ Percent Unmarried	73.8
○ Percent WIC births	77.3
● Percent Breastfeeding at discharge	50.4
● Percent Payment-Private Insurance	11.7
○ Percent Payment-Medicaid	65.5
Percent C-Section	21.9

●Denotes a health status indicator which is significantly lower than the state average.  
○Denotes a health status indicator which is significantly higher than the state average.

\*Due to lack of comparable data from all 50 states, significant differences cannot be calculated.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

○ All Causes	922.6
Heart Disease	172.4
Malignant Neoplasms (cancer)	177.0
Trachea, Bronchus, & Lung	51.1
Colon, Rectum, & Anus	41.2
Female Breast	LNE
Pancreas	LNE
Cerebrovascular Disease	35.3
Chronic Lower Respiratory Diseases	52.5
Accidents	70.7
○ Motor Vehicle Accidents	55.5
Alzheimer's Disease	LNE
Diabetes Mellitus	35.6
○ Influenza & Pneumonia	50.7
Intentional Self-Harm (suicide)	22.0
Chronic Liver Disease and Cirrhosis	46.9
Infant Mortality	11.53

### Leading Causes of Death

Leading Causes of Death	Total Deaths
T1. Heart Disease	29
T1. Malignant Neoplasms (cancer)	29
3. Accidents	12
T4. Chronic Lower Respiratory Diseases	9
T4. Influenza and Pneumonia	9
T6. Chronic Liver Disease and Cirrhosis	6
T6. Cerebrovascular Diseases	6
T6. Diabetes	6
Percent of Deaths due to tobacco use	21.6
Median age at death	73

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Bon Homme County

## Demographic Information



Bon Homme County is located on the Nebraska border in eastern South Dakota and averages 12.5 persons per square mile.

### 2012 Population Information

Subject	Number	Percent
Total population	7,029	100.0
White	6,309	89.8
Black or African American	86	1.2
American Indian & Alaska Native	507	7.2
Asian	19	0.3
Native Hawaiian & Other Pacific Islander	1	0.0
Two or More Races	107	1.5
Under 5 years	314	4.5
Under 18 years	1,353	19.2
65 years and over	1,380	19.6

Source: United States Census Bureau, 2012 Population Estimates

## Health Status Indicators 2008-2012

### Nativity

Percent of Low Birth Weight Infants	6.3
○ Percent of Mothers Receiving Care in 1st Trimester	80.0
● Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	12.4
Percent of Births Less Than 37 Wks. of Gestation	6.9
○ Average Age of Mother	28.0
Teenage Birth Rate <sup>2</sup>	LNE
Percent White Births	94.6
Percent American Indian Births	3.0
● Percent Unmarried	19.9
Percent WIC births	44.0
Percent Breastfeeding at discharge	75.7
○ Percent Payment-Private Insurance	74.3
● Percent Payment-Medicaid	21.1
Percent C-Section	28.3

●Denotes a health status indicator which is significantly lower than the state average.  
 ○Denotes a health status indicator which is significantly higher than the state average.  
 \*Due to lack of comparable data from all 50 states, significant differences cannot be calculated.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

● All Causes	583.6
Heart Disease	139.8
Malignant Neoplasms (cancer)	139.5
Trachea, Bronchus, & Lung	53.5
Colon, Rectum, & Anus	10.9
Female Breast	LNE
Pancreas	LNE
Cerebrovascular Disease	41.6
Chronic Lower Respiratory Diseases	43.6
Accidents	30.2
Motor Vehicle Accidents	17.4
● Alzheimer's Disease	22.4
Diabetes Mellitus	18.5
Influenza & Pneumonia	8.3
Intentional Self-Harm (suicide)	15.3
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	LNE

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	95
2. Malignant Neoplasms (cancer)	77
T3. Chronic Lower Respiratory Diseases	28
T3. Cerebrovascular Diseases	28
5. Alzheimer's Disease	21
6. Accidents	15
7. Diabetes	11
8. Neoplasms of Uncertain or Unknown Behavior	9
9. Unspecified Dementia	7
T10. Influenza and Pneumonia	6
T10. Parkinson's Disease	6
Percent of Deaths due to tobacco use	14.6
Median age at death	84

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Brookings County

## Demographic Information



Brookings County is located on the eastern border of the state and averages 40.3 persons per square mile.

### 2012 Population Information

Subject	Number	Percent
Total population	32,629	100.0
White	30,482	93.4
Black or African American	405	1.2
American Indian & Alaska Native	390	1.2
Asian	901	2.8
Native Hawaiian & Other Pacific Islander	27	0.1
Two or More Races	424	1.3
Under 5 years	1,918	5.9
Under 18 years	6,107	18.7
65 years and over	3,276	10.0

Source: United States Census Bureau, 2012 Population Estimates

## Health Status Indicators 2008-2012

### Natality

● Percent of Low Birth Weight Infants	5.1
○ Percent of Mothers Receiving Care in 1st Trimester	75.7
● Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	11.8
Percent of Births Less Than 37 Wks. of Gestation	6.9
○ Average Age of Mother	27.8
● Teenage Birth Rate <sup>2</sup>	6.9
Percent White Births	90.0
Percent American Indian Births	1.9
● Percent Unmarried	23.9
● Percent WIC births	27.9
○ Percent Breastfeeding at discharge	82.1
○ Percent Payment-Private Insurance	74.4
● Percent Payment-Medicaid	20.4
● Percent C-Section	21.0

●Denotes a health status indicator which is significantly lower than the state average.  
 ○Denotes a health status indicator which is significantly higher than the state average.  
 \*Due to lack of comparable data from all 50 states, significant differences cannot be calculated.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

All Causes	668.8
Heart Disease	144.8
Malignant Neoplasms (cancer)	158.4
Trachea, Bronchus, & Lung	34.3
Colon, Rectum, & Anus	13.2
Female Breast	19.9
Pancreas	9.0
Cerebrovascular Disease	47.9
Chronic Lower Respiratory Diseases	38.6
● Accidents	30.7
● Motor Vehicle Accidents	8.1
Alzheimer's Disease	40.4
● Diabetes Mellitus	12.1
Influenza & Pneumonia	14.7
● Intentional Self-Harm (suicide)	7.0
● Chronic Liver Disease and Cirrhosis	5.5
Infant Mortality	8.20

Leading Causes of Death	Total Deaths
1. Heart Disease	222
2. Malignant Neoplasms (cancer)	217
3. Cerebrovascular Diseases	76
4. Alzheimer's Disease	66
5. Chronic Lower Respiratory Diseases	55
6. Accidents	46
7. Influenza and Pneumonia	23
8. Unspecified Dementia	22
9. Diabetes	17
10. Congenital Malformations, Deformations and Chromosomal Abnormalities	13
Percent of Deaths due to tobacco use	17.9
Median age at death	83

●Denotes a health status indicator which is significantly lower than the state average.  
 ○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Brown County

## Demographic Information



Brown County is located on the North Dakota border in the eastern part of the state and averages 21.3 persons per square mile.

### 2012 Population Information

Subject	Number	Percent
Total population	37,331	100.0
White	34,659	92.8
Black or African American	408	1.1
American Indian & Alaska Native	1,181	3.2
Asian	430	1.2
Native Hawaiian & Other Pacific Islander	48	0.1
Two or More Races	605	1.6
Under 5 years	2,537	6.8
Under 18 years	8,587	23.0
65 years and over	5,921	15.9

Source: United States Census Bureau, 2012 Population Estimates

## Health Status Indicators 2008-2012

### Natality

Percent of Low Birth Weight Infants	5.6
○ Percent of Mothers Receiving Care in 1st Trimester	77.0
Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	17.5
Percent of Births Less Than 37 Wks. of Gestation	8.1
Average Age of Mother	27.1
● Teenage Birth Rate <sup>2</sup>	11.9
Percent White Births	90.9
Percent American Indian Births	5.6
● Percent Unmarried	34.2
● Percent WIC births	33.8
Percent Breastfeeding at discharge	78.3
○ Percent Payment-Private Insurance	64.2
● Percent Payment-Medicaid	29.4
Percent C-Section	28.0

●Denotes a health status indicator which is significantly lower than the state average.  
 ○Denotes a health status indicator which is significantly higher than the state average.  
 \*Due to lack of comparable data from all 50 states, significant differences cannot be calculated.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

All Causes	692.5
Heart Disease	173.9
Malignant Neoplasms (cancer)	155.8
Trachea, Bronchus, & Lung	42.1
Colon, Rectum, & Anus	15.8
Female Breast	23.1
Pancreas	11.1
Cerebrovascular Disease	36.2
Chronic Lower Respiratory Diseases	41.0
● Accidents	32.7
Motor Vehicle Accidents	10.2
Alzheimer's Disease	44.3
Diabetes Mellitus	22.4
Influenza & Pneumonia	17.4
Intentional Self-Harm (suicide)	11.5
Chronic Liver Disease and Cirrhosis	8.8
Infant Mortality	5.03

### Leading Causes of Death

	Total Deaths
1. Heart Disease	491
2. Malignant Neoplasms (cancer)	366
3. Alzheimer's Disease	135
4. Chronic Lower Respiratory Diseases	108
5. Cerebrovascular Diseases	100
6. Accidents	69
7. Diabetes	58
8. Influenza and Pneumonia	51
9. Pneumonitis Due to Solids and Liquids	27
10. Nephritis, Nephrotic Syndrome, and Nephrosis	26

Percent of Deaths due to tobacco use	15.2
Median age at death	83

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Brule County

## Demographic Information



Brule County is located in the south central part of the state and averages 6.4 persons per square mile.

## 2012 Population Information

Subject	Number	Percent
Total population	5,293	100.0
White	4,668	88.2
Black or African American	26	0.5
American Indian & Alaska Native	446	8.4
Asian	11	0.2
Native Hawaiian & Other Pacific Islander	4	0.1
Two or More Races	138	2.6
Under 5 years	361	6.8
Under 18 years	1,322	25.0
65 years and over	905	17.1

Source: United States Census Bureau, 2012 Population Estimates

## Health Status Indicators 2008-2012

### Natality

Percent of Low Birth Weight Infants	5.9
Percent of Mothers Receiving Care in 1st Trimester	65.5
Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	19.0
Percent of Births Less Than 37 Wks. of Gestation	9.2
Average Age of Mother	27.4
• Teenage Birth Rate <sup>2</sup>	9.1
Percent White Births	77.3
Percent American Indian Births	19.4
Percent Unmarried	35.7
Percent WIC births	39.5
Percent Breastfeeding at discharge	72.4
Percent Payment-Private Insurance	64.0
Percent Payment-Medicaid	33.9
Percent C-Section	25.9

•Denotes a health status indicator which is significantly lower than the state average.  
 ○Denotes a health status indicator which is significantly higher than the state average.  
 \*Due to lack of comparable data from all 50 states, significant differences cannot be calculated.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

All Causes	688.0
Heart Disease	148.4
Malignant Neoplasms (cancer)	189.8
Trachea, Bronchus, & Lung	54.0
Colon, Rectum, & Anus	23.0
Female Breast	17.2
Pancreas	7.4
Cerebrovascular Disease	29.6
Chronic Lower Respiratory Diseases	39.1
Accidents	53.2
Motor Vehicle Accidents	13.5
Alzheimer's Disease	32.1
Diabetes Mellitus	35.8
Influenza & Pneumonia	15.7
Intentional Self-Harm (suicide)	LNE
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	10.81

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Malignant Neoplasms (cancer)	70
2. Heart Disease	61
3. Alzheimer's Disease	18
4. Accidents	17
5. Chronic Lower Respiratory Diseases	16
6. Diabetes	13
7. Cerebrovascular Diseases	12
8. Influenza and Pneumonia	7

Percent of Deaths due to tobacco use	23.4
Median age at death	82

•Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Buffalo County

## Demographic Information



Buffalo County is located in the central portion of the state and averages 4.1 persons per square mile.

## 2012 Population Information

Subject	Number	Percent
Total population	2,020	100.0
White	348	17.2
Black or African American	9	0.4
American Indian & Alaska Native	1,627	80.5
Asian	1	0.0
Native Hawaiian & Other Pacific Islander	0	0.0
Two or More Races	35	1.7
Under 5 years	269	13.3
Under 18 years	830	41.1
65 years and over	152	7.5

Source: United States Census Bureau, 2012 Population Estimates

## Health Status Indicators 2008-2012

### Nativity

Percent of Low Birth Weight Infants	6.4
● Percent of Mothers Receiving Care in 1st Trimester	44.8
○ Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	33.5
Percent of Births Less Than 37 Wks. of Gestation	10.8
● Average Age of Mother	25.9
○ Teenage Birth Rate <sup>2</sup>	68.7
Percent White Births	15.9
Percent American Indian Births	77.7
○ Percent Unmarried	74.1
○ Percent WIC births	78.3
● Percent Breastfeeding at discharge	44.8
● Percent Payment-Private Insurance	19.1
○ Percent Payment-Medicaid	74.9
Percent C-Section	26.3

●Denotes a health status indicator which is significantly lower than the state average.  
○Denotes a health status indicator which is significantly higher than the state average.

\*Due to lack of comparable data from all 50 states, significant differences cannot be calculated.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

○ All Causes	1,336.5
Heart Disease	282.5
Malignant Neoplasms (cancer)	243.6
Trachea, Bronchus, & Lung	LNE
Colon, Rectum, & Anus	LNE
Female Breast	LNE
Pancreas	LNE
Cerebrovascular Disease	62.6
Chronic Lower Respiratory Diseases	56.5
Accidents	105.6
Motor Vehicle Accidents	26.7
Alzheimer's Disease	LNE
Diabetes Mellitus	64.5
Influenza & Pneumonia	LNE
Intentional Self-Harm (suicide)	63.1
○ Chronic Liver Disease and Cirrhosis	120.4
Infant Mortality	23.90

### Leading Causes of Death

	Total Deaths
1. Heart Disease	18
2. Malignant Neoplasms (cancer)	15
3. Accidents	10
4. Chronic Liver Disease and Cirrhosis	9
5. Suicide	6
6. Diabetes	5
Percent of Deaths due to tobacco use	27.1
Median age at death	57

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Butte County

## Demographic Information



Butte County is located on the western border of the state and averages 4.5 people per square mile.

### 2012 Population Information

Subject	Number	Percent
Total population	10,228	100.0
White	9,685	94.7
Black or African American	56	0.5
American Indian & Alaska Native	185	1.8
Asian	30	0.3
Native Hawaiian & Other Pacific Islander	12	0.1
Two or More Races	260	2.5
Under 5 years	681	6.7
Under 18 years	2,474	24.2
65 years and over	1,743	17.0

Source: United States Census Bureau, 2012 Population Estimates

## Health Status Indicators 2008-2012

### Natality

Percent of Low Birth Weight Infants	5.1
○ Percent of Mothers Receiving Care in 1st Trimester	77.3
○ Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	25.2
Percent of Births Less Than 37 Wks. of Gestation	7.8
● Average Age of Mother	26.6
Teenage Birth Rate <sup>2</sup>	17.3
Percent White Births	93.5
Percent American Indian Births	1.8
Percent Unmarried	35.4
○ Percent WIC births	46.1
Percent Breastfeeding at discharge	78.5
Percent Payment-Private Insurance	50.1
○ Percent Payment-Medicaid	42.1
Percent C-Section	22.9

●Denotes a health status indicator which is significantly lower than the state average.  
 ○Denotes a health status indicator which is significantly higher than the state average.  
 \*Due to lack of comparable data from all 50 states, significant differences cannot be calculated.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

○ All Causes	822.2
Heart Disease	189.3
○ Malignant Neoplasms (cancer)	209.3
○ Trachea, Bronchus, & Lung	70.0
Colon, Rectum, & Anus	17.5
Female Breast	22.4
Pancreas	12.6
Cerebrovascular Disease	29.9
Chronic Lower Respiratory Diseases	42.1
Accidents	52.6
Motor Vehicle Accidents	28.6
Alzheimer's Disease	39.7
Diabetes Mellitus	23.9
Influenza & Pneumonia	26.7
Intentional Self-Harm (suicide)	25.7
Chronic Liver Disease and Cirrhosis	6.1
Infant Mortality	7.54

Leading Causes of Death	Total Deaths
1. Malignant Neoplasms (cancer)	136
2. Heart Disease	127
3. Chronic Lower Respiratory Diseases	28
4. Accidents	27
5. Alzheimer's Disease	26
6. Cerebrovascular Diseases	20
7. Influenza and Pneumonia	18
8. Diabetes	16
9. Suicide	13
T10. Unspecified Dementia	9
T10. Essential (primary) Hypertension and Hypertensive Renal Disease	9
Percent of Deaths due to tobacco use	19.4
Median age at death	79

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Campbell County

## Demographic Information



Campbell County is located in the north central portion of the state and averages 2.0 persons per square mile.

### 2012 Population Information

Subject	Number	Percent
Total population	1,396	100.0
White	1,371	98.2
Black or African American	5	0.4
American Indian & Alaska Native	7	0.5
Asian	7	0.5
Native Hawaiian & Other Pacific Islander	0	0.0
Two or More Races	6	0.4
Under 5 years	50	3.6
Under 18 years	241	17.3
65 years and over	358	25.6

Source: United States Census Bureau, 2012 Population Estimates

## Health Status Indicators 2008-2012

### Nativity

Percent of Low Birth Weight Infants	LNE
Percent of Mothers Receiving Care in 1st Trimester	73.1
Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	LNE
Percent of Births Less Than 37 Wks. of Gestation	7.6
○ Average Age of Mother	29.1
Teenage Birth Rate <sup>2</sup>	LNE
Percent White Births	96.9
Percent American Indian Births	LNE
● Percent Unmarried	14.7
Percent WIC births	32.3
Percent Breastfeeding at discharge	86.2
○ Percent Payment-Private Insurance	81.0
● Percent Payment-Medicaid	6.3
○ Percent C-Section	44.8

●Denotes a health status indicator which is significantly lower than the state average.  
○Denotes a health status indicator which is significantly higher than the state average.

\*Due to lack of comparable data from all 50 states, significant differences cannot be calculated.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

All Causes	584.5
Heart Disease	163.9
Malignant Neoplasms (cancer)	138.9
Trachea, Bronchus, & Lung	LNE
Colon, Rectum, & Anus	36.5
Female Breast	46.6
Pancreas	LNE
Cerebrovascular Disease	49.8
Chronic Lower Respiratory Diseases	LNE
Accidents	96.5
Motor Vehicle Accidents	LNE
Alzheimer's Disease	20.2
Diabetes Mellitus	LNE
Influenza & Pneumonia	LNE
Intentional Self-Harm (suicide)	LNE
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	LNE

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	21
2. Malignant Neoplasms (cancer)	19
3. Cerebrovascular Diseases	8
4. Accidents	7
Percent of Deaths due to tobacco use	19.5
Median age at death	83

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Charles Mix County

## Demographic Information



Charles Mix County is located in the south central area of the state and averages 8.3 persons per square mile.

### 2012 Population Information

Subject	Number	Percent
Total population	9,216	100.0
White	5,950	64.6
Black or African American	33	0.4
American Indian & Alaska Native	2,944	31.9
Asian	23	0.2
Native Hawaiian & Other Pacific Islander	1	0.0
Two or More Races	265	2.9
Under 5 years	800	8.7
Under 18 years	2,722	29.5
65 years and over	1,628	17.7

Source: United States Census Bureau, 2012 Population Estimates

## Health Status Indicators 2008-2012

### Natality

Percent of Low Birth Weight Infants	4.7
● Percent of Mothers Receiving Care in 1st Trimester	61.8
○ Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	24.5
Percent of Births Less Than 37 Wks. of Gestation	7.7
● Average Age of Mother	25.5
○ Teenage Birth Rate <sup>2</sup>	28.7
Percent White Births	43.5
Percent American Indian Births	51.1
○ Percent Unmarried	58.1
○ Percent WIC births	63.0
● Percent Breastfeeding at discharge	67.0
● Percent Payment-Private Insurance	35.3
○ Percent Payment-Medicaid	58.6
Percent C-Section	25.1

●Denotes a health status indicator which is significantly lower than the state average.  
○Denotes a health status indicator which is significantly higher than the state average.  
\*Due to lack of comparable data from all 50 states, significant differences cannot be calculated.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

All Causes	753.6
Heart Disease	187.6
Malignant Neoplasms (cancer)	180.6
Trachea, Bronchus, & Lung	32.9
Colon, Rectum, & Anus	20.7
Female Breast	17.8
Pancreas	16.7
● Cerebrovascular Disease	22.0
Chronic Lower Respiratory Diseases	38.9
Accidents	57.4
Motor Vehicle Accidents	26.0
Alzheimer's Disease	37.9
○ Diabetes Mellitus	40.5
Influenza & Pneumonia	16.6
Intentional Self-Harm (suicide)	12.8
Chronic Liver Disease and Cirrhosis	21.7
Infant Mortality	6.22

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	129
2. Malignant Neoplasms (cancer)	117
3. Alzheimer's Disease	30
4. Accidents	29
5. Chronic Lower Respiratory Diseases	28
6. Diabetes	25
7. Cerebrovascular Diseases	17
8. Influenza and Pneumonia	12
9. Chronic Liver Disease and Cirrhosis	11
10. Essential (primary) Hypertension and Hypertensive Renal Disease	10
Percent of Deaths due to tobacco use	14.6
Median age at death	80

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Clark County

## Demographic Information



Clark County is located in east central South Dakota and averages 3.9 persons per square mile.

### 2012 Population Information

Subject	Number	Percent
Total population	3,585	100.0
White	3,532	98.5
Black or African American	16	0.4
American Indian & Alaska Native	5	0.1
Asian	2	0.1
Native Hawaiian & Other Pacific Islander	0	0.0
Two or More Races	30	0.8
Under 5 years	266	7.4
Under 18 years	818	22.8
65 years and over	779	21.7

Source: United States Census Bureau, 2012 Population Estimates

## Health Status Indicators 2008-2012

### Nativity

Percent of Low Birth Weight Infants	4.1
● Percent of Mothers Receiving Care in 1st Trimester	57.3
Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	14.9
Percent of Births Less Than 37 Wks. of Gestation	5.8
○ Average Age of Mother	28.5
Teenage Birth Rate <sup>2</sup>	LNE
Percent White Births	97.5
Percent American Indian Births	1.7
● Percent Unmarried	20.2
● Percent WIC births	24.6
Percent Breastfeeding at discharge	85.4
○ Percent Payment-Private Insurance	78.2
● Percent Payment-Medicaid	18.5
Percent C-Section	21.9

●Denotes a health status indicator which is significantly lower than the state average.  
○Denotes a health status indicator which is significantly higher than the state average.  
\*Due to lack of comparable data from all 50 states, significant differences cannot be calculated.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

All Causes	719.3
Heart Disease	195.4
Malignant Neoplasms (cancer)	132.8
Trachea, Bronchus, & Lung	35.6
Colon, Rectum, & Anus	8.1
Female Breast	24.2
Pancreas	7.0
Cerebrovascular Disease	43.9
Chronic Lower Respiratory Diseases	50.1
Accidents	49.5
Motor Vehicle Accidents	LNE
Alzheimer's Disease	34.3
Diabetes Mellitus	25.5
Influenza & Pneumonia	23.9
Intentional Self-Harm (suicide)	LNE
Chronic Liver Disease and Cirrhosis	9.9
Infant Mortality	LNE

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	74
2. Malignant Neoplasms (cancer)	45
T3. Chronic Lower Respiratory Diseases	17
T3. Cerebrovascular Diseases	17
5. Alzheimer's Disease	15
6. Accidents	12
7. Influenza and Pneumonia	9
8. Diabetes	8
9. Gastrointestinal Hemorrhage, Unspecified	5
Percent of Deaths due to tobacco use	17.5
Median age at death	83

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Clay County

## Demographic Information



Clay County is located in the southeastern part of the state and averages 33.6 persons per square mile.

### 2012 Population Information

Subject	Number	Percent
Total population	14,131	100.0
White	12,846	90.9
Black or African American	257	1.8
American Indian & Alaska Native	448	3.2
Asian	283	2.0
Native Hawaiian & Other Pacific Islander	9	0.1
Two or More Races	288	2.0
Under 5 years	827	5.9
Under 18 years	2,650	18.8
65 years and over	1,483	10.5

Source: United States Census Bureau, 2012 Population Estimates

## Health Status Indicators 2008-2012

### Nativity

Percent of Low Birth Weight Infants	6.3
○ Percent of Mothers Receiving Care in 1st Trimester	78.5
● Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	10.8
Percent of Births Less Than 37 Wks. of Gestation	8.2
○ Average Age of Mother	27.7
● Teenage Birth Rate <sup>2</sup>	4.3
Percent White Births	91.4
Percent American Indian Births	4.0
● Percent Unmarried	27.9
Percent WIC births	38.5
Percent Breastfeeding at discharge	79.4
Percent Payment-Private Insurance	59.3
Percent Payment-Medicaid	36.5
○ Percent C-Section	33.2

●Denotes a health status indicator which is significantly lower than the state average.  
 ○Denotes a health status indicator which is significantly higher than the state average.  
 \*Due to lack of comparable data from all 50 states, significant differences cannot be calculated.

<sup>1</sup>Data for mothers who used tobacco are self-reported.  
<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

All Causes	725.7
Heart Disease	186.9
Malignant Neoplasms (cancer)	166.2
Trachea, Bronchus, & Lung	48.5
Colon, Rectum, & Anus	22.4
Female Breast	22.2
Pancreas	5.9
Cerebrovascular Disease	53.6
Chronic Lower Respiratory Diseases	54.0
Accidents	39.1
Motor Vehicle Accidents	11.6
● Alzheimer's Disease	20.9
Diabetes Mellitus	35.6
Influenza & Pneumonia	17.0
Intentional Self-Harm (suicide)	18.1
Chronic Liver Disease and Cirrhosis	8.3
Infant Mortality	6.42

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	125
2. Malignant Neoplasms (cancer)	100
3. Cerebrovascular Diseases	36
4. Chronic Lower Respiratory Diseases	34
T5. Accidents	21
T5. Diabetes	21
7. Alzheimer's Disease	15
8. Influenza and Pneumonia	11
9. Suicide	9
T10. Nephritis, Nephrotic Syndrome, and Nephrosis	6
T10. Parkinson's Disease	6
Percent of Deaths due to tobacco use	18.1
Median age at death	81

●Denotes a health status indicator which is significantly lower than the state average.


○Denotes a health status indicator which is significantly higher than the state average.

<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.  
 Source: South Dakota Department of Health, Office of Health Statistics

# Codington County

## Demographic Information



Codington County is located in eastern South Dakota and averages 39.5 persons per square mile.

### 2012 Population Information

Subject	Number	Percent
Total population	27,606	100.0
White	26,308	95.3
Black or African American	203	0.7
American Indian & Alaska Native	564	2.0
Asian	175	0.6
Native Hawaiian & Other Pacific Islander	1	0.0
Two or More Races	355	1.3
Under 5 years	1,994	7.2
Under 18 years	6,736	24.4
65 years and over	4,142	15.0

Source: United States Census Bureau, 2012 Population Estimates

## Health Status Indicators 2008-2012

### Nativity

● Percent of Low Birth Weight Infants	5.0
○ Percent of Mothers Receiving Care in 1st Trimester	77.9
○ Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	23.8
● Percent of Births Less Than 37 Wks. of Gestation	6.6
Average Age of Mother	26.9
● Teenage Birth Rate <sup>2</sup>	10.7
Percent White Births	92.9
Percent American Indian Births	3.9
Percent Unmarried	37.1
Percent WIC births	39.8
Percent Breastfeeding at discharge	74.6
○ Percent Payment-Private Insurance	64.4
● Percent Payment-Medicaid	31.7
● Percent C-Section	22.6

●Denotes a health status indicator which is significantly lower than the state average.  
○Denotes a health status indicator which is significantly higher than the state average.

\*Due to lack of comparable data from all 50 states, significant differences cannot be calculated.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

All Causes	683.5
Heart Disease	150.9
Malignant Neoplasms (cancer)	175.7
Trachea, Bronchus, & Lung	52.5
Colon, Rectum, & Anus	16.9
Female Breast	18.8
Pancreas	11.0
Cerebrovascular Disease	40.5
Chronic Lower Respiratory Diseases	46.2
Accidents	39.2
Motor Vehicle Accidents	10.1
Alzheimer's Disease	36.5
Diabetes Mellitus	22.8
Influenza & Pneumonia	18.7
Intentional Self-Harm (suicide)	20.2
● Chronic Liver Disease and Cirrhosis	5.8
Infant Mortality	5.26

### Leading Causes of Death

	Total Deaths
1. Malignant Neoplasms (cancer)	291
2. Heart Disease	277
3. Chronic Lower Respiratory Diseases	81
4. Cerebrovascular Diseases	73
5. Alzheimer's Disease	69
6. Accidents	59
7. Diabetes	38
8. Influenza and Pneumonia	34
9. Suicide	25
10. Vascular Dementia	16
Percent of Deaths due to tobacco use	23.2
Median age at death	81

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Corson County

## Demographic Information



Corson County is located in the north central portion of the state and averages 1.6 persons per square mile.

### 2012 Population Information

Subject	Number	Percent
Total population	4,077	100.0
White	1,262	31.0
Black or African American	9	0.2
American Indian & Alaska Native	2,683	65.8
Asian	11	0.3
Native Hawaiian & Other Pacific Islander	2	0.0
Two or More Races	110	2.7
Under 5 years	357	8.8
Under 18 years	1,384	33.9
65 years and over	445	10.9

Source: United States Census Bureau, 2012 Population Estimates

## Health Status Indicators 2008-2012

### Nativity

Percent of Low Birth Weight Infants	5.8
● Percent of Mothers Receiving Care in 1st Trimester	47.6
○ Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	22.5
○ Percent of Births Less Than 37 Wks. of Gestation	12.0
● Average Age of Mother	25.3
○ Teenage Birth Rate <sup>2</sup>	39.7
Percent White Births	23.5
Percent American Indian Births	74.3
○ Percent Unmarried	69.3
○ Percent WIC births	76.7
● Percent Breastfeeding at discharge	55.3
● Percent Payment-Private Insurance	17.4
○ Percent Payment-Medicaid	75.2
Percent C-Section	26.0

●Denotes a health status indicator which is significantly lower than the state average.  
○Denotes a health status indicator which is significantly higher than the state average.

\*Due to lack of comparable data from all 50 states, significant differences cannot be calculated.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

○ All Causes	1,215.7
○ Heart Disease	244.4
Malignant Neoplasms (cancer)	227.3
Trachea, Bronchus, & Lung	53.7
Colon, Rectum, & Anus	24.2
Female Breast	65.9
Pancreas	16.1
Cerebrovascular Disease	26.3
Chronic Lower Respiratory Diseases	61.2
○ Accidents	158.3
○ Motor Vehicle Accidents	65.5
Alzheimer's Disease	34.7
○ Diabetes Mellitus	87.5
Influenza & Pneumonia	25.6
○ Intentional Self-Harm (suicide)	77.3
Chronic Liver Disease and Cirrhosis	25.4
Infant Mortality	19.42

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	42
2. Malignant Neoplasms (cancer)	40
3. Accidents	31
4. Suicide	16
5. Diabetes	15
6. Chronic Lower Respiratory Diseases	10
T7. Chronic Liver Disease and Cirrhosis	5
T7. Alzheimer's Disease	5
Percent of Deaths due to tobacco use	18.6
Median age at death	63

●Denotes a health status indicator which is significantly lower than the state average.


○Denotes a health status indicator which is significantly higher than the state average.

<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.


Source: South Dakota Department of Health, Office of Health Statistics

# Custer County

Demographic Information	Health Status Indicators 2008-2012																																																																																																																					
<div style="text-align: center;">  </div> <p>Custer County is located in the southern Black Hills and averages 5.3 persons per square mile.</p> <p style="text-align: center;"><b>2012 Population Information</b></p> <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="text-align: left;">Subject</th> <th style="text-align: right;">Number</th> <th style="text-align: right;">Percent</th> </tr> </thead> <tbody> <tr> <td>Total population</td> <td style="text-align: right;">8,339</td> <td style="text-align: right;">100.0</td> </tr> <tr> <td>  White</td> <td style="text-align: right;">7,806</td> <td style="text-align: right;">93.6</td> </tr> <tr> <td>  Black or African American</td> <td style="text-align: right;">46</td> <td style="text-align: right;">0.6</td> </tr> <tr> <td>  American Indian &amp; Alaska Native</td> <td style="text-align: right;">273</td> <td style="text-align: right;">3.3</td> </tr> <tr> <td>  Asian</td> <td style="text-align: right;">40</td> <td style="text-align: right;">0.5</td> </tr> <tr> <td>  Native Hawaiian &amp; Other Pacific Islander</td> <td style="text-align: right;">4</td> <td style="text-align: right;">0.0</td> </tr> <tr> <td>  Two or More Races</td> <td style="text-align: right;">170</td> <td style="text-align: right;">2.0</td> </tr> <tr> <td>Under 5 years</td> <td style="text-align: right;">372</td> <td style="text-align: right;">4.5</td> </tr> <tr> <td>Under 18 years</td> <td style="text-align: right;">1,598</td> <td style="text-align: right;">19.2</td> </tr> <tr> <td>65 years and over</td> <td style="text-align: right;">1,935</td> <td style="text-align: right;">23.2</td> </tr> </tbody> </table>	Subject	Number	Percent	Total population	8,339	100.0	White	7,806	93.6	Black or African American	46	0.6	American Indian & Alaska Native	273	3.3	Asian	40	0.5	Native Hawaiian & Other Pacific Islander	4	0.0	Two or More Races	170	2.0	Under 5 years	372	4.5	Under 18 years	1,598	19.2	65 years and over	1,935	23.2	<p style="text-align: center;"><b>Nativity</b></p> <table border="1" style="width: 100%; border-collapse: collapse;"> <tbody> <tr> <td>Percent of Low Birth Weight Infants</td> <td style="text-align: right;">8.2</td> </tr> <tr> <td>Percent of Mothers Receiving Care in 1st Trimester</td> <td style="text-align: right;">70.6</td> </tr> <tr> <td>Percent of Mothers Who Used Tobacco While Pregnant<sup>1</sup></td> <td style="text-align: right;">17.9</td> </tr> <tr> <td>Percent of Births Less Than 37 Wks. of Gestation</td> <td style="text-align: right;">8.4</td> </tr> <tr> <td>Average Age of Mother</td> <td style="text-align: right;">27.2</td> </tr> <tr> <td>Teenage Birth Rate<sup>2</sup></td> <td style="text-align: right;">16.0</td> </tr> <tr> <td>Percent White Births</td> <td style="text-align: right;">89.6</td> </tr> <tr> <td>Percent American Indian Births</td> <td style="text-align: right;">5.7</td> </tr> <tr> <td>  • Percent Unmarried</td> <td style="text-align: right;">30.2</td> </tr> <tr> <td>Percent WIC births</td> <td style="text-align: right;">35.8</td> </tr> <tr> <td>Percent Breastfeeding at discharge</td> <td style="text-align: right;">83.0</td> </tr> <tr> <td>Percent Payment-Private Insurance</td> <td style="text-align: right;">53.8</td> </tr> <tr> <td>Percent Payment-Medicaid</td> <td style="text-align: right;">37.7</td> </tr> <tr> <td>  • Percent C-Section</td> <td style="text-align: right;">17.9</td> </tr> </tbody> </table> <p style="text-align: center;"><b>Mortality<sup>3</sup></b></p> <table border="1" style="width: 100%; border-collapse: collapse;"> <tbody> <tr> <td>All Causes</td> <td style="text-align: right;">639.5</td> </tr> <tr> <td>  • Heart Disease</td> <td style="text-align: right;">121.6</td> </tr> <tr> <td>Malignant Neoplasms (cancer)</td> <td style="text-align: right;">171.9</td> </tr> <tr> <td>  Trachea, Bronchus, &amp; Lung</td> <td style="text-align: right;">37.3</td> </tr> <tr> <td>  Colon, Rectum, &amp; Anus</td> <td style="text-align: right;">17.2</td> </tr> <tr> <td>  Female Breast</td> <td style="text-align: right;">19.2</td> </tr> <tr> <td>  Pancreas</td> <td style="text-align: right;">LNE</td> </tr> <tr> <td>Cerebrovascular Disease</td> <td style="text-align: right;">47.5</td> </tr> <tr> <td>Chronic Lower Respiratory Diseases</td> <td style="text-align: right;">31.5</td> </tr> <tr> <td>Accidents</td> <td style="text-align: right;">54.3</td> </tr> <tr> <td>  Motor Vehicle Accidents</td> <td style="text-align: right;">16.1</td> </tr> <tr> <td>  • Alzheimer's Disease</td> <td style="text-align: right;">19.9</td> </tr> <tr> <td>  • Diabetes Mellitus</td> <td style="text-align: right;">11.0</td> </tr> <tr> <td>Influenza &amp; Pneumonia</td> <td style="text-align: right;">12.7</td> </tr> <tr> <td>Intentional Self-Harm (suicide)</td> <td style="text-align: right;">9.8</td> </tr> <tr> <td>Chronic Liver Disease and Cirrhosis</td> <td style="text-align: right;">8.3</td> </tr> <tr> <td>Infant Mortality</td> <td style="text-align: right;">LNE</td> </tr> </tbody> </table> <p style="text-align: center;"><b>Leading Causes of Death</b></p> <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="text-align: left;">Cause</th> <th style="text-align: right;">Total Deaths</th> </tr> </thead> <tbody> <tr> <td>1. Malignant Neoplasms (cancer)</td> <td style="text-align: right;">110</td> </tr> <tr> <td>2. Heart Disease</td> <td style="text-align: right;">81</td> </tr> <tr> <td>3. Cerebrovascular Diseases</td> <td style="text-align: right;">28</td> </tr> <tr> <td>4. Accidents</td> <td style="text-align: right;">24</td> </tr> <tr> <td>5. Chronic Lower Respiratory Diseases</td> <td style="text-align: right;">20</td> </tr> <tr> <td>6. Alzheimer's Disease</td> <td style="text-align: right;">12</td> </tr> <tr> <td>T7. Disorders of Lipoprotein Metabolism and Other Lipidemias</td> <td style="text-align: right;">8</td> </tr> <tr> <td>T7. Influenza and Pneumonia</td> <td style="text-align: right;">8</td> </tr> <tr> <td>9. Diabetes</td> <td style="text-align: right;">7</td> </tr> <tr> <td>10. Parkinson's Disease</td> <td style="text-align: right;">6</td> </tr> </tbody> </table> <p style="text-align: right;">Percent of Deaths due to tobacco use      20.9 Median age at death      78</p>	Percent of Low Birth Weight Infants	8.2	Percent of Mothers Receiving Care in 1st Trimester	70.6	Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	17.9	Percent of Births Less Than 37 Wks. of Gestation	8.4	Average Age of Mother	27.2	Teenage Birth Rate <sup>2</sup>	16.0	Percent White Births	89.6	Percent American Indian Births	5.7	• Percent Unmarried	30.2	Percent WIC births	35.8	Percent Breastfeeding at discharge	83.0	Percent Payment-Private Insurance	53.8	Percent Payment-Medicaid	37.7	• Percent C-Section	17.9	All Causes	639.5	• Heart Disease	121.6	Malignant Neoplasms (cancer)	171.9	Trachea, Bronchus, & Lung	37.3	Colon, Rectum, & Anus	17.2	Female Breast	19.2	Pancreas	LNE	Cerebrovascular Disease	47.5	Chronic Lower Respiratory Diseases	31.5	Accidents	54.3	Motor Vehicle Accidents	16.1	• Alzheimer's Disease	19.9	• Diabetes Mellitus	11.0	Influenza & Pneumonia	12.7	Intentional Self-Harm (suicide)	9.8	Chronic Liver Disease and Cirrhosis	8.3	Infant Mortality	LNE	Cause	Total Deaths	1. Malignant Neoplasms (cancer)	110	2. Heart Disease	81	3. Cerebrovascular Diseases	28	4. Accidents	24	5. Chronic Lower Respiratory Diseases	20	6. Alzheimer's Disease	12	T7. Disorders of Lipoprotein Metabolism and Other Lipidemias	8	T7. Influenza and Pneumonia	8	9. Diabetes	7	10. Parkinson's Disease	6
Subject	Number	Percent																																																																																																																				
Total population	8,339	100.0																																																																																																																				
White	7,806	93.6																																																																																																																				
Black or African American	46	0.6																																																																																																																				
American Indian & Alaska Native	273	3.3																																																																																																																				
Asian	40	0.5																																																																																																																				
Native Hawaiian & Other Pacific Islander	4	0.0																																																																																																																				
Two or More Races	170	2.0																																																																																																																				
Under 5 years	372	4.5																																																																																																																				
Under 18 years	1,598	19.2																																																																																																																				
65 years and over	1,935	23.2																																																																																																																				
Percent of Low Birth Weight Infants	8.2																																																																																																																					
Percent of Mothers Receiving Care in 1st Trimester	70.6																																																																																																																					
Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	17.9																																																																																																																					
Percent of Births Less Than 37 Wks. of Gestation	8.4																																																																																																																					
Average Age of Mother	27.2																																																																																																																					
Teenage Birth Rate <sup>2</sup>	16.0																																																																																																																					
Percent White Births	89.6																																																																																																																					
Percent American Indian Births	5.7																																																																																																																					
• Percent Unmarried	30.2																																																																																																																					
Percent WIC births	35.8																																																																																																																					
Percent Breastfeeding at discharge	83.0																																																																																																																					
Percent Payment-Private Insurance	53.8																																																																																																																					
Percent Payment-Medicaid	37.7																																																																																																																					
• Percent C-Section	17.9																																																																																																																					
All Causes	639.5																																																																																																																					
• Heart Disease	121.6																																																																																																																					
Malignant Neoplasms (cancer)	171.9																																																																																																																					
Trachea, Bronchus, & Lung	37.3																																																																																																																					
Colon, Rectum, & Anus	17.2																																																																																																																					
Female Breast	19.2																																																																																																																					
Pancreas	LNE																																																																																																																					
Cerebrovascular Disease	47.5																																																																																																																					
Chronic Lower Respiratory Diseases	31.5																																																																																																																					
Accidents	54.3																																																																																																																					
Motor Vehicle Accidents	16.1																																																																																																																					
• Alzheimer's Disease	19.9																																																																																																																					
• Diabetes Mellitus	11.0																																																																																																																					
Influenza & Pneumonia	12.7																																																																																																																					
Intentional Self-Harm (suicide)	9.8																																																																																																																					
Chronic Liver Disease and Cirrhosis	8.3																																																																																																																					
Infant Mortality	LNE																																																																																																																					
Cause	Total Deaths																																																																																																																					
1. Malignant Neoplasms (cancer)	110																																																																																																																					
2. Heart Disease	81																																																																																																																					
3. Cerebrovascular Diseases	28																																																																																																																					
4. Accidents	24																																																																																																																					
5. Chronic Lower Respiratory Diseases	20																																																																																																																					
6. Alzheimer's Disease	12																																																																																																																					
T7. Disorders of Lipoprotein Metabolism and Other Lipidemias	8																																																																																																																					
T7. Influenza and Pneumonia	8																																																																																																																					
9. Diabetes	7																																																																																																																					
10. Parkinson's Disease	6																																																																																																																					
<p>Source: United States Census Bureau, 2012 Population Estimates</p>	<p>•Denotes a health status indicator which is significantly lower than the state average. ○Denotes a health status indicator which is significantly higher than the state average. *Due to lack of comparable data from all 50 states, significant differences cannot be calculated. <sup>1</sup>Data for mothers who used tobacco are self-reported. <sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.</p>																																																																																																																					
	<p>•Denotes a health status indicator which is significantly lower than the state average. ○Denotes a health status indicator which is significantly higher than the state average. <sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.</p> <p>See technical notes for more information. Source: South Dakota Department of Health, Office of Health Statistics</p>																																																																																																																					

# Davison County

## Demographic Information



Davison County is located in eastern South Dakota and averages 44.8 persons per square mile.

### 2012 Population Information

Subject	Number	Percent
Total population	19,769	100.0
White	18,646	94.3
Black or African American	172	0.9
American Indian & Alaska Native	558	2.8
Asian	104	0.5
Native Hawaiian & Other Pacific Islander	19	0.1
Two or More Races	270	1.4
Under 5 years	1,303	6.6
Under 18 years	4,563	23.1
65 years and over	3,380	17.1

Source: United States Census Bureau, 2012 Population Estimates

## Health Status Indicators 2008-2012

### Natality

Percent of Low Birth Weight Infants	5.7
○ Percent of Mothers Receiving Care in 1st Trimester	76.4
○ Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	20.6
Percent of Births Less Than 37 Wks. of Gestation	8.3
● Average Age of Mother	26.4
Teenage Birth Rate <sup>2</sup>	13.1
Percent White Births	90.3
Percent American Indian Births	7.0
Percent Unmarried	40.0
○ Percent WIC births	43.9
● Percent Breastfeeding at discharge	68.2
○ Percent Payment-Private Insurance	60.9
Percent Payment-Medicaid	36.3
Percent C-Section	28.7

### Mortality<sup>3</sup>

All Causes	670.8
Heart Disease	179.5
Malignant Neoplasms (cancer)	157.7
Trachea, Bronchus, & Lung	51.0
Colon, Rectum, & Anus	12.2
Female Breast	21.8
Pancreas	8.6
Cerebrovascular Disease	42.9
Chronic Lower Respiratory Diseases	42.8
● Accidents	28.5
● Motor Vehicle Accidents	7.9
● Alzheimer's Disease	25.7
● Diabetes Mellitus	12.2
Influenza & Pneumonia	20.2
Intentional Self-Harm (suicide)	19.1
Chronic Liver Disease and Cirrhosis	9.3
Infant Mortality	6.64

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	266
2. Malignant Neoplasms (cancer)	209
3. Cerebrovascular Diseases	67
4. Chronic Lower Respiratory Diseases	62
5. Alzheimer's Disease	46
6. Influenza and Pneumonia	40
7. Accidents	35
8. Unspecified Dementia	20
9. Pneumonitis Due to Solids and Liquids	19
10. Suicide	17
Percent of Deaths due to tobacco use	18.3
Median age at death	82

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

\*Due to lack of comparable data from all 50 states, significant differences cannot be calculated.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

●Denotes a health status indicator which is significantly lower than the state average.


○Denotes a health status indicator which is significantly higher than the state average.

<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.


Source: South Dakota Department of Health, Office of Health Statistics

# Day County

Demographic Information	Health Status Indicators 2008-2012																																																																																																																								
<div style="text-align: center;">  </div> <p>Day County is located in the northeastern part of the state and averages 5.6 persons per square mile.</p> <p style="text-align: center;"><b>2012 Population Information</b></p> <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="text-align: left;">Subject</th> <th style="text-align: right;">Number</th> <th style="text-align: right;">Percent</th> </tr> </thead> <tbody> <tr> <td>Total population</td> <td style="text-align: right;">5,613</td> <td style="text-align: right;">100.0</td> </tr> <tr> <td>White</td> <td style="text-align: right;">4,973</td> <td style="text-align: right;">88.6</td> </tr> <tr> <td>Black or African American</td> <td style="text-align: right;">29</td> <td style="text-align: right;">0.5</td> </tr> <tr> <td>American Indian &amp; Alaska Native</td> <td style="text-align: right;">499</td> <td style="text-align: right;">8.9</td> </tr> <tr> <td>Asian</td> <td style="text-align: right;">16</td> <td style="text-align: right;">0.3</td> </tr> <tr> <td>Native Hawaiian &amp; Other Pacific Islander</td> <td style="text-align: right;">0</td> <td style="text-align: right;">0.0</td> </tr> <tr> <td>Two or More Races</td> <td style="text-align: right;">96</td> <td style="text-align: right;">1.7</td> </tr> <tr> <td>Under 5 years</td> <td style="text-align: right;">328</td> <td style="text-align: right;">5.8</td> </tr> <tr> <td>Under 18 years</td> <td style="text-align: right;">1,198</td> <td style="text-align: right;">21.3</td> </tr> <tr> <td>65 years and over</td> <td style="text-align: right;">1,332</td> <td style="text-align: right;">23.7</td> </tr> </tbody> </table>	Subject	Number	Percent	Total population	5,613	100.0	White	4,973	88.6	Black or African American	29	0.5	American Indian & Alaska Native	499	8.9	Asian	16	0.3	Native Hawaiian & Other Pacific Islander	0	0.0	Two or More Races	96	1.7	Under 5 years	328	5.8	Under 18 years	1,198	21.3	65 years and over	1,332	23.7	<p style="text-align: center;"><b>Nativity</b></p> <table border="1" style="width: 100%; border-collapse: collapse;"> <tbody> <tr> <td>Percent of Low Birth Weight Infants</td> <td style="text-align: right;">6.4</td> </tr> <tr> <td>Percent of Mothers Receiving Care in 1st Trimester</td> <td style="text-align: right;">67.5</td> </tr> <tr> <td>○ Percent of Mothers Who Used Tobacco While Pregnant<sup>1</sup></td> <td style="text-align: right;">30.8</td> </tr> <tr> <td>Percent of Births Less Than 37 Wks. of Gestation</td> <td style="text-align: right;">7.6</td> </tr> <tr> <td>Average Age of Mother</td> <td style="text-align: right;">27.3</td> </tr> <tr> <td>● Teenage Birth Rate<sup>2</sup></td> <td style="text-align: right;">5.5</td> </tr> <tr> <td>Percent White Births</td> <td style="text-align: right;">72.7</td> </tr> <tr> <td>Percent American Indian Births</td> <td style="text-align: right;">21.5</td> </tr> <tr> <td>Percent Unmarried</td> <td style="text-align: right;">40.5</td> </tr> <tr> <td>Percent WIC births</td> <td style="text-align: right;">41.1</td> </tr> <tr> <td>Percent Breastfeeding at discharge</td> <td style="text-align: right;">70.9</td> </tr> <tr> <td>Percent Payment-Private Insurance</td> <td style="text-align: right;">56.9</td> </tr> <tr> <td>Percent Payment-Medicaid</td> <td style="text-align: right;">37.0</td> </tr> <tr> <td>Percent C-Section</td> <td style="text-align: right;">23.2</td> </tr> </tbody> </table>	Percent of Low Birth Weight Infants	6.4	Percent of Mothers Receiving Care in 1st Trimester	67.5	○ Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	30.8	Percent of Births Less Than 37 Wks. of Gestation	7.6	Average Age of Mother	27.3	● Teenage Birth Rate <sup>2</sup>	5.5	Percent White Births	72.7	Percent American Indian Births	21.5	Percent Unmarried	40.5	Percent WIC births	41.1	Percent Breastfeeding at discharge	70.9	Percent Payment-Private Insurance	56.9	Percent Payment-Medicaid	37.0	Percent C-Section	23.2	<p style="text-align: center;"><b>Mortality<sup>3</sup></b></p> <table border="1" style="width: 100%; border-collapse: collapse;"> <tbody> <tr> <td>All Causes</td> <td style="text-align: right;">720.0</td> </tr> <tr> <td>Heart Disease</td> <td style="text-align: right;">195.7</td> </tr> <tr> <td>Malignant Neoplasms (cancer)</td> <td style="text-align: right;">162.7</td> </tr> <tr> <td>    Trachea, Bronchus, &amp; Lung</td> <td style="text-align: right;">43.4</td> </tr> <tr> <td>    Colon, Rectum, &amp; Anus</td> <td style="text-align: right;">12.3</td> </tr> <tr> <td>    Female Breast</td> <td style="text-align: right;">21.6</td> </tr> <tr> <td>    Pancreas</td> <td style="text-align: right;">13.4</td> </tr> <tr> <td>Cerebrovascular Disease</td> <td style="text-align: right;">53.6</td> </tr> <tr> <td>Chronic Lower Respiratory Diseases</td> <td style="text-align: right;">43.1</td> </tr> <tr> <td>Accidents</td> <td style="text-align: right;">56.9</td> </tr> <tr> <td>    Motor Vehicle Accidents</td> <td style="text-align: right;">14.9</td> </tr> <tr> <td>Alzheimer's Disease</td> <td style="text-align: right;">37.0</td> </tr> <tr> <td>Diabetes Mellitus</td> <td style="text-align: right;">14.5</td> </tr> <tr> <td>Influenza &amp; Pneumonia</td> <td style="text-align: right;">16.2</td> </tr> <tr> <td>Intentional Self-Harm (suicide)</td> <td style="text-align: right;">LNE</td> </tr> <tr> <td>Chronic Liver Disease and Cirrhosis</td> <td style="text-align: right;">8.9</td> </tr> <tr> <td>Infant Mortality</td> <td style="text-align: right;">LNE</td> </tr> </tbody> </table> <p style="text-align: center;"><b>Leading Causes of Death</b></p> <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="text-align: left;">Leading Causes of Death</th> <th style="text-align: right;">Total Deaths</th> </tr> </thead> <tbody> <tr> <td>1. Heart Disease</td> <td style="text-align: right;">109</td> </tr> <tr> <td>2. Malignant Neoplasms (cancer)</td> <td style="text-align: right;">86</td> </tr> <tr> <td>3. Cerebrovascular Diseases</td> <td style="text-align: right;">31</td> </tr> <tr> <td>4. Alzheimer's Disease</td> <td style="text-align: right;">25</td> </tr> <tr> <td>T5. Accidents</td> <td style="text-align: right;">24</td> </tr> <tr> <td>T5. Chronic Lower Respiratory Diseases</td> <td style="text-align: right;">24</td> </tr> <tr> <td>7. Influenza and Pneumonia</td> <td style="text-align: right;">10</td> </tr> <tr> <td>8. Diabetes</td> <td style="text-align: right;">7</td> </tr> <tr> <td>9. Nephritis, Nephrotic Syndrome, and Nephrosis</td> <td style="text-align: right;">5</td> </tr> </tbody> </table> <table border="1" style="width: 100%; border-collapse: collapse; margin-top: 10px;"> <tbody> <tr> <td>Percent of Deaths due to tobacco use</td> <td style="text-align: right;">18.2</td> </tr> <tr> <td>Median age at death</td> <td style="text-align: right;">83</td> </tr> </tbody> </table> <p>●Denotes a health status indicator which is significantly lower than the state average.  ○Denotes a health status indicator which is significantly higher than the state average.  <sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.</p> <p>See technical notes for more information.  Source: South Dakota Department of Health, Office of Health Statistics</p>	All Causes	720.0	Heart Disease	195.7	Malignant Neoplasms (cancer)	162.7	Trachea, Bronchus, & Lung	43.4	Colon, Rectum, & Anus	12.3	Female Breast	21.6	Pancreas	13.4	Cerebrovascular Disease	53.6	Chronic Lower Respiratory Diseases	43.1	Accidents	56.9	Motor Vehicle Accidents	14.9	Alzheimer's Disease	37.0	Diabetes Mellitus	14.5	Influenza & Pneumonia	16.2	Intentional Self-Harm (suicide)	LNE	Chronic Liver Disease and Cirrhosis	8.9	Infant Mortality	LNE	Leading Causes of Death	Total Deaths	1. Heart Disease	109	2. Malignant Neoplasms (cancer)	86	3. Cerebrovascular Diseases	31	4. Alzheimer's Disease	25	T5. Accidents	24	T5. Chronic Lower Respiratory Diseases	24	7. Influenza and Pneumonia	10	8. Diabetes	7	9. Nephritis, Nephrotic Syndrome, and Nephrosis	5	Percent of Deaths due to tobacco use	18.2	Median age at death	83
Subject	Number	Percent																																																																																																																							
Total population	5,613	100.0																																																																																																																							
White	4,973	88.6																																																																																																																							
Black or African American	29	0.5																																																																																																																							
American Indian & Alaska Native	499	8.9																																																																																																																							
Asian	16	0.3																																																																																																																							
Native Hawaiian & Other Pacific Islander	0	0.0																																																																																																																							
Two or More Races	96	1.7																																																																																																																							
Under 5 years	328	5.8																																																																																																																							
Under 18 years	1,198	21.3																																																																																																																							
65 years and over	1,332	23.7																																																																																																																							
Percent of Low Birth Weight Infants	6.4																																																																																																																								
Percent of Mothers Receiving Care in 1st Trimester	67.5																																																																																																																								
○ Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	30.8																																																																																																																								
Percent of Births Less Than 37 Wks. of Gestation	7.6																																																																																																																								
Average Age of Mother	27.3																																																																																																																								
● Teenage Birth Rate <sup>2</sup>	5.5																																																																																																																								
Percent White Births	72.7																																																																																																																								
Percent American Indian Births	21.5																																																																																																																								
Percent Unmarried	40.5																																																																																																																								
Percent WIC births	41.1																																																																																																																								
Percent Breastfeeding at discharge	70.9																																																																																																																								
Percent Payment-Private Insurance	56.9																																																																																																																								
Percent Payment-Medicaid	37.0																																																																																																																								
Percent C-Section	23.2																																																																																																																								
All Causes	720.0																																																																																																																								
Heart Disease	195.7																																																																																																																								
Malignant Neoplasms (cancer)	162.7																																																																																																																								
Trachea, Bronchus, & Lung	43.4																																																																																																																								
Colon, Rectum, & Anus	12.3																																																																																																																								
Female Breast	21.6																																																																																																																								
Pancreas	13.4																																																																																																																								
Cerebrovascular Disease	53.6																																																																																																																								
Chronic Lower Respiratory Diseases	43.1																																																																																																																								
Accidents	56.9																																																																																																																								
Motor Vehicle Accidents	14.9																																																																																																																								
Alzheimer's Disease	37.0																																																																																																																								
Diabetes Mellitus	14.5																																																																																																																								
Influenza & Pneumonia	16.2																																																																																																																								
Intentional Self-Harm (suicide)	LNE																																																																																																																								
Chronic Liver Disease and Cirrhosis	8.9																																																																																																																								
Infant Mortality	LNE																																																																																																																								
Leading Causes of Death	Total Deaths																																																																																																																								
1. Heart Disease	109																																																																																																																								
2. Malignant Neoplasms (cancer)	86																																																																																																																								
3. Cerebrovascular Diseases	31																																																																																																																								
4. Alzheimer's Disease	25																																																																																																																								
T5. Accidents	24																																																																																																																								
T5. Chronic Lower Respiratory Diseases	24																																																																																																																								
7. Influenza and Pneumonia	10																																																																																																																								
8. Diabetes	7																																																																																																																								
9. Nephritis, Nephrotic Syndrome, and Nephrosis	5																																																																																																																								
Percent of Deaths due to tobacco use	18.2																																																																																																																								
Median age at death	83																																																																																																																								
<p>Source: United States Census Bureau, 2012 Population Estimates</p>	<p>●Denotes a health status indicator which is significantly lower than the state average.  ○Denotes a health status indicator which is significantly higher than the state average.  *Due to lack of comparable data from all 50 states, significant differences cannot be calculated.  <sup>1</sup>Data for mothers who used tobacco are self-reported.  <sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.</p>																																																																																																																								

# Deuel County

## Demographic Information



Deuel County borders Minnesota and averages 7.0 persons per square mile.

## 2012 Population Information

Subject	Number	Percent
Total population	4,380	100.0
White	4,282	97.8
Black or African American	40	0.9
American Indian & Alaska Native	17	0.4
Asian	7	0.2
Native Hawaiian & Other Pacific Islander	0	0.0
Two or More Races	34	0.8
Under 5 years	257	5.9
Under 18 years	1,006	23.0
65 years and over	877	20.0

Source: United States Census Bureau, 2012 Population Estimates

## Health Status Indicators 2008-2012

### Nativity

Percent of Low Birth Weight Infants	5.2
○ Percent of Mothers Receiving Care in 1st Trimester	83.8
Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	17.6
Percent of Births Less Than 37 Wks. of Gestation	7.1
Average Age of Mother	27.3
Teenage Birth Rate <sup>2</sup>	10.6
Percent White Births	99.2
Percent American Indian Births	LNE
● Percent Unmarried	24.9
● Percent WIC births	30.0
Percent Breastfeeding at discharge	79.7
○ Percent Payment-Private Insurance	78.8
● Percent Payment-Medicaid	15.1
Percent C-Section	22.5

●Denotes a health status indicator which is significantly lower than the state average.  
○Denotes a health status indicator which is significantly higher than the state average.

\*Due to lack of comparable data from all 50 states, significant differences cannot be calculated.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

All Causes	661.5
Heart Disease	169.1
● Malignant Neoplasms (cancer)	114.7
Trachea, Bronchus, & Lung	55.1
Colon, Rectum, & Anus	11.2
Female Breast	16.3
Pancreas	LNE
Cerebrovascular Disease	29.7
Chronic Lower Respiratory Diseases	66.8
Accidents	59.1
Motor Vehicle Accidents	16.4
Alzheimer's Disease	22.2
Diabetes Mellitus	17.6
Influenza & Pneumonia	13.6
Intentional Self-Harm (suicide)	22.1
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	LNE

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	61
2. Malignant Neoplasms (cancer)	39
3. Chronic Lower Respiratory Diseases	23
4. Accidents	14
5. Septicemia	12
6. Cerebrovascular Diseases	11
7. Alzheimer's Disease	9
T8. Influenza and Pneumonia	6
T8. Diabetes	6
Percent of Deaths due to tobacco use	21.3
Median age at death	81

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Dewey County

## Demographic Information



Dewey County is located in the north central region of the state and averages 2.3 persons per square mile.

### 2012 Population Information

Subject	Number	Percent
Total population	5,538	100.0
White	1,225	22.1
Black or African American	10	0.2
American Indian & Alaska Native	4,088	73.8
Asian	9	0.2
Native Hawaiian & Other Pacific Islander	0	0.0
Two or More Races	206	3.7
Under 5 years	566	10.2
Under 18 years	1,917	34.6
65 years and over	545	9.8

Source: United States Census Bureau, 2012 Population Estimates

## Health Status Indicators 2008-2012

### Natality

Percent of Low Birth Weight Infants	6.0
● Percent of Mothers Receiving Care in 1st Trimester	51.0
○ Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	26.4
Percent of Births Less Than 37 Wks. of Gestation	8.6
● Average Age of Mother	25.2
○ Teenage Birth Rate <sup>2</sup>	41.8
Percent White Births	10.2
Percent American Indian Births	84.6
○ Percent Unmarried	67.7
○ Percent WIC births	83.0
● Percent Breastfeeding at discharge	52.6
● Percent Payment-Private Insurance	16.2
○ Percent Payment-Medicaid	76.8
○ Percent C-Section	34.4

●Denotes a health status indicator which is significantly lower than the state average.  
○Denotes a health status indicator which is significantly higher than the state average.  
\*Due to lack of comparable data from all 50 states, significant differences cannot be calculated.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

○ All Causes	1,196.9
○ Heart Disease	268.4
○ Malignant Neoplasms (cancer)	241.5
○ Trachea, Bronchus, & Lung	97.8
Colon, Rectum, & Anus	28.1
Female Breast	25.9
Pancreas	LNE
Cerebrovascular Disease	29.9
Chronic Lower Respiratory Diseases	55.6
○ Accidents	92.0
Motor Vehicle Accidents	40.1
Alzheimer's Disease	39.8
○ Diabetes Mellitus	76.3
Influenza & Pneumonia	37.6
○ Intentional Self-Harm (suicide)	58.4
○ Chronic Liver Disease and Cirrhosis	57.2
Infant Mortality	5.82

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	58
2. Malignant Neoplasms (cancer)	53
3. Accidents	22
4. Diabetes	18
5. Suicide	16
6. Chronic Liver Disease and Cirrhosis	14
7. Chronic Lower Respiratory Diseases	12
T8. Influenza and Pneumonia	8
T8. Alzheimer's Disease	8
10. Mental and Behavioral Disorders Due to Use of Alcohol	7
Percent of Deaths due to tobacco use	21.5
Median age at death	65

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Douglas County

## Demographic Information



Douglas County is located in south central South Dakota and averages 7.0 persons per square mile.

### 2012 Population Information

Subject	Number	Percent
Total population	2,970	100.0
White	2,857	96.2
Black or African American	12	0.4
American Indian & Alaska Native	67	2.3
Asian	7	0.2
Native Hawaiian & Other Pacific Islander	0	0.0
Two or More Races	27	0.9
Under 5 years	163	5.5
Under 18 years	658	22.2
65 years and over	725	24.4

Source: United States Census Bureau, 2012 Population Estimates

## Health Status Indicators 2008-2012

### Nativity

Percent of Low Birth Weight Infants	4.3
Percent of Mothers Receiving Care in 1st Trimester	69.4
● Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	10.4
● Percent of Births Less Than 37 Wks. of Gestation	3.7
○ Average Age of Mother	28.2
Teenage Birth Rate <sup>2</sup>	8.7
Percent White Births	96.3
Percent American Indian Births	LNE
● Percent Unmarried	13.5
● Percent WIC births	26.5
Percent Breastfeeding at discharge	85.3
Percent Payment-Private Insurance	61.7
● Percent Payment-Medicaid	22.2
Percent C-Section	25.2

●Denotes a health status indicator which is significantly lower than the state average.  
○Denotes a health status indicator which is significantly higher than the state average.

\*Due to lack of comparable data from all 50 states, significant differences cannot be calculated.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

All Causes	672.7
Heart Disease	153.3
Malignant Neoplasms (cancer)	180.1
Trachea, Bronchus, & Lung	26.7
Colon, Rectum, & Anus	19.7
Female Breast	27.8
Pancreas	25.6
Cerebrovascular Disease	46.6
Chronic Lower Respiratory Diseases	40.6
Accidents	25.1
Motor Vehicle Accidents	LNE
Alzheimer's Disease	51.5
Diabetes Mellitus	28.5
Influenza & Pneumonia	15.8
Intentional Self-Harm (suicide)	LNE
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	LNE

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	57
2. Malignant Neoplasms (cancer)	45
3. Alzheimer's Disease	24
4. Cerebrovascular Diseases	19
5. Chronic Lower Respiratory Diseases	14
6. Diabetes	11
7. Accidents	8
8. Influenza and Pneumonia	7
Percent of Deaths due to tobacco use	10.8
Median age at death	85

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Edmunds County

## Demographic Information



Edmunds County is located in the north central region of the state and averages 3.6 persons per square mile.

### 2012 Population Information

Subject	Number	Percent
Total population	4,026	100.0
White	3,949	98.1
Black or African American	8	0.2
American Indian & Alaska Native	20	0.5
Asian	7	0.2
Native Hawaiian & Other Pacific Islander	0	0.0
Two or More Races	42	1.0
Under 5 years	225	5.6
Under 18 years	953	23.7
65 years and over	865	21.5

Source: United States Census Bureau, 2012 Population Estimates

## Health Status Indicators 2008-2012

### Nativity

Percent of Low Birth Weight Infants	3.9
Percent of Mothers Receiving Care in 1st Trimester	70.7
● Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	11.2
Percent of Births Less Than 37 Wks. of Gestation	5.3
○ Average Age of Mother	27.9
● Teenage Birth Rate <sup>2</sup>	8.2
Percent White Births	97.8
Percent American Indian Births	LNE
● Percent Unmarried	18.4
● Percent WIC births	22.6
Percent Breastfeeding at discharge	79.9
○ Percent Payment-Private Insurance	73.2
● Percent Payment-Medicaid	17.0
Percent C-Section	28.1

●Denotes a health status indicator which is significantly lower than the state average.  
 ○Denotes a health status indicator which is significantly higher than the state average.  
 \*Due to lack of comparable data from all 50 states, significant differences cannot be calculated.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

All Causes	676.0
Heart Disease	155.9
Malignant Neoplasms (cancer)	151.5
● Trachea, Bronchus, & Lung	24.3
Colon, Rectum, & Anus	16.6
Female Breast	27.6
Pancreas	11.4
Cerebrovascular Disease	27.3
Chronic Lower Respiratory Diseases	56.1
Accidents	34.1
Motor Vehicle Accidents	23.0
Alzheimer's Disease	34.9
○ Diabetes Mellitus	47.3
Influenza & Pneumonia	17.2
Intentional Self-Harm (suicide)	15.1
Chronic Liver Disease and Cirrhosis	11.8
Infant Mortality	LNE

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	61
2. Malignant Neoplasms (cancer)	51
3. Chronic Lower Respiratory Diseases	21
4. Diabetes	18
5. Alzheimer's Disease	16
T6. Accidents	9
T6. Cerebrovascular Diseases	9
8. Influenza and Pneumonia	7
9. Essential (primary) Hypertension and Hypertensive Renal Disease	6
Percent of Deaths due to tobacco use	14.6
Median age at death	83

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Fall River County

## Demographic Information



Fall River County is located in the southwestern corner of the state and averages 4.1 persons per square mile.

## 2012 Population Information

Subject	Number	Percent
Total population	6,971	100.0
White	6,188	88.8
Black or African American	77	1.1
American Indian & Alaska Native	470	6.7
Asian	43	0.6
Native Hawaiian & Other Pacific Islander	3	0.0
Two or More Races	190	2.7
Under 5 years	236	3.4
Under 18 years	1,187	17.0
65 years and over	1,708	24.5

Source: United States Census Bureau, 2012 Population Estimates

## Health Status Indicators 2008-2012

### Nativity

Percent of Low Birth Weight Infants	7.9
● Percent of Mothers Receiving Care in 1st Trimester	56.7
○ Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	28.3
Percent of Births Less Than 37 Wks. of Gestation	7.9
Average Age of Mother	26.6
Teenage Birth Rate <sup>2</sup>	15.8
Percent White Births	81.9
Percent American Indian Births	10.6
Percent Unmarried	41.0
○ Percent WIC births	53.4
Percent Breastfeeding at discharge	79.3
● Percent Payment-Private Insurance	45.1
○ Percent Payment-Medicaid	48.9
● Percent C-Section	15.0

●Denotes a health status indicator which is significantly lower than the state average.  
○Denotes a health status indicator which is significantly higher than the state average.  
\*Due to lack of comparable data from all 50 states, significant differences cannot be calculated.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

○ All Causes	832.3
Heart Disease	194.5
Malignant Neoplasms (cancer)	184.7
Trachea, Bronchus, & Lung	61.7
● Colon, Rectum, & Anus	8.1
Female Breast	28.8
Pancreas	7.2
Cerebrovascular Disease	31.8
○ Chronic Lower Respiratory Diseases	77.6
Accidents	63.5
Motor Vehicle Accidents	20.1
● Alzheimer's Disease	20.9
○ Diabetes Mellitus	47.5
Influenza & Pneumonia	10.3
Intentional Self-Harm (suicide)	7.8
Chronic Liver Disease and Cirrhosis	21.4
Infant Mortality	LNE

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	132
2. Malignant Neoplasms (cancer)	126
3. Chronic Lower Respiratory Diseases	55
4. Diabetes	31
5. Accidents	26
6. Cerebrovascular Diseases	22
7. Alzheimer's Disease	16
8. Chronic Liver Disease and Cirrhosis	10
T9. Unspecified Dementia	9
T9. Pneumonitis Due to Solids and Liquids	9
T9. Parkinson's Disease	9
T9. Septicemia	9
Percent of Deaths due to tobacco use	20.3
Median age at death	79

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Faulk County

## Demographic Information



Faulk County is located in north central South Dakota and averages 2.4 persons per square mile.

### 2012 Population Information

Subject	Number	Percent
Total population	2,377	100.0
White	2,343	98.6
Black or African American	5	0.2
American Indian & Alaska Native	3	0.1
Asian	7	0.3
Native Hawaiian & Other Pacific Islander	0	0.0
Two or More Races	19	0.8
Under 5 years	141	5.9
Under 18 years	550	23.1
65 years and over	553	23.3

Source: United States Census Bureau, 2012 Population Estimates

## Health Status Indicators 2008-2012

### Natality

● Percent of Low Birth Weight Infants	3.1
Percent of Mothers Receiving Care in 1st Trimester	57.5
● Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	5.5
● Percent of Births Less Than 37 Wks. of Gestation	3.9
○ Average Age of Mother	29.5
Teenage Birth Rate <sup>2</sup>	LNE
Percent White Births	100
Percent American Indian Births	LNE
● Percent Unmarried	12.4
● Percent WIC births	18.0
Percent Breastfeeding at discharge	86.7
○ Percent Payment-Private Insurance	78.7
● Percent Payment-Medicaid	11.8
Percent C-Section	20.2

●Denotes a health status indicator which is significantly lower than the state average.  
○Denotes a health status indicator which is significantly higher than the state average.  
\*Due to lack of comparable data from all 50 states, significant differences cannot be calculated.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

● All Causes	531.2
● Heart Disease	91.3
● Malignant Neoplasms (cancer)	106.6
Trachea, Bronchus, & Lung	25.9
Colon, Rectum, & Anus	10.3
Female Breast	LNE
Pancreas	LNE
Cerebrovascular Disease	23.2
Chronic Lower Respiratory Diseases	44.3
● Accidents	9.3
Motor Vehicle Accidents	LNE
Alzheimer's Disease	56.1
Diabetes Mellitus	27.1
Influenza & Pneumonia	37.5
Intentional Self-Harm (suicide)	LNE
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	LNE

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Malignant Neoplasms (cancer)	25
2. Heart Disease	23
3. Alzheimer's Disease	18
4. Chronic Lower Respiratory Diseases	11
5. Influenza and Pneumonia	8
6. Diabetes	7
7. Cerebrovascular Diseases	6
Percent of Deaths due to tobacco use	9.8
Median age at death	86

●Denotes a health status indicator which is significantly lower than the state average.  
○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Grant County

## Demographic Information



Grant County borders Minnesota in northeastern South Dakota and averages 10.8 persons per square mile.

### 2012 Population Information

Subject	Number	Percent
Total population	7,259	100.0
White	7,073	97.4
Black or African American	27	0.4
American Indian & Alaska Native	59	0.8
Asian	29	0.4
Native Hawaiian & Other Pacific Islander	3	0.0
Two or More Races	68	0.9
Under 5 years	398	5.5
Under 18 years	1,612	22.2
65 years and over	1,425	19.6

Source: United States Census Bureau, 2012 Population Estimates

## Health Status Indicators 2008-2012

### Nativity

Percent of Low Birth Weight Infants	4.2
Percent of Mothers Receiving Care in 1st Trimester	75.4
Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	15.6
• Percent of Births Less Than 37 Wks. of Gestation	4.5
Average Age of Mother	27.4
Teenage Birth Rate <sup>2</sup>	20.2
Percent White Births	96.1
Percent American Indian Births	0.8
• Percent Unmarried	30.5
Percent WIC births	37.0
Percent Breastfeeding at discharge	72.5
Percent Payment-Private Insurance	60.6
• Percent Payment-Medicaid	22.3
Percent C-Section	24.0

•Denotes a health status indicator which is significantly lower than the state average.  
 ○Denotes a health status indicator which is significantly higher than the state average.  
 \*Due to lack of comparable data from all 50 states, significant differences cannot be calculated.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

All Causes	636.4
Heart Disease	149.0
Malignant Neoplasms (cancer)	144.2
• Trachea, Bronchus, & Lung	26.4
Colon, Rectum, & Anus	21.2
Female Breast	19.2
Pancreas	9.0
Cerebrovascular Disease	42.8
Chronic Lower Respiratory Diseases	40.7
Accidents	43.1
Motor Vehicle Accidents	19.8
Alzheimer's Disease	38.1
Diabetes Mellitus	24.8
• Influenza & Pneumonia	6.0
Intentional Self-Harm (suicide)	LNE
Chronic Liver Disease and Cirrhosis	6.1
Infant Mortality	LNE

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	89
2. Malignant Neoplasms (cancer)	81
3. Cerebrovascular Diseases	29
4. Alzheimer's Disease	28
5. Chronic Lower Respiratory Diseases	25
6. Accidents	15
7. Diabetes	14
8. Aortic Aneurysm and Dissection	6
9. Disorders of Lipoprotein Metabolism and Other Lipidemias	5
Percent of Deaths due to tobacco use	15.3
Median age at death	82

•Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Gregory County

## Demographic Information



Gregory County borders the west bank of the Missouri River and the state of Nebraska and averages 4.2 persons per square mile.

### 2012 Population Information

Subject	Number	Percent
Total population	4,265	100.0
White	3,823	89.6
Black or African American	12	0.3
American Indian & Alaska Native	316	7.4
Asian	17	0.4
Native Hawaiian & Other Pacific Islander	7	0.2
Two or More Races	90	2.1
Under 5 years	250	5.9
Under 18 years	952	22.3
65 years and over	1,025	24.0

Source: United States Census Bureau, 2012 Population Estimates

## Health Status Indicators 2008-2012

### Natality

Percent of Low Birth Weight Infants	4.6
Percent of Mothers Receiving Care in 1st Trimester	61.6
Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	19.8
Percent of Births Less Than 37 Wks. of Gestation	9.1
Average Age of Mother	27.2
Teenage Birth Rate <sup>2</sup>	11.9
Percent White Births	79.9
Percent American Indian Births	16.0
Percent Unmarried	37.0
○ Percent WIC births	49.8
Percent Breastfeeding at discharge	69.9
Percent Payment-Private Insurance	48.9
Percent Payment-Medicaid	43.8
Percent C-Section	31.5

●Denotes a health status indicator which is significantly lower than the state average.  
○Denotes a health status indicator which is significantly higher than the state average.  
\*Due to lack of comparable data from all 50 states, significant differences cannot be calculated.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

○ All Causes	809.0
○ Heart Disease	233.9
Malignant Neoplasms (cancer)	186.1
Trachea, Bronchus, & Lung	57.8
Colon, Rectum, & Anus	19.1
Female Breast	12.1
Pancreas	LNE
Cerebrovascular Disease	36.0
Chronic Lower Respiratory Diseases	57.5
Accidents	60.1
Motor Vehicle Accidents	24.6
Alzheimer's Disease	22.9
○ Diabetes Mellitus	47.6
● Influenza & Pneumonia	5.8
Intentional Self-Harm (suicide)	15.5
Chronic Liver Disease and Cirrhosis	12.9
Infant Mortality	LNE

### Leading Causes of Death

	Total Deaths
1. Heart Disease	112
2. Malignant Neoplasms (cancer)	75
3. Chronic Lower Respiratory Diseases	26
4. Diabetes	21
5. Accidents	18
T6. Disorders of Lipoprotein Metabolism and Other Lipidemias	14
T6. Cerebrovascular Diseases	14
8. Alzheimer's Disease	13
Percent of Deaths due to tobacco use	20.6
Median age at death	84

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Haakon County

## Demographic Information



Haakon County is located in the west central region of the state and averages 1.1 persons per square mile.

### 2012 Population Information

Subject	Number	Percent
Total population	1,939	100.0
White	1,808	93.2
Black or African American	15	0.8
American Indian & Alaska Native	40	2.1
Asian	20	1.0
Native Hawaiian & Other Pacific Islander	2	0.1
Two or More Races	54	2.8
Under 5 years	125	6.4
Under 18 years	442	22.8
65 years and over	422	21.8

Source: United States Census Bureau, 2012 Population Estimates

## Health Status Indicators 2008-2012

### Nativity

● Percent of Low Birth Weight Infants	2.7
Percent of Mothers Receiving Care in 1st Trimester	69.4
● Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	10.9
Percent of Births Less Than 37 Wks. of Gestation	6.3
Average Age of Mother	27.8
Teenage Birth Rate <sup>2</sup>	21.1
Percent White Births	91.9
Percent American Indian Births	2.7
● Percent Unmarried	23.4
● Percent WIC births	23.9
Percent Breastfeeding at discharge	90.0
○ Percent Payment-Private Insurance	78.4
● Percent Payment-Medicaid	15.3
Percent C-Section	27.0

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

\*Due to lack of comparable data from all 50 states, significant differences cannot be calculated.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

All Causes	641.2
Heart Disease	136.6
Malignant Neoplasms (cancer)	163.8
Trachea, Bronchus, & Lung	35.0
Colon, Rectum, & Anus	29.2
Female Breast	LNE
Pancreas	LNE
Cerebrovascular Disease	38.3
Chronic Lower Respiratory Diseases	30.7
Accidents	44.5
Motor Vehicle Accidents	36.7
Alzheimer's Disease	LNE
Diabetes Mellitus	LNE
○ Influenza & Pneumonia	53.9
Intentional Self-Harm (suicide)	LNE
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	LNE

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	27
2. Malignant Neoplasms (cancer)	26
3. Influenza and Pneumonia	12
T4. Accidents	6
T4. Chronic Lower Respiratory Diseases	6
T4. Cerebrovascular Diseases	6
Percent of Deaths due to tobacco use	11.5
Median age at death	85

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Hamlin County

## Demographic Information



Hamlin County is located in the northeastern region of the state and averages 11.6 persons per square mile.

### 2012 Population Information

Subject	Number	Percent
Total population	5,918	100.0
White	5,810	98.2
Black or African American	32	0.5
American Indian & Alaska Native	23	0.4
Asian	14	0.2
Native Hawaiian & Other Pacific Islander	1	0.0
Two or More Races	38	0.6
Under 5 years	531	9.0
Under 18 years	1,770	29.9
65 years and over	1,010	17.1

Source: United States Census Bureau, 2012 Population Estimates

## Health Status Indicators 2008-2012

### Natality

● Percent of Low Birth Weight Infants	4.0
○ Percent of Mothers Receiving Care in 1st Trimester	68.1
● Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	14.8
● Percent of Births Less Than 37 Wks. of Gestation	5.3
Average Age of Mother	27.4
Teenage Birth Rate <sup>2</sup>	16.1
Percent White Births	99.1
Percent American Indian Births	LNE
● Percent Unmarried	12.0
○ Percent WIC births	41.9
○ Percent Breastfeeding at discharge	87.5
○ Percent Payment-Private Insurance	71.8
● Percent Payment-Medicaid	22.9
● Percent C-Section	17.9

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

\*Due to lack of comparable data from all 50 states, significant differences cannot be calculated.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

○ All Causes	812.6
Heart Disease	154.2
Malignant Neoplasms (cancer)	155.3
Trachea, Bronchus, & Lung	34.5
Colon, Rectum, & Anus	20.9
Female Breast	LNE
Pancreas	15.1
Cerebrovascular Disease	55.3
Chronic Lower Respiratory Diseases	51.8
Accidents	41.8
Motor Vehicle Accidents	15.7
○ Alzheimer's Disease	117.0
Diabetes Mellitus	26.8
Influenza & Pneumonia	17.2
Intentional Self-Harm (suicide)	LNE
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	5.46

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	70
2. Malignant Neoplasms (cancer)	59
3. Alzheimer's Disease	58
4. Cerebrovascular Diseases	25
5. Chronic Lower Respiratory Diseases	23
6. Accidents	16
7. Diabetes	12
8. Vascular Dementia	11
T9. Influenza and Pneumonia	8
T9. Parkinson's Disease	8
Percent of Deaths due to tobacco use	13.4
Median age at death	84

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Hand County

## Demographic Information



Hand County is located in central South Dakota and averages 2.4 persons per square mile.

### 2012 Population Information

Subject	Number	Percent
Total population	3,388	100.0
White	3,332	98.3
Black or African American	3	0.1
American Indian & Alaska Native	14	0.4
Asian	11	0.3
Native Hawaiian & Other Pacific Islander	2	0.1
Two or More Races	26	0.8
Under 5 years	190	5.6
Under 18 years	706	20.8
65 years and over	864	25.5

Source: United States Census Bureau, 2012 Population Estimates

## Health Status Indicators 2008-2012

### Nativity

Percent of Low Birth Weight Infants	8.1
Percent of Mothers Receiving Care in 1st Trimester	70.9
● Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	8.2
Percent of Births Less Than 37 Wks. of Gestation	8.7
Average Age of Mother	27.8
Teenage Birth Rate <sup>2</sup>	LNE
Percent White Births	97.7
Percent American Indian Births	1.7
● Percent Unmarried	19.7
● Percent WIC births	25.0
Percent Breastfeeding at discharge	78.4
○ Percent Payment-Private Insurance	78.0
● Percent Payment-Medicaid	16.8
○ Percent C-Section	41.6

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

\*Due to lack of comparable data from all 50 states, significant differences cannot be calculated.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

● All Causes	539.6
● Heart Disease	114.6
● Malignant Neoplasms (cancer)	113.1
Trachea, Bronchus, & Lung	29.4
Colon, Rectum, & Anus	8.3
Female Breast	LNE
Pancreas	LNE
Cerebrovascular Disease	41.2
Chronic Lower Respiratory Diseases	42.8
Accidents	28.8
Motor Vehicle Accidents	LNE
● Alzheimer's Disease	15.3
Diabetes Mellitus	19.8
● Influenza & Pneumonia	5.6
Intentional Self-Harm (suicide)	LNE
Chronic Liver Disease and Cirrhosis	16.7
Infant Mortality	LNE

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	48
2. Malignant Neoplasms (cancer)	35
3. Cerebrovascular Diseases	17
4. Chronic Lower Respiratory Diseases	15
5. Atherosclerosis	14
6. Diabetes	7
7. Alzheimer's Disease	6
T8. Accidents	5
T8. Essential (primary) Hypertension and Hypertensive Renal Disease	5
Percent of Deaths due to tobacco use	18.8
Median age at death	83

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Hanson County

## Demographic Information



Hanson County is located in southeastern South Dakota and averages 7.7 people per square mile.

### 2012 Population Information

Subject	Number	Percent
Total population	3,377	100.0
White	3,318	98.3
Black or African American	8	0.2
American Indian & Alaska Native	16	0.5
Asian	15	0.4
Native Hawaiian & Other Pacific Islander	2	0.1
Two or More Races	18	0.5
Under 5 years	297	8.8
Under 18 years	1,075	31.8
65 years and over	466	13.8

Source: United States Census Bureau, 2012 Population Estimates

## Health Status Indicators 2008-2012

### Nativity

Percent of Low Birth Weight Infants	7.9
Percent of Mothers Receiving Care in 1st Trimester	61.9
● Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	6.6
Percent of Births Less Than 37 Wks. of Gestation	7.1
○ Average Age of Mother	29.6
Teenage Birth Rate <sup>2</sup>	LNE
Percent White Births	99.6
Percent American Indian Births	LNE
● Percent Unmarried	11.2
● Percent WIC births	19.1
Percent Breastfeeding at discharge	79.5
○ Percent Payment-Private Insurance	79.8
● Percent Payment-Medicaid	11.3
Percent C-Section	30.6

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

\*Due to lack of comparable data from all 50 states, significant differences cannot be calculated.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

All Causes	612.8
Heart Disease	137.4
Malignant Neoplasms (cancer)	181.7
Trachea, Bronchus, & Lung	61.1
Colon, Rectum, & Anus	LNE
Female Breast	LNE
Pancreas	LNE
Cerebrovascular Disease	55.2
● Chronic Lower Respiratory Diseases	18.1
Accidents	45.7
Motor Vehicle Accidents	LNE
Alzheimer's Disease	LNE
Diabetes Mellitus	LNE
Influenza & Pneumonia	LNE
Intentional Self-Harm (suicide)	LNE
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	LNE

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Malignant Neoplasms (cancer)	32
2. Heart Disease	21
3. Cerebrovascular Diseases	8
4. Accidents	7
Percent of Deaths due to tobacco use	12.9
Median age at death	74

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Harding County

## Demographic Information



Harding County is located in the northwestern corner of the state and averages 0.5 persons per square mile.

## 2012 Population Information

Subject	Number	Percent
Total population	1,316	100.0
White	1,267	96.3
Black or African American	5	0.4
American Indian & Alaska Native	22	1.7
Asian	1	0.1
Native Hawaiian & Other Pacific Islander	0	0.0
Two or More Races	21	1.6
Under 5 years	85	6.5
Under 18 years	299	22.7
65 years and over	200	15.2

Source: United States Census Bureau, 2012 Population Estimates

## Health Status Indicators 2008-2012

### Nativity

Percent of Low Birth Weight Infants	LNE
Percent of Mothers Receiving Care in 1st Trimester	75.0
● Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	6.2
Percent of Births Less Than 37 Wks. of Gestation	4.6
Average Age of Mother	28.3
Teenage Birth Rate <sup>2</sup>	LNE
Percent White Births	98.5
Percent American Indian Births	LNE
● Percent Unmarried	10.8
● Percent WIC births	13.8
Percent Breastfeeding at discharge	87.7
Percent Payment-Private Insurance	73.8
● Percent Payment-Medicaid	12.3
Percent C-Section	33.8

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

\*Due to lack of comparable data from all 50 states, significant differences cannot be calculated.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

All Causes	629.9
Heart Disease	218.4
● Malignant Neoplasms (cancer)	60.7
Trachea, Bronchus, & Lung	LNE
Colon, Rectum, & Anus	LNE
Female Breast	LNE
Pancreas	LNE
Cerebrovascular Disease	41.2
Chronic Lower Respiratory Diseases	54.4
Accidents	102.8
Motor Vehicle Accidents	LNE
Alzheimer's Disease	LNE
Diabetes Mellitus	LNE
Influenza & Pneumonia	LNE
Intentional Self-Harm (suicide)	LNE
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	LNE

Leading Causes of Death	Total Deaths
1. Heart Disease	17
2. Accidents	6
3. Malignant Neoplasms (cancer)	5
Percent of Deaths due to tobacco use	17.0
Median age at death	77

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Hughes County

## Demographic Information



Hughes County is located in the center of the state and averages 23.0 persons per square mile.

### 2012 Population Information

Subject	Number	Percent
Total population	17,450	100.0
White	14,871	85.2
Black or African American	251	1.4
American Indian & Alaska Native	1,812	10.4
Asian	98	0.6
Native Hawaiian & Other Pacific Islander	8	0.0
Two or More Races	410	2.3
Under 5 years	1,190	6.8
Under 18 years	4,040	23.2
65 years and over	2,454	14.1

Source: United States Census Bureau, 2012 Population Estimates

## Health Status Indicators 2008-2012

### Nativity

Percent of Low Birth Weight Infants	5.1
Percent of Mothers Receiving Care in 1st Trimester	67.9
Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	20.3
Percent of Births Less Than 37 Wks. of Gestation	7.5
Average Age of Mother	27.2
Teenage Birth Rate <sup>2</sup>	11.7
Percent White Births	74.3
Percent American Indian Births	20.3
Percent Unmarried	40.3
Percent WIC births	37.2
Percent Breastfeeding at discharge	73.5
Percent Payment-Private Insurance	59.1
Percent Payment-Medicaid	36.9
○ Percent C-Section	37.1

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

\*Due to lack of comparable data from all 50 states, significant differences cannot be calculated.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

All Causes	676.1
Heart Disease	149.0
Malignant Neoplasms (cancer)	166.5
Trachea, Bronchus, & Lung	38.1
Colon, Rectum, & Anus	18.7
Female Breast	13.7
Pancreas	14.4
Cerebrovascular Disease	45.1
Chronic Lower Respiratory Diseases	50.3
● Accidents	30.2
Motor Vehicle Accidents	10.7
Alzheimer's Disease	39.4
Diabetes Mellitus	20.2
Influenza & Pneumonia	10.6
Intentional Self-Harm (suicide)	17.4
Chronic Liver Disease and Cirrhosis	9.9
Infant Mortality	7.71

### Leading Causes of Death

Leading Causes of Death	Total Deaths
T1. Heart Disease	166
T1. Malignant Neoplasms (cancer)	166
3. Chronic Lower Respiratory Diseases	52
4. Cerebrovascular Diseases	50
5. Alzheimer's Disease	43
6. Accidents	31
7. Diabetes	22
8. Suicide	14
T9. Unspecified Dementia	13
T9. Influenza and Pneumonia	13
Percent of Deaths due to tobacco use	20.7
Median age at death	81

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Hutchinson County

## Demographic Information



Hutchinson County is located in the southeastern region of the state and averages 9.0 persons per square mile.

### 2012 Population Information

Subject	Number	Percent
Total population	7,187	100.0
White	6,989	97.2
Black or African American	46	0.6
American Indian & Alaska Native	75	1.0
Asian	18	0.3
Native Hawaiian & Other Pacific Islander	0	0.0
Two or More Races	59	0.8
Under 5 years	441	6.1
Under 18 years	1,726	24.0
65 years and over	1,738	24.2

Source: United States Census Bureau, 2012 Population Estimates

## Health Status Indicators 2008-2012

### Nativity

● Percent of Low Birth Weight Infants	3.5
Percent of Mothers Receiving Care in 1st Trimester	61.3
● Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	10.0
Percent of Births Less Than 37 Wks. of Gestation	6.5
○ Average Age of Mother	28.3
Teenage Birth Rate <sup>2</sup>	LNE
Percent White Births	98.8
Percent American Indian Births	LNE
● Percent Unmarried	19.3
● Percent WIC births	21.9
Percent Breastfeeding at discharge	81.3
○ Percent Payment-Private Insurance	78.1
● Percent Payment-Medicaid	18.2
Percent C-Section	23.8

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

\*Due to lack of comparable data from all 50 states, significant differences cannot be calculated.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

All Causes	669.0
Heart Disease	154.9
Malignant Neoplasms (cancer)	154.2
Trachea, Bronchus, & Lung	39.4
Colon, Rectum, & Anus	9.8
Female Breast	22.8
Pancreas	7.9
Cerebrovascular Disease	38.0
Chronic Lower Respiratory Diseases	39.5
Accidents	44.2
Motor Vehicle Accidents	12.6
Alzheimer's Disease	28.5
Diabetes Mellitus	16.3
Influenza & Pneumonia	18.6
Intentional Self-Harm (suicide)	12.7
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	LNE

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	154
2. Malignant Neoplasms (cancer)	106
3. Cerebrovascular Diseases	39
4. Alzheimer's Disease	37
5. Chronic Lower Respiratory Diseases	28
6. Accidents	25
7. Influenza and Pneumonia	18
8. Nephritis, Nephrotic Syndrome, and Nephrosis	14
9. Pneumonitis Due to Solids and Liquids	12
10. Diabetes	11

Percent of Deaths due to tobacco use	10.4
Median age at death	85

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Hyde County

## Demographic Information



Hyde County is located in the central region of the state and averages 1.7 people per square mile.

### 2012 Population Information

Subject	Number	Percent
Total population	1,437	100.0
White	1,283	89.3
Black or African American	6	0.4
American Indian & Alaska Native	115	8.0
Asian	5	0.3
Native Hawaiian & Other Pacific Islander	1	0.1
Two or More Races	27	1.9
Under 5 years	77	5.4
Under 18 years	323	22.5
65 years and over	322	22.4

Source: United States Census Bureau, 2012 Population Estimates

## Health Status Indicators 2008-2012

### Nativity

Percent of Low Birth Weight Infants	LNE
Percent of Mothers Receiving Care in 1st Trimester	72.7
Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	18.2
Percent of Births Less Than 37 Wks. of Gestation	7.3
Average Age of Mother	26.5
Teenage Birth Rate <sup>2</sup>	LNE
Percent White Births	83.6
Percent American Indian Births	10.9
Percent Unmarried	32.7
Percent WIC births	45.5
Percent Breastfeeding at discharge	73.6
Percent Payment-Private Insurance	61.8
Percent Payment-Medicaid	36.4
Percent C-Section	25.5

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

\*Due to lack of comparable data from all 50 states, significant differences cannot be calculated.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

All Causes	762.9
○ Heart Disease	332.0
Malignant Neoplasms (cancer)	128.0
Trachea, Bronchus, & Lung	24.7
Colon, Rectum, & Anus	LNE
Female Breast	LNE
Pancreas	LNE
Cerebrovascular Disease	59.7
● Chronic Lower Respiratory Diseases	19.1
Accidents	95.4
Motor Vehicle Accidents	65.8
Alzheimer's Disease	LNE
Diabetes Mellitus	LNE
Influenza & Pneumonia	LNE
Intentional Self-Harm (suicide)	LNE
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	LNE

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	49
2. Malignant Neoplasms (cancer)	16
3. Cerebrovascular Diseases	11
4. Accidents	6
Percent of Deaths due to tobacco use	9.8
Median age at death	85

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Jackson County

## Demographic Information



Jackson County is located in western South Dakota and averages 1.6 persons per square mile.

### 2012 Population Information

Subject	Number	Percent
Total population	3,191	100.0
White	1,368	42.9
Black or African American	17	0.5
American Indian & Alaska Native	1,647	51.6
Asian	3	0.1
Native Hawaiian & Other Pacific Islander	0	0.0
Two or More Races	156	4.9
Under 5 years	317	9.9
Under 18 years	1,036	32.5
65 years and over	423	13.3

Source: United States Census Bureau, 2012 Population Estimates

## Health Status Indicators 2008-2012

### Nativity

Percent of Low Birth Weight Infants	8.4
● Percent of Mothers Receiving Care in 1st Trimester	46.5
Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	20.4
○ Percent of Births Less Than 37 Wks. of Gestation	13.3
● Average Age of Mother	25.2
○ Teenage Birth Rate <sup>2</sup>	42.6
Percent White Births	19.7
Percent American Indian Births	75.8
○ Percent Unmarried	75.8
○ Percent WIC births	76.3
● Percent Breastfeeding at discharge	53.7
● Percent Payment-Private Insurance	14.4
○ Percent Payment-Medicaid	65.4
Percent C-Section	23.5

●Denotes a health status indicator which is significantly lower than the state average.  
○Denotes a health status indicator which is significantly higher than the state average.

\*Due to lack of comparable data from all 50 states, significant differences cannot be calculated.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

○ All Causes	1,053.6
Heart Disease	165.2
Malignant Neoplasms (cancer)	152.6
Trachea, Bronchus, & Lung	69.9
Colon, Rectum, & Anus	25.3
Female Breast	LNE
Pancreas	LNE
Cerebrovascular Disease	17.9
Chronic Lower Respiratory Diseases	81.8
○ Accidents	159.9
○ Motor Vehicle Accidents	81.2
Alzheimer's Disease	34.1
Diabetes Mellitus	35.8
Influenza & Pneumonia	27.4
Intentional Self-Harm (suicide)	43.4
○ Chronic Liver Disease and Cirrhosis	66.8
Infant Mortality	14.01

Leading Causes of Death	Total Deaths
1. Heart Disease	26
T2. Accidents	22
T2. Malignant Neoplasms (cancer)	22
4. Chronic Lower Respiratory Diseases	13
5. Chronic Liver Disease and Cirrhosis	9
T6. Suicide	6
T6. Alzheimer's Disease	6
T6. Diabetes	6
9. Influenza and Pneumonia	5
Percent of Deaths due to tobacco use	14.7
Median age at death	71

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Jerauld County

## Demographic Information



Jerauld County is located in the central region of the state and averages 3.9 persons per square mile.

### 2012 Population Information

Subject	Number	Percent
Total population	2,047	100.0
White	2,011	98.2
Black or African American	4	0.2
American Indian & Alaska Native	7	0.3
Asian	5	0.2
Native Hawaiian & Other Pacific Islander	2	0.1
Two or More Races	18	0.9
Under 5 years	145	7.1
Under 18 years	448	21.9
65 years and over	507	24.8

Source: United States Census Bureau, 2012 Population Estimates

## Health Status Indicators 2008-2012

### Nativity

Percent of Low Birth Weight Infants	5.4
Percent of Mothers Receiving Care in 1st Trimester	66.7
Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	12.5
Percent of Births Less Than 37 Wks. of Gestation	4.7
Average Age of Mother	27.9
Teenage Birth Rate <sup>2</sup>	21.0
Percent White Births	99.2
Percent American Indian Births	LNE
● Percent Unmarried	24.0
Percent WIC births	32.8
Percent Breastfeeding at discharge	82.9
Percent Payment-Private Insurance	61.2
Percent Payment-Medicaid	27.1
Percent C-Section	25.6

●Denotes a health status indicator which is significantly lower than the state average.  
○Denotes a health status indicator which is significantly higher than the state average.

\*Due to lack of comparable data from all 50 states, significant differences cannot be calculated.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

All Causes	704.6
Heart Disease	134.4
Malignant Neoplasms (cancer)	206.5
Trachea, Bronchus, & Lung	36.9
Colon, Rectum, & Anus	23.8
Female Breast	LNE
Pancreas	LNE
Cerebrovascular Disease	65.2
Chronic Lower Respiratory Diseases	36.0
Accidents	59.8
Motor Vehicle Accidents	33.3
○ Alzheimer's Disease	81.0
Diabetes Mellitus	LNE
Influenza & Pneumonia	LNE
Intentional Self-Harm (suicide)	29.4
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	LNE

### Leading Causes of Death

Leading Causes of Death	Total Deaths
T1. Heart Disease	35
T1. Malignant Neoplasms (cancer)	35
3. Alzheimer's Disease	24
4. Cerebrovascular Diseases	16
5. Chronic Lower Respiratory Diseases	9
6. Accidents	8
Percent of Deaths due to tobacco use	12.5
Median age at death	83

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Jones County

## Demographic Information



Jones County is located in western South Dakota and averages 1.0 persons per square mile.

### 2012 Population Information

Subject	Number	Percent
Total population	1,013	100.0
White	945	93.3
Black or African American	4	0.4
American Indian & Alaska Native	29	2.9
Asian	1	0.1
Native Hawaiian & Other Pacific Islander	2	0.2
Two or More Races	32	3.2
Under 5 years	69	6.8
Under 18 years	231	22.8
65 years and over	222	21.9

Source: United States Census Bureau, 2012 Population Estimates

## Health Status Indicators 2008-2012

### Natality

Percent of Low Birth Weight Infants	6.9
Percent of Mothers Receiving Care in 1st Trimester	81.0
Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	15.5
Percent of Births Less Than 37 Wks. of Gestation	15.5
Average Age of Mother	26.3
Teenage Birth Rate <sup>2</sup>	25.6
Percent White Births	96.6
Percent American Indian Births	LNE
● Percent Unmarried	19.0
Percent WIC births	44.8
Percent Breastfeeding at discharge	82.8
Percent Payment-Private Insurance	71.9
Percent Payment-Medicaid	24.6
Percent C-Section	36.2

●Denotes a health status indicator which is significantly lower than the state average.  
○Denotes a health status indicator which is significantly higher than the state average.

\*Due to lack of comparable data from all 50 states, significant differences cannot be calculated.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

● All Causes	514.3
● Heart Disease	62.9
Malignant Neoplasms (cancer)	129.1
Trachea, Bronchus, & Lung	LNE
Colon, Rectum, & Anus	LNE
Female Breast	63.4
Pancreas	LNE
Cerebrovascular Disease	35.6
Chronic Lower Respiratory Diseases	58.6
Accidents	LNE
Motor Vehicle Accidents	LNE
Alzheimer's Disease	LNE
Diabetes Mellitus	LNE
Influenza & Pneumonia	27.0
Intentional Self-Harm (suicide)	LNE
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	LNE

### Leading Causes of Death

	Total Deaths
1. Malignant Neoplasms (cancer)	12
2. Heart Disease	6
3. Chronic Lower Respiratory Diseases	5
Percent of Deaths due to tobacco use	18.2
Median age at death	81

●Denotes a health status indicator which is significantly lower than the state average.  
○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Kingsbury County

## Demographic Information



Kingsbury County is located in east central South Dakota and averages 6.2 persons per square mile.

### 2012 Population Information

Subject	Number	Percent
Total population	5,220	100.0
White	5,083	97.4
Black or African American	24	0.5
American Indian & Alaska Native	32	0.6
Asian	23	0.4
Native Hawaiian & Other Pacific Islander	0	0.0
Two or More Races	58	1.1
Under 5 years	352	6.7
Under 18 years	1,191	22.8
65 years and over	1,083	20.7

Source: United States Census Bureau, 2012 Population Estimates

## Health Status Indicators 2008-2012

### Nativity

Percent of Low Birth Weight Infants	4.1
Percent of Mothers Receiving Care in 1st Trimester	71.5
Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	14.0
Percent of Births Less Than 37 Wks. of Gestation	5.7
○ Average Age of Mother	28.1
Teenage Birth Rate <sup>2</sup>	LNE
Percent White Births	97.5
Percent American Indian Births	1.6
● Percent Unmarried	20.9
● Percent WIC births	23.7
Percent Breastfeeding at discharge	79.2
○ Percent Payment-Private Insurance	79.7
● Percent Payment-Medicaid	16.8
Percent C-Section	21.5

### Mortality<sup>3</sup>

All Causes	747.9
Heart Disease	169.4
Malignant Neoplasms (cancer)	160.5
Trachea, Bronchus, & Lung	51.7
Colon, Rectum, & Anus	19.4
Female Breast	11.2
Pancreas	8.5
Cerebrovascular Disease	56.7
Chronic Lower Respiratory Diseases	32.1
Accidents	62.5
Motor Vehicle Accidents	37.7
Alzheimer's Disease	37.2
Diabetes Mellitus	25.1
Influenza & Pneumonia	19.7
Intentional Self-Harm (suicide)	13.9
Chronic Liver Disease and Cirrhosis	7.4
Infant Mortality	9.49

### Leading Causes of Death

	Total Deaths
1. Heart Disease	96
2. Malignant Neoplasms (cancer)	72
3. Cerebrovascular Diseases	34
4. Alzheimer's Disease	23
5. Accidents	20
6. Chronic Lower Respiratory Diseases	16
7. Influenza and Pneumonia	12
8. Diabetes	11
9. Unspecified Dementia	8
10. Vascular Dementia	7
Percent of Deaths due to tobacco use	13.1
Median age at death	83

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

\*Due to lack of comparable data from all 50 states, significant differences cannot be calculated.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

●Denotes a health status indicator which is significantly lower than the state average.  
○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Lake County

## Demographic Information



Lake County is located in the east central region of the state and averages 19.9 persons per square mile.

### 2012 Population Information

Subject	Number	Percent
Total population	11,771	100.0
White	11,331	96.3
Black or African American	106	0.9
American Indian & Alaska Native	94	0.8
Asian	86	0.7
Native Hawaiian & Other Pacific Islander	5	0.0
Two or More Races	149	1.3
Under 5 years	659	5.6
Under 18 years	2,398	20.4
65 years and over	2,127	18.1

Source: United States Census Bureau, 2012 Population Estimates

## Health Status Indicators 2008-2012

### Nativity

Percent of Low Birth Weight Infants	5.1
Percent of Mothers Receiving Care in 1st Trimester	72.6
Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	15.6
Percent of Births Less Than 37 Wks. of Gestation	6.5
○ Average Age of Mother	28.1
● Teenage Birth Rate <sup>2</sup>	8.8
Percent White Births	97.2
Percent American Indian Births	1.6
● Percent Unmarried	22.2
● Percent WIC births	27.0
Percent Breastfeeding at discharge	78.0
○ Percent Payment-Private Insurance	72.6
● Percent Payment-Medicaid	21.7
Percent C-Section	29.1

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

\*Due to lack of comparable data from all 50 states, significant differences cannot be calculated.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

● All Causes	641.9
Heart Disease	151.3
Malignant Neoplasms (cancer)	162.7
Trachea, Bronchus, & Lung	32.8
● Colon, Rectum, & Anus	8.3
Female Breast	31.6
Pancreas	13.2
Cerebrovascular Disease	46.9
Chronic Lower Respiratory Diseases	50.9
● Accidents	23.1
Motor Vehicle Accidents	8.6
○ Alzheimer's Disease	56.7
Diabetes Mellitus	14.6
● Influenza & Pneumonia	2.8
Intentional Self-Harm (suicide)	11.7
Chronic Liver Disease and Cirrhosis	8.9
Infant Mortality	6.43

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Malignant Neoplasms (cancer)	134
2. Heart Disease	129
3. Alzheimer's Disease	53
4. Chronic Lower Respiratory Diseases	44
5. Cerebrovascular Diseases	41
6. Accidents	17
7. Diabetes	12
8. Essential (primary) Hypertension and Hypertensive Renal Disease	8
9. Vascular Dementia	7
T10. Suicide	6
T10. Chronic Liver Disease and Cirrhosis	6
T10. Aortic Aneurysm and Dissection	6
T10. Septicemia	6

Percent of Deaths due to tobacco use 19.8

Median age at death 81

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Lawrence County

## Demographic Information



Lawrence County is located along the Wyoming border and averages 30.1 persons per square mile.

### 2012 Population Information

Subject	Number	Percent
Total population	24,397	100.0
White	22,963	94.1
Black or African American	209	0.9
American Indian & Alaska Native	548	2.2
Asian	180	0.7
Native Hawaiian & Other Pacific Islander	14	0.1
Two or More Races	483	2.0
Under 5 years	1,288	5.3
Under 18 years	4,587	18.8
65 years and over	4,181	17.1

Source: United States Census Bureau, 2012 Population Estimates

## Health Status Indicators 2008-2012

### Nativity

Percent of Low Birth Weight Infants	8.1
○ Percent of Mothers Receiving Care in 1st Trimester	79.3
Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	20.3
Percent of Births Less Than 37 Wks. of Gestation	7.6
Average Age of Mother	27.4
Teenage Birth Rate <sup>2</sup>	14.4
Percent White Births	93.3
Percent American Indian Births	2.1
Percent Unmarried	36.6
Percent WIC births	38.4
○ Percent Breastfeeding at discharge	80.6
Percent Payment-Private Insurance	54.6
Percent Payment-Medicaid	37.1
● Percent C-Section	22.1

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

\*Due to lack of comparable data from all 50 states, significant differences cannot be calculated.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

● All Causes	635.7
Heart Disease	140.1
Malignant Neoplasms (cancer)	164.3
Trachea, Bronchus, & Lung	43.6
Colon, Rectum, & Anus	11.9
Female Breast	23.1
Pancreas	14.1
Cerebrovascular Disease	36.2
Chronic Lower Respiratory Diseases	49.0
● Accidents	29.3
● Motor Vehicle Accidents	7.5
● Alzheimer's Disease	25.1
Diabetes Mellitus	21.9
● Influenza & Pneumonia	7.3
Intentional Self-Harm (suicide)	15.0
Chronic Liver Disease and Cirrhosis	9.1
Infant Mortality	7.38

### Leading Causes of Death

	Total Deaths
1. Malignant Neoplasms (cancer)	267
2. Heart Disease	240
3. Chronic Lower Respiratory Diseases	83
4. Cerebrovascular Diseases	64
5. Alzheimer's Disease	47
6. Accidents	42
7. Diabetes	37
T8. Unspecified Dementia	19
T8. Suicide	19
T10. Vascular Dementia	14
T10. Chronic Liver Disease and Cirrhosis	14
T10. Influenza and Pneumonia	14
Percent of Deaths due to tobacco use	19.5
Median age at death	81

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Lincoln County

## Demographic Information



Lincoln County is located in southeastern South Dakota and averages 77.7 persons per square mile.

### 2012 Population Information

Subject	Number	Percent
Total population	48,296	100.0
White	46,270	95.8
Black or African American	479	1.0
American Indian & Alaska Native	303	0.6
Asian	544	1.1
Native Hawaiian & Other Pacific Islander	13	0.0
Two or More Races	687	1.4
Under 5 years	4,279	8.9
Under 18 years	14,065	29.1
65 years and over	4,529	9.4

Source: United States Census Bureau, 2012 Population Estimates

## Health Status Indicators 2008-2012

### Nativity

Percent of Low Birth Weight Infants	6.0
○ Percent of Mothers Receiving Care in 1st Trimester	79.8
● Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	7.7
Percent of Births Less Than 37 Wks. of Gestation	8.1
○ Average Age of Mother	28.9
● Teenage Birth Rate <sup>2</sup>	6.7
Percent White Births	97.6
Percent American Indian Births	0.4
● Percent Unmarried	15.6
● Percent WIC births	10.9
○ Percent Breastfeeding at discharge	79.4
○ Percent Payment-Private Insurance	86.4
● Percent Payment-Medicaid	11.4
Percent C-Section	27.0

### Mortality<sup>3</sup>

● All Causes	577.1
● Heart Disease	128.9
● Malignant Neoplasms (cancer)	138.5
Trachea, Bronchus, & Lung	39.2
Colon, Rectum, & Anus	13.6
Female Breast	22.6
Pancreas	9.5
Cerebrovascular Disease	31.6
Chronic Lower Respiratory Diseases	36.3
● Accidents	28.4
● Motor Vehicle Accidents	7.7
Alzheimer's Disease	34.9
Diabetes Mellitus	21.8
● Influenza & Pneumonia	8.2
● Intentional Self-Harm (suicide)	10.2
● Chronic Liver Disease and Cirrhosis	3.1
Infant Mortality	5.46

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Malignant Neoplasms (cancer)	228
2. Heart Disease	216
3. Alzheimer's Disease	58
4. Chronic Lower Respiratory Diseases	55
5. Accidents	52
6. Cerebrovascular Diseases	48
7. Diabetes	33
8. Suicide	22
9. Unspecified Dementia	21
10. Parkinson's Disease	17
Percent of Deaths due to tobacco use	15.4
Median age at death	81

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

\*Due to lack of comparable data from all 50 states, significant differences cannot be calculated.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

●Denotes a health status indicator which is significantly lower than the state average.  
○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Lyman County

## Demographic Information



Lyman County is located in central South Dakota and averages 2.3 persons per square mile.

### 2012 Population Information

Subject	Number	Percent
Total population	3,789	100.0
White	2,187	57.7
Black or African American	12	0.3
American Indian & Alaska Native	1,480	39.1
Asian	10	0.3
Native Hawaiian & Other Pacific Islander	1	0.0
Two or More Races	99	2.6
Under 5 years	288	7.6
Under 18 years	1,096	28.9
65 years and over	564	14.9

Source: United States Census Bureau, 2012 Population Estimates

## Health Status Indicators 2008-2012

### Nativity

Percent of Low Birth Weight Infants	4.4
● Percent of Mothers Receiving Care in 1st Trimester	48.9
○ Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	26.3
Percent of Births Less Than 37 Wks. of Gestation	6.2
● Average Age of Mother	25.9
Teenage Birth Rate <sup>2</sup>	35.5
Percent White Births	37.1
Percent American Indian Births	59.5
○ Percent Unmarried	62.9
○ Percent WIC births	67.1
● Percent Breastfeeding at discharge	62.7
● Percent Payment-Private Insurance	35.6
○ Percent Payment-Medicaid	59.1
○ Percent C-Section	35.5

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

\*Due to lack of comparable data from all 50 states, significant differences cannot be calculated.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

All Causes	834.9
Heart Disease	192.2
Malignant Neoplasms (cancer)	144.5
Trachea, Bronchus, & Lung	70.1
Colon, Rectum, & Anus	21.9
Female Breast	LNE
Pancreas	LNE
Cerebrovascular Disease	23.5
Chronic Lower Respiratory Diseases	48.5
○ Accidents	155.0
○ Motor Vehicle Accidents	95.0
Alzheimer's Disease	49.3
Diabetes Mellitus	25.1
Influenza & Pneumonia	19.0
Intentional Self-Harm (suicide)	LNE
Chronic Liver Disease and Cirrhosis	13.5
Infant Mortality	LNE

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	39
2. Malignant Neoplasms (cancer)	32
3. Accidents	28
T4. Chronic Lower Respiratory Diseases	10
T4. Alzheimer's Disease	10
6. Diabetes	6
7. Cerebrovascular Diseases	5
Percent of Deaths due to tobacco use	25.1
Median age at death	72

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# McCook County

## Demographic Information



McCook County is located in eastern South Dakota and averages 9.8 persons per square mile.

## 2012 Population Information

Subject	Number	Percent
Total population	5,610	100.0
White	5,498	98.0
Black or African American	21	0.4
American Indian & Alaska Native	35	0.6
Asian	14	0.2
Native Hawaiian & Other Pacific Islander	3	0.1
Two or More Races	39	0.7
Under 5 years	390	7.0
Under 18 years	1,420	25.3
65 years and over	1,071	19.1

Source: United States Census Bureau, 2012 Population Estimates

## Health Status Indicators 2008-2012

### Nativity

Percent of Low Birth Weight Infants	6.0
Percent of Mothers Receiving Care in 1st Trimester	67.8
● Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	10.7
Percent of Births Less Than 37 Wks. of Gestation	8.2
○ Average Age of Mother	28.6
● Teenage Birth Rate <sup>2</sup>	5.6
Percent White Births	98.1
Percent American Indian Births	0.8
● Percent Unmarried	20.4
● Percent WIC births	31.6
Percent Breastfeeding at discharge	75.5
○ Percent Payment-Private Insurance	75.3
● Percent Payment-Medicaid	21.1
Percent C-Section	25.3

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

\*Due to lack of comparable data from all 50 states, significant differences cannot be calculated.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

○ All Causes	843.8
○ Heart Disease	214.5
○ Malignant Neoplasms (cancer)	213.3
Trachea, Bronchus, & Lung	62.8
Colon, Rectum, & Anus	9.4
Female Breast	15.9
Pancreas	9.4
Cerebrovascular Disease	61.1
Chronic Lower Respiratory Diseases	49.6
Accidents	41.1
Motor Vehicle Accidents	24.8
Alzheimer's Disease	52.4
Diabetes Mellitus	26.1
Influenza & Pneumonia	16.9
Intentional Self-Harm (suicide)	LNE
Chronic Liver Disease and Cirrhosis	7.1
Infant Mortality	8.15

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	101
2. Malignant Neoplasms (cancer)	89
3. Cerebrovascular Diseases	30
4. Alzheimer's Disease	28
5. Chronic Lower Respiratory Diseases	23
6. Diabetes	12
7. Accidents	11
8. Influenza and Pneumonia	9
9. Septicemia	6
T10. Vascular Dementia	5
T10. Nephritis, Nephrotic Syndrome, and Nephrosis	5
T10. Parkinson's Disease	5
Percent of Deaths due to tobacco use	15.4
Median age at death	82

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# McPherson County

## Demographic Information



McPherson County is located in the north central region of the state and averages 2.2 persons per square mile.

### 2012 Population Information

Subject	Number	Percent
Total population	2,439	100.0
White	2,399	98.4
Black or African American	5	0.2
American Indian & Alaska Native	2	0.1
Asian	11	0.5
Native Hawaiian & Other Pacific Islander	3	0.1
Two or More Races	19	0.8
Under 5 years	120	4.9
Under 18 years	512	21.0
65 years and over	721	29.6

Source: United States Census Bureau, 2012 Population Estimates

## Health Status Indicators 2008-2012

### Nativity

Percent of Low Birth Weight Infants	5.1
Percent of Mothers Receiving Care in 1st Trimester	68.0
Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	11.2
Percent of Births Less Than 37 Wks. of Gestation	6.1
○ Average Age of Mother	30.0
Teenage Birth Rate <sup>2</sup>	LNE
Percent White Births	98.0
Percent American Indian Births	LNE
● Percent Unmarried	14.1
● Percent WIC births	18.4
Percent Breastfeeding at discharge	84.7
○ Percent Payment-Private Insurance	80.8
● Percent Payment-Medicaid	14.1
● Percent C-Section	17.2

●Denotes a health status indicator which is significantly lower than the state average.  
○Denotes a health status indicator which is significantly higher than the state average.

\*Due to lack of comparable data from all 50 states, significant differences cannot be calculated.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

● All Causes	525.5
Heart Disease	154.5
● Malignant Neoplasms (cancer)	76.4
Trachea, Bronchus, & Lung	32.6
Colon, Rectum, & Anus	11.6
Female Breast	LNE
Pancreas	LNE
Cerebrovascular Disease	65.3
● Chronic Lower Respiratory Diseases	19.3
Accidents	44.0
Motor Vehicle Accidents	15.7
● Alzheimer's Disease	14.2
Diabetes Mellitus	23.8
Influenza & Pneumonia	8.5
Intentional Self-Harm (suicide)	LNE
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	LNE

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	54
T2. Cerebrovascular Diseases	22
T2. Malignant Neoplasms (cancer)	22
4. Essential (primary) Hypertension and Hypertensive Renal Disease	10
T5. Accidents	8
T5. Diabetes	8
7. Chronic Lower Respiratory Diseases	7
8. Alzheimer's Disease	6
Percent of Deaths due to tobacco use	11.0
Median age at death	85

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Marshall County

## Demographic Information



Marshall County is located in the northeastern part of the state and averages 5.6 persons per square mile.

### 2012 Population Information

Subject	Number	Percent
Total population	4,671	100.0
White	4,114	88.1
Black or African American	41	0.9
American Indian & Alaska Native	430	9.2
Asian	15	0.3
Native Hawaiian & Other Pacific Islander	1	0.0
Two or More Races	70	1.5
Under 5 years	271	5.8
Under 18 years	1,016	21.8
65 years and over	927	19.8

Source: United States Census Bureau, 2012 Population Estimates

## Health Status Indicators 2008-2012

### Nativity

Percent of Low Birth Weight Infants	7.4
Percent of Mothers Receiving Care in 1st Trimester	63.2
Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	17.8
Percent of Births Less Than 37 Wks. of Gestation	5.9
Average Age of Mother	27.3
Teenage Birth Rate <sup>2</sup>	10.4
Percent White Births	81.1
Percent American Indian Births	17.4
Percent Unmarried	31.4
● Percent WIC births	31.1
Percent Breastfeeding at discharge	73.1
Percent Payment-Private Insurance	61.0
Percent Payment-Medicaid	29.0
Percent C-Section	29.0

### Mortality<sup>3</sup>

○ All Causes	812.3
Heart Disease	165.5
Malignant Neoplasms (cancer)	199.2
Trachea, Bronchus, & Lung	49.4
Colon, Rectum, & Anus	26.5
Female Breast	25.8
Pancreas	25.8
Cerebrovascular Disease	55.4
Chronic Lower Respiratory Diseases	59.4
Accidents	46.5
Motor Vehicle Accidents	22.7
Alzheimer's Disease	40.5
Diabetes Mellitus	29.6
Influenza & Pneumonia	20.7
Intentional Self-Harm (suicide)	LNE
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	25.74

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Malignant Neoplasms (cancer)	70
2. Heart Disease	65
3. Chronic Lower Respiratory Diseases	26
4. Cerebrovascular Diseases	24
5. Alzheimer's Disease	20
6. Accidents	11
7. Influenza and Pneumonia	9
8. Diabetes	8
Percent of Deaths due to tobacco use	16.4
Median age at death	83

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

\*Due to lack of comparable data from all 50 states, significant differences cannot be calculated.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

○Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Meade County

## Demographic Information



Meade County is located in west central South Dakota and averages 7.3 persons per square mile.

### 2012 Population Information

Subject	Number	Percent
Total population	26,052	100.0
White	23,885	91.7
Black or African American	468	1.8
American Indian & Alaska Native	740	2.8
Asian	208	0.8
Native Hawaiian & Other Pacific Islander	25	0.1
Two or More Races	726	2.8
Under 5 years	1,727	6.6
Under 18 years	6,173	23.7
65 years and over	3,344	12.8

Source: United States Census Bureau, 2012 Population Estimates

## Health Status Indicators 2008-2012

### Nativity

Percent of Low Birth Weight Infants	6.4
○ Percent of Mothers Receiving Care in 1st Trimester	77.5
Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	15.6
Percent of Births Less Than 37 Wks. of Gestation	7.5
Average Age of Mother	27.1
Teenage Birth Rate <sup>2</sup>	13.3
Percent White Births	92.0
Percent American Indian Births	3.4
● Percent Unmarried	26.0
● Percent WIC births	34.6
○ Percent Breastfeeding at discharge	83.0
Percent Payment-Private Insurance	51.9
● Percent Payment-Medicaid	27.9
● Percent C-Section	21.8

●Denotes a health status indicator which is significantly lower than the state average.  
○Denotes a health status indicator which is significantly higher than the state average.

\*Due to lack of comparable data from all 50 states, significant differences cannot be calculated.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

All Causes	728.2
Heart Disease	163.7
Malignant Neoplasms (cancer)	168.7
Trachea, Bronchus, & Lung	50.3
Colon, Rectum, & Anus	10.5
Female Breast	13.7
Pancreas	11.8
Cerebrovascular Disease	44.7
Chronic Lower Respiratory Diseases	53.8
Accidents	37.5
Motor Vehicle Accidents	14.7
● Alzheimer's Disease	24.4
Diabetes Mellitus	26.6
Influenza & Pneumonia	15.2
Intentional Self-Harm (suicide)	20.5
Chronic Liver Disease and Cirrhosis	14.5
Infant Mortality	8.44

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Malignant Neoplasms (cancer)	216
2. Heart Disease	200
3. Chronic Lower Respiratory Diseases	68
4. Cerebrovascular Diseases	54
5. Accidents	47
6. Diabetes	32
7. Alzheimer's Disease	28
8. Suicide	25
9. Chronic Liver Disease and Cirrhosis	20
10. Influenza and Pneumonia	18
Percent of Deaths due to tobacco use	20.9
Median age at death	77

●Denotes a health status indicator which is significantly lower than the state average.  
○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Mellette County

## Demographic Information



Mellette County is located in the south central region of the state and averages 1.6 persons per square mile.

### 2012 Population Information

Subject	Number	Percent
Total population	2,101	100.0
White	871	41.5
Black or African American	3	0.1
American Indian & Alaska Native	1,101	52.4
Asian	7	0.3
Native Hawaiian & Other Pacific Islander	0	0.0
Two or More Races	119	5.7
Under 5 years	197	9.4
Under 18 years	648	30.8
65 years and over	315	15.0

Source: United States Census Bureau, 2012 Population Estimates

## Health Status Indicators 2008-2012

### Nativity

Percent of Low Birth Weight Infants	6.7
● Percent of Mothers Receiving Care in 1st Trimester	45.3
Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	23.0
Percent of Births Less Than 37 Wks. of Gestation	6.7
● Average Age of Mother	24.3
Teenage Birth Rate <sup>2</sup>	41.4
Percent White Births	21.0
Percent American Indian Births	65.4
○ Percent Unmarried	70.6
○ Percent WIC births	66.0
Percent Breastfeeding at discharge	72.8
● Percent Payment-Private Insurance	12.9
○ Percent Payment-Medicaid	57.1
Percent C-Section	35.0

### Mortality<sup>3</sup>

○ All Causes	1,062.1
Heart Disease	147.7
Malignant Neoplasms (cancer)	235.2
Trachea, Bronchus, & Lung	59.7
Colon, Rectum, & Anus	31.3
Female Breast	53.0
Pancreas	LNE
Cerebrovascular Disease	65.2
Chronic Lower Respiratory Diseases	100.4
Accidents	102.8
○ Motor Vehicle Accidents	67.8
Alzheimer's Disease	23.7
Diabetes Mellitus	54.4
Influenza & Pneumonia	LNE
Intentional Self-Harm (suicide)	28.3
○ Chronic Liver Disease and Cirrhosis	70.0
Infant Mortality	18.40

### Leading Causes of Death

	Total Deaths
1. Malignant Neoplasms (cancer)	27
2. Heart Disease	17
T3. Accidents	11
T3. Chronic Lower Respiratory Diseases	11
5. Cerebrovascular Diseases	8
T6. Chronic Liver Disease and Cirrhosis	6
T6. Diabetes	6
Percent of Deaths due to tobacco use	21.7
Median age at death	75

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

\*Due to lack of comparable data from all 50 states, significant differences cannot be calculated.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Miner County

## Demographic Information



Miner County is located in the east central region of the state and averages 4.2 persons per square mile.

### 2012 Population Information

Subject	Number	Percent
Total population	2,326	100.0
White	2,282	98.1
Black or African American	8	0.3
American Indian & Alaska Native	5	0.2
Asian	9	0.4
Native Hawaiian & Other Pacific Islander	0	0.0
Two or More Races	22	0.9
Under 5 years	136	5.8
Under 18 years	542	23.3
65 years and over	498	21.4

Source: United States Census Bureau, 2012 Population Estimates

## Health Status Indicators 2008-2012

### Nativity

Percent of Low Birth Weight Infants	LNE
Percent of Mothers Receiving Care in 1st Trimester	74.2
Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	13.2
● Percent of Births Less Than 37 Wks. of Gestation	3.3
○ Average Age of Mother	28.4
Teenage Birth Rate <sup>2</sup>	14.0
Percent White Births	95.9
Percent American Indian Births	2.5
● Percent Unmarried	18.2
● Percent WIC births	26.4
Percent Breastfeeding at discharge	79.7
○ Percent Payment-Private Insurance	75.2
● Percent Payment-Medicaid	21.5
Percent C-Section	28.1

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

\*Due to lack of comparable data from all 50 states, significant differences cannot be calculated.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

All Causes	694.2
Heart Disease	166.5
Malignant Neoplasms (cancer)	189.2
Trachea, Bronchus, & Lung	44.2
Colon, Rectum, & Anus	22.5
Female Breast	LNE
Pancreas	17.8
Cerebrovascular Disease	37.0
Chronic Lower Respiratory Diseases	58.5
Accidents	31.8
Motor Vehicle Accidents	31.8
○ Alzheimer's Disease	91.0
Diabetes Mellitus	LNE
Influenza & Pneumonia	11.7
Intentional Self-Harm (suicide)	LNE
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	LNE

### Leading Causes of Death

Leading Causes of Death	Total Deaths
T1. Heart Disease	44
T1. Malignant Neoplasms (cancer)	44
3. Alzheimer's Disease	28
4. Chronic Lower Respiratory Diseases	14
5. Cerebrovascular Diseases	12
Percent of Deaths due to tobacco use	20.3
Median age at death	86

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Minnehaha County

## Demographic Information



Minnehaha County is located in southeastern South Dakota and averages 210.0 persons per square mile.

### 2012 Population Information

Subject	Number	Percent
Total population	175,037	100.0
White	155,918	89.1
Black or African American	7,411	4.2
American Indian & Alaska Native	4,838	2.8
Asian	2,770	1.6
Native Hawaiian & Other Pacific Islander	223	0.1
Two or More Races	3,877	2.2
Under 5 years	13,186	7.5
Under 18 years	43,372	24.8
65 years and over	20,281	11.6

Source: United States Census Bureau, 2012 Population Estimates

## Health Status Indicators 2008-2012

### Nativity

Percent of Low Birth Weight Infants	7.0
Percent of Mothers Receiving Care in 1st Trimester	69.8
● Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	15.4
Percent of Births Less Than 37 Wks. of Gestation	8.6
○ Average Age of Mother	27.6
Teenage Birth Rate <sup>2</sup>	16.4
Percent White Births	84.0
Percent American Indian Births	4.9
● Percent Unmarried	36.6
● Percent WIC births	31.5
Percent Breastfeeding at discharge	74.1
○ Percent Payment-Private Insurance	63.5
● Percent Payment-Medicaid	32.8
Percent C-Section	26.8

●Denotes a health status indicator which is significantly lower than the state average.  
○Denotes a health status indicator which is significantly higher than the state average.

\*Due to lack of comparable data from all 50 states, significant differences cannot be calculated.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

All Causes	698.6
● Heart Disease	145.4
Malignant Neoplasms (cancer)	173.9
Trachea, Bronchus, & Lung	49.1
Colon, Rectum, & Anus	15.5
Female Breast	19.6
Pancreas	10.1
Cerebrovascular Disease	41.0
Chronic Lower Respiratory Diseases	50.0
● Accidents	33.0
● Motor Vehicle Accidents	9.2
Alzheimer's Disease	42.2
● Diabetes Mellitus	17.4
Influenza & Pneumonia	13.3
Intentional Self-Harm (suicide)	14.1
Chronic Liver Disease and Cirrhosis	8.7
Infant Mortality	5.67

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Malignant Neoplasms (cancer)	1,520
2. Heart Disease	1,328
3. Chronic Lower Respiratory Diseases	439
4. Alzheimer's Disease	393
5. Cerebrovascular Diseases	367
6. Accidents	297
7. Diabetes	157
8. Suicide	125
9. Influenza and Pneumonia	123
10. Essential (primary) Hypertension and Hypertensive Renal Disease	86
Percent of Deaths due to tobacco use	18.0
Median age at death	79

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Moody County

## Demographic Information



Moody County is located on the Minnesota border and averages 12.5 persons per square mile.

### 2012 Population Information

Subject	Number	Percent
Total population	6,446	100.0
White	5,288	82.0
Black or African American	54	0.8
American Indian & Alaska Native	854	13.2
Asian	81	1.3
Native Hawaiian & Other Pacific Islander	0	0.0
Two or More Races	169	2.6
Under 5 years	442	6.9
Under 18 years	1,651	25.6
65 years and over	1,041	16.1

Source: United States Census Bureau, 2012 Population Estimates

## Health Status Indicators 2008-2012

### Nativity

Percent of Low Birth Weight Infants	5.9
Percent of Mothers Receiving Care in 1st Trimester	61.7
Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	20.7
Percent of Births Less Than 37 Wks. of Gestation	9.9
Average Age of Mother	27.3
Teenage Birth Rate <sup>2</sup>	14.2
Percent White Births	78.2
Percent American Indian Births	16.2
Percent Unmarried	36.2
Percent WIC births	40.1
Percent Breastfeeding at discharge	75.8
Percent Payment-Private Insurance	62.0
Percent Payment-Medicaid	32.1
Percent C-Section	25.2

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

\*Due to lack of comparable data from all 50 states, significant differences cannot be calculated.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

● All Causes	622.4
Heart Disease	177.0
Malignant Neoplasms (cancer)	162.2
Trachea, Bronchus, & Lung	34.9
Colon, Rectum, & Anus	20.7
Female Breast	26.7
Pancreas	12.2
Cerebrovascular Disease	28.6
● Chronic Lower Respiratory Diseases	22.7
Accidents	40.1
Motor Vehicle Accidents	13.0
Alzheimer's Disease	59.6
Diabetes Mellitus	27.4
Influenza & Pneumonia	LNE
Intentional Self-Harm (suicide)	16.0
Chronic Liver Disease and Cirrhosis	9.6
Infant Mortality	LNE

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	75
2. Malignant Neoplasms (cancer)	67
3. Alzheimer's Disease	28
4. Accidents	15
T5. Cerebrovascular Diseases	11
T5. Diabetes	11
7. Chronic Lower Respiratory Diseases	9
8. Suicide	6
Percent of Deaths due to tobacco use	21.6
Median age at death	80

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Pennington County

## Demographic Information



Pennington County is located on the Wyoming border and averages 36.4 persons per square mile.

### 2012 Population Information

Subject	Number	Percent
Total population	104,347	100.0
White	87,714	84.1
Black or African American	1,590	1.5
American Indian & Alaska Native	10,172	9.7
Asian	1,187	1.1
Native Hawaiian & Other Pacific Islander	98	0.1
Two or More Races	3,586	3.4
Under 5 years	7,410	7.1
Under 18 years	25,200	24.2
65 years and over	14,968	14.3

Source: United States Census Bureau, 2012 Population Estimates

## Health Status Indicators 2008-2012

### Nativity

Percent of Low Birth Weight Infants	6.8
Percent of Mothers Receiving Care in 1st Trimester	71.9
○ Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	19.2
Percent of Births Less Than 37 Wks. of Gestation	8.1
● Average Age of Mother	26.5
Teenage Birth Rate <sup>2</sup>	18.6
Percent White Births	72.0
Percent American Indian Births	19.0
○ Percent Unmarried	43.2
○ Percent WIC births	46.2
○ Percent Breastfeeding at discharge	79.7
● Percent Payment-Private Insurance	37.6
○ Percent Payment-Medicaid	42.9
● Percent C-Section	19.9

●Denotes a health status indicator which is significantly lower than the state average.  
○Denotes a health status indicator which is significantly higher than the state average.

\*Due to lack of comparable data from all 50 states, significant differences cannot be calculated.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

● All Causes	655.4
Heart Disease	145.6
Malignant Neoplasms (cancer)	160.2
Trachea, Bronchus, & Lung	45.0
● Colon, Rectum, & Anus	11.4
Female Breast	23.9
Pancreas	11.0
● Cerebrovascular Disease	29.2
Chronic Lower Respiratory Diseases	43.8
Accidents	45.4
● Motor Vehicle Accidents	9.8
● Alzheimer's Disease	29.4
● Diabetes Mellitus	11.1
Influenza & Pneumonia	16.5
Intentional Self-Harm (suicide)	19.2
Chronic Liver Disease and Cirrhosis	9.2
Infant Mortality	7.39

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Malignant Neoplasms (cancer)	916
2. Heart Disease	854
3. Chronic Lower Respiratory Diseases	254
4. Accidents	242
5. Alzheimer's Disease	177
6. Cerebrovascular Diseases	169
7. Unspecified Dementia	103
8. Suicide	99
9. Influenza and Pneumonia	97
10. Diabetes	64
Percent of Deaths due to tobacco use	16.2
Median age at death	78

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Perkins County

## Demographic Information



Perkins County is located in northwestern South Dakota and averages 1.0 persons per square mile.

### 2012 Population Information

Subject	Number	Percent
Total population	3,037	100.0
White	2,939	96.8
Black or African American	10	0.3
American Indian & Alaska Native	48	1.6
Asian	8	0.3
Native Hawaiian & Other Pacific Islander	2	0.1
Two or More Races	30	1.0
Under 5 years	153	5.0
Under 18 years	628	20.7
65 years and over	691	22.8

Source: United States Census Bureau, 2012 Population Estimates

## Health Status Indicators 2008-2012

### Natality

Percent of Low Birth Weight Infants	4.4
Percent of Mothers Receiving Care in 1st Trimester	62.3
● Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	11.3
Percent of Births Less Than 37 Wks. of Gestation	5.7
○ Average Age of Mother	28.5
Teenage Birth Rate <sup>2</sup>	LNE
Percent White Births	96.8
Percent American Indian Births	2.5
● Percent Unmarried	15.7
Percent WIC births	40.5
Percent Breastfeeding at discharge	80.8
Percent Payment-Private Insurance	61.8
● Percent Payment-Medicaid	21.0
Percent C-Section	32.1

●Denotes a health status indicator which is significantly lower than the state average.  
○Denotes a health status indicator which is significantly higher than the state average.

\*Due to lack of comparable data from all 50 states, significant differences cannot be calculated.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

All Causes	779.2
Heart Disease	137.3
Malignant Neoplasms (cancer)	187.5
Trachea, Bronchus, & Lung	41.9
Colon, Rectum, & Anus	23.9
Female Breast	41.4
Pancreas	9.7
Cerebrovascular Disease	30.6
Chronic Lower Respiratory Diseases	47.3
○ Accidents	99.1
Motor Vehicle Accidents	30.0
Alzheimer's Disease	28.2
Diabetes Mellitus	22.9
Influenza & Pneumonia	17.5
Intentional Self-Harm (suicide)	16.7
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	LNE

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Malignant Neoplasms (cancer)	51
2. Heart Disease	46
3. Accidents	19
4. Chronic Lower Respiratory Diseases	15
5. Cerebrovascular Diseases	12
6. Alzheimer's Disease	10
7. Diabetes	8
Percent of Deaths due to tobacco use	16.5
Median age at death	83

●Denotes a health status indicator which is significantly lower than the state average.  
○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Potter County

## Demographic Information



Potter County is located in north central South Dakota and averages 2.7 persons per square mile.

### 2012 Population Information

Subject	Number	Percent
Total population	2,359	100.0
White	2,293	97.2
Black or African American	9	0.4
American Indian & Alaska Native	26	1.1
Asian	8	0.3
Native Hawaiian & Other Pacific Islander	1	0.0
Two or More Races	22	0.9
Under 5 years	132	5.6
Under 18 years	468	19.8
65 years and over	613	26.0

Source: United States Census Bureau, 2012 Population Estimates

## Health Status Indicators 2008-2012

### Nativity

Percent of Low Birth Weight Infants	3.3
Percent of Mothers Receiving Care in 1st Trimester	70.2
Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	11.6
Percent of Births Less Than 37 Wks. of Gestation	5.0
Average Age of Mother	27.8
Teenage Birth Rate <sup>2</sup>	LNE
Percent White Births	94.2
Percent American Indian Births	4.1
● Percent Unmarried	24.0
● Percent WIC births	24.2
Percent Breastfeeding at discharge	70.8
○ Percent Payment-Private Insurance	75.2
● Percent Payment-Medicaid	22.3
Percent C-Section	28.1

●Denotes a health status indicator which is significantly lower than the state average.  
○Denotes a health status indicator which is significantly higher than the state average.

\*Due to lack of comparable data from all 50 states, significant differences cannot be calculated.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

All Causes	748.9
Heart Disease	142.6
Malignant Neoplasms (cancer)	204.7
Trachea, Bronchus, & Lung	63.5
Colon, Rectum, & Anus	23.2
Female Breast	37.9
Pancreas	LNE
Cerebrovascular Disease	32.4
Chronic Lower Respiratory Diseases	35.3
Accidents	70.3
Motor Vehicle Accidents	LNE
Alzheimer's Disease	33.1
Diabetes Mellitus	28.7
Influenza & Pneumonia	22.5
Intentional Self-Harm (suicide)	13.6
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	LNE

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Malignant Neoplasms (cancer)	48
2. Heart Disease	44
3. Alzheimer's Disease	12
T4. Accidents	11
T4. Chronic Lower Respiratory Diseases	11
T6. Influenza and Pneumonia	9
T6. Cerebrovascular Diseases	9
8. Diabetes	7
9. Unspecified Dementia	6
10. Parkinson's Disease	5
Percent of Deaths due to tobacco use	18.4
Median age at death	84

●Denotes a health status indicator which is significantly lower than the state average.  
○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Roberts County

## Demographic Information



Roberts County is located in the extreme northeastern corner of the state and averages 9.2 persons per square mile.

### 2012 Population Information

Subject	Number	Percent
Total population	10,303	100.0
White	6,228	60.4
Black or African American	24	0.2
American Indian & Alaska Native	3,697	35.9
Asian	35	0.3
Native Hawaiian & Other Pacific Islander	1	0.0
Two or More Races	318	3.1
Under 5 years	878	8.5
Under 18 years	2,935	28.5
65 years and over	1,826	17.7

Source: United States Census Bureau, 2012 Population Estimates

## Health Status Indicators 2008-2012

### Natality

Percent of Low Birth Weight Infants	6.6
● Percent of Mothers Receiving Care in 1st Trimester	51.7
○ Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	28.9
Percent of Births Less Than 37 Wks. of Gestation	8.9
● Average Age of Mother	25.1
○ Teenage Birth Rate <sup>2</sup>	38.0
Percent White Births	37.4
Percent American Indian Births	55.2
○ Percent Unmarried	61.6
○ Percent WIC births	61.1
● Percent Breastfeeding at discharge	66.9
● Percent Payment-Private Insurance	32.4
○ Percent Payment-Medicaid	60.6
Percent C-Section	24.1

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

\*Due to lack of comparable data from all 50 states, significant differences cannot be calculated.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

○ All Causes	862.1
○ Heart Disease	215.7
Malignant Neoplasms (cancer)	198.6
Trachea, Bronchus, & Lung	43.4
Colon, Rectum, & Anus	27.7
Female Breast	26.8
Pancreas	11.8
Cerebrovascular Disease	49.4
Chronic Lower Respiratory Diseases	34.2
○ Accidents	76.5
○ Motor Vehicle Accidents	40.0
Alzheimer's Disease	25.0
Diabetes Mellitus	32.4
Influenza & Pneumonia	11.9
Intentional Self-Harm (suicide)	20.9
Chronic Liver Disease and Cirrhosis	23.7
Infant Mortality	15.06

### Leading Causes of Death

	Total Deaths
1. Heart Disease	161
2. Malignant Neoplasms (cancer)	133
3. Accidents	39
4. Cerebrovascular Diseases	38
5. Chronic Lower Respiratory Diseases	25
T6. Alzheimer's Disease	20
T6. Diabetes	20
8. Chronic Liver Disease and Cirrhosis	11
9. Suicide	9
T10. Influenza and Pneumonia	8
T10. Septicemia	8
Percent of Deaths due to tobacco use	16.7
Median age at death	80

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Sanborn County

## Demographic Information



Sanborn County is located in east central South Dakota and averages 4.1 persons per square mile.

### 2012 Population Information

Subject	Number	Percent
Total population	2,324	100.0
White	2,267	97.5
Black or African American	2	0.1
American Indian & Alaska Native	12	0.5
Asian	6	0.3
Native Hawaiian & Other Pacific Islander	1	0.0
Two or More Races	36	1.5
Under 5 years	127	5.5
Under 18 years	497	21.4
65 years and over	455	19.6

Source: United States Census Bureau, 2012 Population Estimates

## Health Status Indicators 2008-2012

### Nativity

Percent of Low Birth Weight Infants	6.7
Percent of Mothers Receiving Care in 1st Trimester	69.2
Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	17.0
Percent of Births Less Than 37 Wks. of Gestation	5.9
○ Average Age of Mother	28.3
Teenage Birth Rate <sup>2</sup>	LNE
Percent White Births	99.3
Percent American Indian Births	LNE
● Percent Unmarried	26.7
● Percent WIC births	29.3
Percent Breastfeeding at discharge	79.1
○ Percent Payment-Private Insurance	73.1
● Percent Payment-Medicaid	22.4
Percent C-Section	26.7

●Denotes a health status indicator which is significantly lower than the state average.  
○Denotes a health status indicator which is significantly higher than the state average.

\*Due to lack of comparable data from all 50 states, significant differences cannot be calculated.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

All Causes	648.1
Heart Disease	135.2
● Malignant Neoplasms (cancer)	104.8
● Trachea, Bronchus, & Lung	19.8
Colon, Rectum, & Anus	27.9
Female Breast	LNE
Pancreas	LNE
Cerebrovascular Disease	46.4
Chronic Lower Respiratory Diseases	29.7
Accidents	74.7
Motor Vehicle Accidents	48.3
Alzheimer's Disease	65.5
Diabetes Mellitus	15.4
Influenza & Pneumonia	17.4
Intentional Self-Harm (suicide)	LNE
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	LNE

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	27
2. Malignant Neoplasms (cancer)	19
3. Alzheimer's Disease	15
4. Accidents	11
5. Cerebrovascular Diseases	10
6. Chronic Lower Respiratory Diseases	6
Percent of Deaths due to tobacco use	17.2
Median age at death	82

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Shannon County

## Demographic Information



Shannon County is located in the southwestern part of the state, along the Nebraska border and averages 6.5 persons per square mile.

### 2012 Population Information

Subject	Number	Percent
Total population	14,059	100.0
White	752	5.3
Black or African American	46	0.3
American Indian & Alaska Native	12,971	92.3
Asian	71	0.5
Native Hawaiian & Other Pacific Islander	6	0.0
Two or More Races	213	1.5
Under 5 years	1,634	11.6
Under 18 years	5,434	38.7
65 years and over	895	6.4

Source: United States Census Bureau, 2012 Population Estimates

## Health Status Indicators 2008-2012

### Nativity

Percent of Low Birth Weight Infants	7.2
● Percent of Mothers Receiving Care in 1st Trimester	51.9
○ Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	23.3
○ Percent of Births Less Than 37 Wks. of Gestation	10.5
● Average Age of Mother	24.2
○ Teenage Birth Rate <sup>2</sup>	65.7
Percent White Births	2.1
Percent American Indian Births	97.1
○ Percent Unmarried	86.7
○ Percent WIC births	75.4
● Percent Breastfeeding at discharge	51.8
● Percent Payment-Private Insurance	3.7
○ Percent Payment-Medicaid	65.3
● Percent C-Section	23.0

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

\*Due to lack of comparable data from all 50 states, significant differences cannot be calculated.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

○ All Causes	1375.2
Heart Disease	198.5
Malignant Neoplasms (cancer)	216.8
Trachea, Bronchus, & Lung	51.0
Colon, Rectum, & Anus	32.5
Female Breast	14.9
Pancreas	16.1
Cerebrovascular Disease	56.4
○ Chronic Lower Respiratory Diseases	90.6
○ Accidents	134.8
○ Motor Vehicle Accidents	64.5
● Alzheimer's Disease	10.7
○ Diabetes Mellitus	98.0
○ Influenza & Pneumonia	54.1
○ Intentional Self-Harm (suicide)	35.4
○ Chronic Liver Disease and Cirrhosis	96.1
○ Infant Mortality	18.03

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Accidents	85
2. Malignant Neoplasms (cancer)	74
3. Heart Disease	70
4. Chronic Liver Disease and Cirrhosis	49
5. Diabetes	38
6. Suicide	28
7. Chronic Lower Respiratory Diseases	27
8. Influenza and Pneumonia	20
9. Cerebrovascular Diseases	18
10. Septicemia	13
Percent of Deaths due to tobacco use	11.2
Median age at death	54

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Spink County

## Demographic Information



Spink County is located in the center of eastern South Dakota and averages 4.3 persons per square mile.

### 2012 Population Information

Subject	Number	Percent
Total population	6,611	100.0
White	6,428	97.2
Black or African American	32	0.5
American Indian & Alaska Native	85	1.3
Asian	7	0.1
Native Hawaiian & Other Pacific Islander	1	0.0
Two or More Races	58	0.9
Under 5 years	383	5.8
Under 18 years	1,614	24.4
65 years and over	1,277	19.3

Source: United States Census Bureau, 2012 Population Estimates

## Health Status Indicators 2008-2012

### Nativity

○ Percent of Low Birth Weight Infants	10.2
Percent of Mothers Receiving Care in 1st Trimester	67.2
Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	16.7
Percent of Births Less Than 37 Wks. of Gestation	10.7
Average Age of Mother	27.4
● Teenage Birth Rate <sup>2</sup>	6.8
Percent White Births	97.2
Percent American Indian Births	0.8
● Percent Unmarried	26.1
Percent WIC births	34.2
Percent Breastfeeding at discharge	77.8
○ Percent Payment-Private Insurance	69.9
● Percent Payment-Medicaid	24.4
Percent C-Section	25.4

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

\*Due to lack of comparable data from all 50 states, significant differences cannot be calculated.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

All Causes	732.0
Heart Disease	167.3
Malignant Neoplasms (cancer)	136.7
Trachea, Bronchus, & Lung	35.7
Colon, Rectum, & Anus	20.1
Female Breast	10.7
Pancreas	7.2
Cerebrovascular Disease	34.3
Chronic Lower Respiratory Diseases	59.8
Accidents	67.0
Motor Vehicle Accidents	20.8
Alzheimer's Disease	54.1
Diabetes Mellitus	25.6
Influenza & Pneumonia	22.6
Intentional Self-Harm (suicide)	9.6
Chronic Liver Disease and Cirrhosis	11.8
Infant Mortality	7.61

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	96
2. Malignant Neoplasms (cancer)	74
3. Alzheimer's Disease	37
4. Chronic Lower Respiratory Diseases	33
5. Accidents	24
6. Cerebrovascular Diseases	20
7. Influenza and Pneumonia	15
8. Diabetes	11
T9. Other Interstitial Pulmonary Diseases with Fibrosis	5
T9. Chronic Liver Disease and Cirrhosis	5
Percent of Deaths due to tobacco use	13.7
Median age at death	82

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Stanley County

## Demographic Information



Stanley County is located in the center of the state and averages 2.1 persons per square mile.

### 2012 Population Information

Subject	Number	Percent
Total population	2,969	100.0
White	2,641	89.0
Black or African American	18	0.6
American Indian & Alaska Native	224	7.5
Asian	8	0.3
Native Hawaiian & Other Pacific Islander	0	0.0
Two or More Races	78	2.6
Under 5 years	172	5.8
Under 18 years	675	22.7
65 years and over	487	16.4

Source: United States Census Bureau, 2012 Population Estimates

## Health Status Indicators 2008-2012

### Nativity

Percent of Low Birth Weight Infants	6.7
Percent of Mothers Receiving Care in 1st Trimester	68.9
Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	23.1
Percent of Births Less Than 37 Wks. of Gestation	7.7
Average Age of Mother	27.7
Teenage Birth Rate <sup>2</sup>	16.6
Percent White Births	84.6
Percent American Indian Births	11.8
Percent Unmarried	41.5
Percent WIC births	48.2
Percent Breastfeeding at discharge	73.7
Percent Payment-Private Insurance	60.5
Percent Payment-Medicaid	37.9
○ Percent C-Section	42.6

●Denotes a health status indicator which is significantly lower than the state average.  
○Denotes a health status indicator which is significantly higher than the state average.

\*Due to lack of comparable data from all 50 states, significant differences cannot be calculated.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

All Causes	584.7
● Heart Disease	99.0
Malignant Neoplasms (cancer)	196.6
Trachea, Bronchus, & Lung	67.5
Colon, Rectum, & Anus	18.3
Female Breast	47.3
Pancreas	LNE
Cerebrovascular Disease	40.7
Chronic Lower Respiratory Diseases	52.9
Accidents	39.5
Motor Vehicle Accidents	18.2
Alzheimer's Disease	LNE
Diabetes Mellitus	16.6
Influenza & Pneumonia	LNE
Intentional Self-Harm (suicide)	22.0
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	LNE

### Leading Causes of Death

	Total Deaths
1. Malignant Neoplasms (cancer)	35
2. Heart Disease	16
3. Chronic Lower Respiratory Diseases	10
T4. Accidents	6
T4. Cerebrovascular Diseases	6
Percent of Deaths due to tobacco use	35.4
Median age at death	69

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Sully County

## Demographic Information



Sully County is located in the central region of the state and averages 1.4 persons per square mile.

### 2012 Population Information

Subject	Number	Percent
Total population	1,427	100.0
White	1,364	95.6
Black or African American	10	0.7
American Indian & Alaska Native	22	1.5
Asian	0	0.0
Native Hawaiian & Other Pacific Islander	0	0.0
Two or More Races	31	2.2
Under 5 years	76	5.3
Under 18 years	295	20.7
65 years and over	269	18.9

Source: United States Census Bureau, 2012 Population Estimates

## Health Status Indicators 2008-2012

### Nativity

Percent of Low Birth Weight Infants	10.0
Percent of Mothers Receiving Care in 1st Trimester	60.8
Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	12.7
Percent of Births Less Than 37 Wks. of Gestation	11.3
Average Age of Mother	28.3
Teenage Birth Rate <sup>2</sup>	LNE
Percent White Births	87.5
Percent American Indian Births	10.0
● Percent Unmarried	23.8
● Percent WIC births	26.6
Percent Breastfeeding at discharge	73.4
Percent Payment-Private Insurance	67.5
Percent Payment-Medicaid	28.8
Percent C-Section	32.5

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

\*Due to lack of comparable data from all 50 states, significant differences cannot be calculated.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

● All Causes	503.3
Heart Disease	107.8
Malignant Neoplasms (cancer)	173.8
Trachea, Bronchus, & Lung	52.1
Colon, Rectum, & Anus	LNE
Female Breast	LNE
Pancreas	31.6
Cerebrovascular Disease	LNE
Chronic Lower Respiratory Diseases	LNE
Accidents	30.2
Motor Vehicle Accidents	LNE
Alzheimer's Disease	LNE
Diabetes Mellitus	LNE
Influenza & Pneumonia	LNE
Intentional Self-Harm (suicide)	LNE
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	LNE

### Leading Causes of Death

	Total Deaths
1. Malignant Neoplasms (cancer)	17
2. Heart Disease	10
Percent of Deaths due to tobacco use	27.1
Median age at death	73

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Todd County

## Demographic Information



Todd County is located in south central South Dakota, along the Nebraska border and averages 6.9 persons per square mile.

### 2012 Population Information

Subject	Number	Percent
Total population	9,942	100.0
White	1,049	10.6
Black or African American	26	0.3
American Indian & Alaska Native	8,630	86.8
Asian	31	0.3
Native Hawaiian & Other Pacific Islander	0	0.0
Two or More Races	206	2.1
Under 5 years	1,292	13.0
Under 18 years	4,034	40.6
65 years and over	663	6.7

Source: United States Census Bureau, 2012 Population Estimates

## Health Status Indicators 2008-2012

### Nativity

Percent of Low Birth Weight Infants	7.4
● Percent of Mothers Receiving Care in 1st Trimester	37.6
○ Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	27.2
○ Percent of Births Less Than 37 Wks. of Gestation	10.1
● Average Age of Mother	24.6
○ Teenage Birth Rate <sup>2</sup>	69.4
Percent White Births	4.4
Percent American Indian Births	92.6
○ Percent Unmarried	83.1
○ Percent WIC births	73.6
● Percent Breastfeeding at discharge	57.3
● Percent Payment-Private Insurance	4.5
○ Percent Payment-Medicaid	45.0
○ Percent C-Section	34.4

●Denotes a health status indicator which is significantly lower than the state average.  
○Denotes a health status indicator which is significantly higher than the state average.

\*Due to lack of comparable data from all 50 states, significant differences cannot be calculated.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

○ All Causes	1,265.6
Heart Disease	198.9
○ Malignant Neoplasms (cancer)	240.9
Trachea, Bronchus, & Lung	80.4
Colon, Rectum, & Anus	28.7
Female Breast	LNE
Pancreas	10.2
Cerebrovascular Disease	72.6
Chronic Lower Respiratory Diseases	75.4
○ Accidents	140.3
○ Motor Vehicle Accidents	78.3
● Alzheimer's Disease	13.8
○ Diabetes Mellitus	85.4
Influenza & Pneumonia	21.9
○ Intentional Self-Harm (suicide)	61.5
○ Chronic Liver Disease and Cirrhosis	61.0
Infant Mortality	13.43

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Malignant Neoplasms (cancer)	63
2. Heart Disease	62
3. Accidents	58
4. Suicide	30
5. Diabetes	28
6. Chronic Liver Disease and Cirrhosis	21
7. Chronic Lower Respiratory Diseases	20
8. Cerebrovascular Diseases	18
9. Nephritis, Nephrotic Syndrome, and Nephrosis	10
10. Assault (homicide)	9
Percent of Deaths due to tobacco use	17.1
Median age at death	59

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Tripp County

## Demographic Information



Tripp County is located along the Nebraska border in south central South Dakota and averages 3.5 persons per square mile.

### 2012 Population Information

Subject	Number	Percent
Total population	5,485	100.0
White	4,577	83.4
Black or African American	19	0.3
American Indian & Alaska Native	751	13.7
Asian	16	0.3
Native Hawaiian & Other Pacific Islander	1	0.0
Two or More Races	121	2.2
Under 5 years	329	6.0
Under 18 years	1,243	22.7
65 years and over	1,184	21.6

Source: United States Census Bureau, 2012 Population Estimates

## Health Status Indicators 2008-2012

### Natality

Percent of Low Birth Weight Infants	8.4
Percent of Mothers Receiving Care in 1st Trimester	61.4
○ Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	25.9
Percent of Births Less Than 37 Wks. of Gestation	8.4
Average Age of Mother	26.7
Teenage Birth Rate <sup>2</sup>	20.3
Percent White Births	61.6
Percent American Indian Births	33.0
○ Percent Unmarried	49.2
○ Percent WIC births	54.7
Percent Breastfeeding at discharge	66.2
● Percent Payment-Private Insurance	39.5
○ Percent Payment-Medicaid	52.4
○ Percent C-Section	35.4

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

\*Due to lack of comparable data from all 50 states, significant differences cannot be calculated.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

All Causes	733.7
Heart Disease	127.3
Malignant Neoplasms (cancer)	148.5
● Trachea, Bronchus, & Lung	25.2
Colon, Rectum, & Anus	19.9
Female Breast	20.6
Pancreas	8.1
Cerebrovascular Disease	43.2
Chronic Lower Respiratory Diseases	39.6
Accidents	49.1
Motor Vehicle Accidents	21.3
Alzheimer's Disease	38.6
Diabetes Mellitus	21.4
Influenza & Pneumonia	19.8
Intentional Self-Harm (suicide)	14.7
Chronic Liver Disease and Cirrhosis	23.0
Infant Mortality	9.01

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	100
2. Malignant Neoplasms (cancer)	68
3. Alzheimer's Disease	27
4. Cerebrovascular Diseases	23
5. Chronic Lower Respiratory Diseases	20
6. Accidents	18
7. Influenza and Pneumonia	12
8. Diabetes	11
9. Chronic Liver Disease and Cirrhosis	7
T10. Essential (primary) Hypertension and Hypertensive Renal Disease	5
T10. Parkinson's Disease	5
Percent of Deaths due to tobacco use	15.1
Median age at death	82

●Denotes a health status indicator which is significantly lower than the state average.


○Denotes a health status indicator which is significantly higher than the state average.

<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.


Source: South Dakota Department of Health, Office of Health Statistics

# Turner County

Demographic Information	Health Status Indicators 2008-2012																																																																																																																										
<div style="text-align: center;">  </div> <p>Turner County is located in southeastern South Dakota and averages 13.5 persons per square mile.</p> <p style="text-align: center;"><b>2012 Population Information</b></p> <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="text-align: left;">Subject</th> <th style="text-align: right;">Number</th> <th style="text-align: right;">Percent</th> </tr> </thead> <tbody> <tr> <td>Total population</td> <td style="text-align: right;">8,308</td> <td style="text-align: right;">100.0</td> </tr> <tr> <td>White</td> <td style="text-align: right;">8,102</td> <td style="text-align: right;">97.5</td> </tr> <tr> <td>Black or African American</td> <td style="text-align: right;">39</td> <td style="text-align: right;">0.5</td> </tr> <tr> <td>American Indian &amp; Alaska Native</td> <td style="text-align: right;">78</td> <td style="text-align: right;">0.9</td> </tr> <tr> <td>Asian</td> <td style="text-align: right;">14</td> <td style="text-align: right;">0.2</td> </tr> <tr> <td>Native Hawaiian &amp; Other Pacific Islander</td> <td style="text-align: right;">2</td> <td style="text-align: right;">0.0</td> </tr> <tr> <td>Two or More Races</td> <td style="text-align: right;">73</td> <td style="text-align: right;">0.9</td> </tr> <tr> <td>Under 5 years</td> <td style="text-align: right;">494</td> <td style="text-align: right;">5.9</td> </tr> <tr> <td>Under 18 years</td> <td style="text-align: right;">1,962</td> <td style="text-align: right;">23.6</td> </tr> <tr> <td>65 years and over</td> <td style="text-align: right;">1,611</td> <td style="text-align: right;">19.4</td> </tr> </tbody> </table> <p>Source: United States Census Bureau, 2012 Population Estimates</p>	Subject	Number	Percent	Total population	8,308	100.0	White	8,102	97.5	Black or African American	39	0.5	American Indian & Alaska Native	78	0.9	Asian	14	0.2	Native Hawaiian & Other Pacific Islander	2	0.0	Two or More Races	73	0.9	Under 5 years	494	5.9	Under 18 years	1,962	23.6	65 years and over	1,611	19.4	<p style="text-align: center;"><b>Nativity</b></p> <table border="1" style="width: 100%; border-collapse: collapse;"> <tbody> <tr> <td>Percent of Low Birth Weight Infants</td> <td style="text-align: right;">6.3</td> </tr> <tr> <td>Percent of Mothers Receiving Care in 1st Trimester</td> <td style="text-align: right;">74.1</td> </tr> <tr> <td>● Percent of Mothers Who Used Tobacco While Pregnant<sup>1</sup></td> <td style="text-align: right;">12.8</td> </tr> <tr> <td>Percent of Births Less Than 37 Wks. of Gestation</td> <td style="text-align: right;">9.8</td> </tr> <tr> <td>○ Average Age of Mother</td> <td style="text-align: right;">28.0</td> </tr> <tr> <td>Teenage Birth Rate<sup>2</sup></td> <td style="text-align: right;">LNE</td> </tr> <tr> <td>Percent White Births</td> <td style="text-align: right;">99.1</td> </tr> <tr> <td>Percent American Indian Births</td> <td style="text-align: right;">LNE</td> </tr> <tr> <td>● Percent Unmarried</td> <td style="text-align: right;">24.7</td> </tr> <tr> <td>● Percent WIC births</td> <td style="text-align: right;">28.7</td> </tr> <tr> <td>Percent Breastfeeding at discharge</td> <td style="text-align: right;">77.9</td> </tr> <tr> <td>○ Percent Payment-Private Insurance</td> <td style="text-align: right;">73.4</td> </tr> <tr> <td>● Percent Payment-Medicaid</td> <td style="text-align: right;">22.4</td> </tr> <tr> <td>Percent C-Section</td> <td style="text-align: right;">30.4</td> </tr> </tbody> </table> <p>●Denotes a health status indicator which is significantly lower than the state average. ○Denotes a health status indicator which is significantly higher than the state average. *Due to lack of comparable data from all 50 states, significant differences cannot be calculated. <sup>1</sup>Data for mothers who used tobacco are self-reported. <sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.</p>	Percent of Low Birth Weight Infants	6.3	Percent of Mothers Receiving Care in 1st Trimester	74.1	● Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	12.8	Percent of Births Less Than 37 Wks. of Gestation	9.8	○ Average Age of Mother	28.0	Teenage Birth Rate <sup>2</sup>	LNE	Percent White Births	99.1	Percent American Indian Births	LNE	● Percent Unmarried	24.7	● Percent WIC births	28.7	Percent Breastfeeding at discharge	77.9	○ Percent Payment-Private Insurance	73.4	● Percent Payment-Medicaid	22.4	Percent C-Section	30.4	<p style="text-align: center;"><b>Mortality<sup>3</sup></b></p> <table border="1" style="width: 100%; border-collapse: collapse;"> <tbody> <tr> <td>All Causes</td> <td style="text-align: right;">680.3</td> </tr> <tr> <td>● Heart Disease</td> <td style="text-align: right;">162.9</td> </tr> <tr> <td>Malignant Neoplasms (cancer)</td> <td style="text-align: right;">148.3</td> </tr> <tr> <td>    Trachea, Bronchus, &amp; Lung</td> <td style="text-align: right;">32.3</td> </tr> <tr> <td>    Colon, Rectum, &amp; Anus</td> <td style="text-align: right;">15.2</td> </tr> <tr> <td>    Female Breast</td> <td style="text-align: right;">20.0</td> </tr> <tr> <td>    Pancreas</td> <td style="text-align: right;">9.0</td> </tr> <tr> <td>Cerebrovascular Disease</td> <td style="text-align: right;">34.4</td> </tr> <tr> <td>Chronic Lower Respiratory Diseases</td> <td style="text-align: right;">47.8</td> </tr> <tr> <td>Accidents</td> <td style="text-align: right;">43.1</td> </tr> <tr> <td>    Motor Vehicle Accidents</td> <td style="text-align: right;">19.3</td> </tr> <tr> <td>○ Alzheimer's Disease</td> <td style="text-align: right;">59.6</td> </tr> <tr> <td>Diabetes Mellitus</td> <td style="text-align: right;">16.5</td> </tr> <tr> <td>Influenza &amp; Pneumonia</td> <td style="text-align: right;">12.8</td> </tr> <tr> <td>Intentional Self-Harm (suicide)</td> <td style="text-align: right;">8.4</td> </tr> <tr> <td>Chronic Liver Disease and Cirrhosis</td> <td style="text-align: right;">LNE</td> </tr> <tr> <td>Infant Mortality</td> <td style="text-align: right;">LNE</td> </tr> </tbody> </table> <p style="text-align: center;"><b>Leading Causes of Death</b></p> <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="text-align: left;">Cause</th> <th style="text-align: right;">Total Deaths</th> </tr> </thead> <tbody> <tr> <td>1. Heart Disease</td> <td style="text-align: right;">120</td> </tr> <tr> <td>2. Malignant Neoplasms (cancer)</td> <td style="text-align: right;">96</td> </tr> <tr> <td>3. Alzheimer's Disease</td> <td style="text-align: right;">52</td> </tr> <tr> <td>4. Chronic Lower Respiratory Diseases</td> <td style="text-align: right;">34</td> </tr> <tr> <td>5. Cerebrovascular Diseases</td> <td style="text-align: right;">25</td> </tr> <tr> <td>6. Accidents</td> <td style="text-align: right;">20</td> </tr> <tr> <td>7. Influenza and Pneumonia</td> <td style="text-align: right;">12</td> </tr> <tr> <td>8. Diabetes</td> <td style="text-align: right;">11</td> </tr> <tr> <td>9. Unspecified Dementia</td> <td style="text-align: right;">8</td> </tr> <tr> <td>10. Nephritis, Nephrotic Syndrome and Nephrosis</td> <td style="text-align: right;">7</td> </tr> <tr> <td>Percent of Deaths due to tobacco use</td> <td style="text-align: right;">15.0</td> </tr> <tr> <td>Median age at death</td> <td style="text-align: right;">83</td> </tr> </tbody> </table> <p>●Denotes a health status indicator which is significantly lower than the state average. ○Denotes a health status indicator which is significantly higher than the state average. <sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.</p> <p>See technical notes for more information. Source: South Dakota Department of Health, Office of Health Statistics</p>	All Causes	680.3	● Heart Disease	162.9	Malignant Neoplasms (cancer)	148.3	Trachea, Bronchus, & Lung	32.3	Colon, Rectum, & Anus	15.2	Female Breast	20.0	Pancreas	9.0	Cerebrovascular Disease	34.4	Chronic Lower Respiratory Diseases	47.8	Accidents	43.1	Motor Vehicle Accidents	19.3	○ Alzheimer's Disease	59.6	Diabetes Mellitus	16.5	Influenza & Pneumonia	12.8	Intentional Self-Harm (suicide)	8.4	Chronic Liver Disease and Cirrhosis	LNE	Infant Mortality	LNE	Cause	Total Deaths	1. Heart Disease	120	2. Malignant Neoplasms (cancer)	96	3. Alzheimer's Disease	52	4. Chronic Lower Respiratory Diseases	34	5. Cerebrovascular Diseases	25	6. Accidents	20	7. Influenza and Pneumonia	12	8. Diabetes	11	9. Unspecified Dementia	8	10. Nephritis, Nephrotic Syndrome and Nephrosis	7	Percent of Deaths due to tobacco use	15.0	Median age at death	83
Subject	Number	Percent																																																																																																																									
Total population	8,308	100.0																																																																																																																									
White	8,102	97.5																																																																																																																									
Black or African American	39	0.5																																																																																																																									
American Indian & Alaska Native	78	0.9																																																																																																																									
Asian	14	0.2																																																																																																																									
Native Hawaiian & Other Pacific Islander	2	0.0																																																																																																																									
Two or More Races	73	0.9																																																																																																																									
Under 5 years	494	5.9																																																																																																																									
Under 18 years	1,962	23.6																																																																																																																									
65 years and over	1,611	19.4																																																																																																																									
Percent of Low Birth Weight Infants	6.3																																																																																																																										
Percent of Mothers Receiving Care in 1st Trimester	74.1																																																																																																																										
● Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	12.8																																																																																																																										
Percent of Births Less Than 37 Wks. of Gestation	9.8																																																																																																																										
○ Average Age of Mother	28.0																																																																																																																										
Teenage Birth Rate <sup>2</sup>	LNE																																																																																																																										
Percent White Births	99.1																																																																																																																										
Percent American Indian Births	LNE																																																																																																																										
● Percent Unmarried	24.7																																																																																																																										
● Percent WIC births	28.7																																																																																																																										
Percent Breastfeeding at discharge	77.9																																																																																																																										
○ Percent Payment-Private Insurance	73.4																																																																																																																										
● Percent Payment-Medicaid	22.4																																																																																																																										
Percent C-Section	30.4																																																																																																																										
All Causes	680.3																																																																																																																										
● Heart Disease	162.9																																																																																																																										
Malignant Neoplasms (cancer)	148.3																																																																																																																										
Trachea, Bronchus, & Lung	32.3																																																																																																																										
Colon, Rectum, & Anus	15.2																																																																																																																										
Female Breast	20.0																																																																																																																										
Pancreas	9.0																																																																																																																										
Cerebrovascular Disease	34.4																																																																																																																										
Chronic Lower Respiratory Diseases	47.8																																																																																																																										
Accidents	43.1																																																																																																																										
Motor Vehicle Accidents	19.3																																																																																																																										
○ Alzheimer's Disease	59.6																																																																																																																										
Diabetes Mellitus	16.5																																																																																																																										
Influenza & Pneumonia	12.8																																																																																																																										
Intentional Self-Harm (suicide)	8.4																																																																																																																										
Chronic Liver Disease and Cirrhosis	LNE																																																																																																																										
Infant Mortality	LNE																																																																																																																										
Cause	Total Deaths																																																																																																																										
1. Heart Disease	120																																																																																																																										
2. Malignant Neoplasms (cancer)	96																																																																																																																										
3. Alzheimer's Disease	52																																																																																																																										
4. Chronic Lower Respiratory Diseases	34																																																																																																																										
5. Cerebrovascular Diseases	25																																																																																																																										
6. Accidents	20																																																																																																																										
7. Influenza and Pneumonia	12																																																																																																																										
8. Diabetes	11																																																																																																																										
9. Unspecified Dementia	8																																																																																																																										
10. Nephritis, Nephrotic Syndrome and Nephrosis	7																																																																																																																										
Percent of Deaths due to tobacco use	15.0																																																																																																																										
Median age at death	83																																																																																																																										

# Union County

## Demographic Information



Union County is located in the southeastern corner of the state and averages 31.3 persons per square mile.

### 2012 Population Information

Subject	Number	Percent
Total population	14,855	100.0
White	14,211	95.7
Black or African American	158	1.1
American Indian & Alaska Native	89	0.6
Asian	179	1.2
Native Hawaiian & Other Pacific Islander	7	0.0
Two or More Races	211	1.4
Under 5 years	908	6.1
Under 18 years	3,738	25.2
65 years and over	2,134	14.4

Source: United States Census Bureau, 2012 Population Estimates

## Health Status Indicators 2008-2012

### Nativity

Percent of Low Birth Weight Infants	7.7
○ Percent of Mothers Receiving Care in 1st Trimester	83.7
● Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	13.3
Percent of Births Less Than 37 Wks. of Gestation	10.3
○ Average Age of Mother	28.6
● Teenage Birth Rate <sup>2</sup>	6.7
Percent White Births	95.9
Percent American Indian Births	0.7
● Percent Unmarried	23.6
● Percent WIC births	20.9
Percent Breastfeeding at discharge	79.5
○ Percent Payment-Private Insurance	74.9
● Percent Payment-Medicaid	21.7
○ Percent C-Section	33.3

●Denotes a health status indicator which is significantly lower than the state average.  
○Denotes a health status indicator which is significantly higher than the state average.

\*Due to lack of comparable data from all 50 states, significant differences cannot be calculated.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

All Causes	654.9
Heart Disease	148.0
Malignant Neoplasms (cancer)	177.4
Trachea, Bronchus, & Lung	52.5
Colon, Rectum, & Anus	18.3
Female Breast	23.3
Pancreas	6.4
Cerebrovascular Disease	29.9
Chronic Lower Respiratory Diseases	41.9
Accidents	40.4
Motor Vehicle Accidents	11.1
Alzheimer's Disease	33.5
Diabetes Mellitus	18.1
Influenza & Pneumonia	13.3
Intentional Self-Harm (suicide)	9.2
● Chronic Liver Disease and Cirrhosis	3.8
Infant Mortality	LNE

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Malignant Neoplasms (cancer)	150
2. Heart Disease	131
3. Chronic Lower Respiratory Diseases	36
4. Accidents	33
5. Alzheimer's Disease	30
6. Cerebrovascular Diseases	27
7. Unspecified Dementia	17
8. Diabetes	15
9. Influenza and Pneumonia	11
10. Suicide	8
Percent of Deaths due to tobacco use	16.1
Median age at death	81

●Denotes a health status indicator which is significantly lower than the state average.  
○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Walworth County

## Demographic Information



Walworth County is located in north central South Dakota, near the North Dakota border and averages 7.7 persons per square mile.

### 2012 Population Information

Subject	Number	Percent
Total population	5,459	100.0
White	4,510	82.6
Black or African American	20	0.4
American Indian & Alaska Native	740	13.6
Asian	19	0.3
Native Hawaiian & Other Pacific Islander	0	0.0
Two or More Races	170	3.1
Under 5 years	330	6.0
Under 18 years	1,183	21.7
65 years and over	1,314	24.1

Source: United States Census Bureau, 2012 Population Estimates

## Health Status Indicators 2008-2012

### Natality

● Percent of Low Birth Weight Infants	3.8
● Percent of Mothers Receiving Care in 1st Trimester	59.5
○ Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	23.5
Percent of Births Less Than 37 Wks. of Gestation	8.2
● Average Age of Mother	25.8
Teenage Birth Rate <sup>2</sup>	23.9
Percent White Births	61.6
Percent American Indian Births	34.6
○ Percent Unmarried	49.8
○ Percent WIC births	49.0
Percent Breastfeeding at discharge	67.8
● Percent Payment-Private Insurance	40.6
○ Percent Payment-Medicaid	51.4
Percent C-Section	23.3

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

\*Due to lack of comparable data from all 50 states, significant differences cannot be calculated.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

All Causes	731.5
Heart Disease	127.7
Malignant Neoplasms (cancer)	138.0
Trachea, Bronchus, & Lung	37.0
Colon, Rectum, & Anus	25.9
● Female Breast	8.6
Pancreas	5.5
Cerebrovascular Disease	40.4
Chronic Lower Respiratory Diseases	50.5
● Accidents	21.8
Motor Vehicle Accidents	LNE
○ Alzheimer's Disease	62.5
Diabetes Mellitus	32.2
Influenza & Pneumonia	33.4
Intentional Self-Harm (suicide)	9.5
Chronic Liver Disease and Cirrhosis	16.6
Infant Mortality	9.46

### Leading Causes of Death

	Total Deaths
1. Heart Disease	73
2. Malignant Neoplasms (cancer)	71
3. Alzheimer's Disease	42
4. Chronic Lower Respiratory Diseases	27
5. Cerebrovascular Diseases	23
6. Diabetes	17
7. Influenza and Pneumonia	13
8. Accidents	7
9. Neoplasms of Uncertain or Unknown Behavior	6
10. Chronic Liver Disease and Cirrhosis	5

Percent of Deaths due to tobacco use	19.0
Median age at death	82

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Yankton County

## Demographic Information



Yankton County is located in southeastern South Dakota on the Nebraska border and averages 43.1 persons per square mile.

## 2012 Population Information

Subject	Number	Percent
Total population	22,603	100.0
White	21,040	93.1
Black or African American	449	2.0
American Indian & Alaska Native	648	2.9
Asian	143	0.6
Native Hawaiian & Other Pacific Islander	10	0.0
Two or More Races	313	1.4
Under 5 years	1,313	5.8
Under 18 years	4,831	21.4
65 years and over	3,739	16.5

Source: United States Census Bureau, 2012 Population Estimates

## Health Status Indicators 2008-2012

### Nativity

Percent of Low Birth Weight Infants	6.1
○ Percent of Mothers Receiving Care in 1st Trimester	82.1
○ Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	21.7
Percent of Births Less Than 37 Wks. of Gestation	7.4
Average Age of Mother	27.0
● Teenage Birth Rate <sup>2</sup>	11.3
Percent White Births	92.0
Percent American Indian Births	4.0
Percent Unmarried	39.9
Percent WIC births	39.7
● Percent Breastfeeding at discharge	68.2
○ Percent Payment-Private Insurance	63.9
Percent Payment-Medicaid	33.0
Percent C-Section	26.1

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

\*Due to lack of comparable data from all 50 states, significant differences cannot be calculated.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

● All Causes	641.8
Heart Disease	145.0
● Malignant Neoplasms (cancer)	141.1
Trachea, Bronchus, & Lung	48.5
Colon, Rectum, & Anus	13.5
Female Breast	15.8
Pancreas	8.2
Cerebrovascular Disease	31.2
Chronic Lower Respiratory Diseases	49.3
Accidents	41.0
Motor Vehicle Accidents	14.4
Alzheimer's Disease	34.7
Diabetes Mellitus	32.5
Influenza & Pneumonia	13.4
Intentional Self-Harm (suicide)	18.2
Chronic Liver Disease and Cirrhosis	6.7
Infant Mortality	4.64

### Leading Causes of Death

	Total Deaths
1. Heart Disease	247
2. Malignant Neoplasms (cancer)	207
3. Chronic Lower Respiratory Diseases	73
4. Alzheimer's Disease	67
5. Accidents	55
6. Cerebrovascular Diseases	54
7. Diabetes	53
8. Influenza and Pneumonia	25
9. Suicide	21
T10. Unspecified Dementia	16
T10. Essential (primary) Hypertension and Hypertensive Renal Disease	16
Percent of Deaths due to tobacco use	20.2
Median age at death	82

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Ziebach County

## Demographic Information



Ziebach County is located in north central South Dakota and averages 1.4 persons per square mile.

### 2012 Population Information

Subject	Number	Percent
Total population	2,869	100.0
White	697	24.3
Black or African American	10	0.3
American Indian & Alaska Native	2,061	71.8
Asian	10	0.3
Native Hawaiian & Other Pacific Islander	1	0.0
Two or More Races	90	3.1
Under 5 years	255	8.9
Under 18 years	1,022	35.6
65 years and over	203	7.1

Source: United States Census Bureau, 2012 Population Estimates

## Health Status Indicators 2008-2012

### Nativity

Percent of Low Birth Weight Infants	8.2
● Percent of Mothers Receiving Care in 1st Trimester	44.6
Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	22.5
Percent of Births Less Than 37 Wks. of Gestation	9.9
● Average Age of Mother	25.5
○ Teenage Birth Rate <sup>2</sup>	42.4
Percent White Births	13.7
Percent American Indian Births	82.0
○ Percent Unmarried	69.5
○ Percent WIC births	83.1
● Percent Breastfeeding at discharge	50.7
● Percent Payment-Private Insurance	13.1
○ Percent Payment-Medicaid	81.2
Percent C-Section	33.5

●Denotes a health status indicator which is significantly lower than the state average.  
○Denotes a health status indicator which is significantly higher than the state average.  
\*Due to lack of comparable data from all 50 states, significant differences cannot be calculated.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

○ All Causes	959.4
Heart Disease	225.7
Malignant Neoplasms (cancer)	138.5
Trachea, Bronchus, & Lung	44.8
Colon, Rectum, & Anus	LNE
Female Breast	LNE
Pancreas	LNE
Cerebrovascular Disease	34.9
Chronic Lower Respiratory Diseases	56.7
○ Accidents	123.7
Motor Vehicle Accidents	60.0
Alzheimer's Disease	LNE
Diabetes Mellitus	71.8
Influenza & Pneumonia	21.1
Intentional Self-Harm (suicide)	LNE
Chronic Liver Disease and Cirrhosis	33.1
Infant Mortality	21.46

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	18
2. Accidents	14
3. Malignant Neoplasms (cancer)	10
4. Diabetes	5
Percent of Deaths due to tobacco use	14.8
Median age at death	71

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics