


South Dakota

Health Status Indicators 2007-2011

Demographic Information



South Dakota is located in the north central portion of the United States and averages 10.7 persons per square mile.

2011 Population Information

Subject	Number	Percent
Total population	824,082	100.0
White	713,655	86.6
Black or African American	11,537	1.4
American Indian & Alaska Native	73,343	8.9
Asian	8,241	1.0
Native Hawaiian & Other Pacific Islander	824	0.1
Two or More Races	16,482	2.0
Under 5 years	59,334	7.2
Under 18 years	203,548	24.7
65 years and over	118,668	14.4

Source: United States Census Bureau, 2011 American Community Survey estimates

Natality

● Percent of Low Birth Weight Infants	6.5
* Percent of Mothers Receiving Care in 1st Trimester	69.4
* Percent of Mothers Who Used Tobacco While Pregnant ¹	18.1
● Percent of Births Less Than 37 Wks. of Gestation	8.5
Average Age of Mother	27.0
Teenage Birth Rate ²	18.1
Percent White Births	78.0
Percent American Indian Births	16.3
Percent Unmarried	38.3
Percent WIC births	38.9
Percent Breastfeeding at discharge	73.1
Percent Payment-Private Insurance	54.4
Percent Payment-Medicaid	35.2
Percent C-Section	26.3

- Denotes a health status indicator which is significantly lower than the national average.
- Denotes a health status indicator which is significantly higher than the national average.
- * Due to lack of comparable data from all 50 states, significant differences cannot be calculated.
- ¹Data for mothers who used tobacco are self-reported.
- ²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

● ALL CAUSES	704.9
● Heart Disease	159.6
● Malignant Neoplasms (cancer)	166.1
● Trachea, Bronchus, & Lung	45.3
Colon, Rectum, & Anus	15.9
Female Breast	21.0
Pancreas	9.9
Cerebrovascular Disease	39.3
○ Chronic Lower Respiratory Diseases	46.6
○ Accidents	42.4
○ Motor Vehicle Accidents	15.7
○ Alzheimer's Disease	34.9
○ Diabetes Mellitus	23.8
Influenza & Pneumonia	15.9
○ Intentional Self-Harm (suicide)	15.3
○ Chronic Liver Disease and Cirrhosis	10.5
Infant Mortality	6.96

Leading Causes of Death

Total Deaths

1. Heart Disease	8,304
2. Cancer	7,980
3. Chronic Lower Respiratory Disease	2,320
4. Cerebrovascular Disease	2,069
5. Alzheimer's Disease	1,972
6. Accidents	1,875
7. Diabetes	1,170
8. Influenza and Pneumonia	854
9. Suicide	616
10. Chronic Liver Disease and Cirrhosis	454

Percent of Deaths due to tobacco use 17.6
Median age at death 80

- Denotes a health status indicator which is significantly lower than the national average.
- Denotes a health status indicator which is significantly higher than the national average.
- ³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.


See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Aurora County

Health Status Indicators 2007-2011

Demographic Information



Aurora County is located in the south central portion of the state and averages 3.8 persons per square mile.

2011 Population Information

Subject	Number	Percent
Total population	2,694	100.0
White	2,597	96.4
Black or African American	11	0.4
American Indian and Alaska Native	43	1.6
Asian	19	0.7
Native Hawaiian & Other Pacific Islander	0	0.0
Two or More Races	24	0.9
Under 5 years	183	6.8
Under 18 years	714	26.5
65 years and over	525	19.5

Source: United States Census Bureau, 2011 American Community Survey estimates

Nativity

Percent of Low Birth Weight Infants	8.9
Percent of Mothers Receiving Care in 1st Trimester	73.8
Percent of Mothers Who Used Tobacco While Pregnant ¹	14.8
Percent of Births Less Than 37 Wks. of Gestation	10.1
○ Average Age of Mother	28.0
Teenage Birth Rate ²	LNE
Percent White Births	99.4
Percent American Indian Births	LNE
● Percent Unmarried	21.9
Percent WIC births	39.6
Percent Breastfeeding at discharge	75.1
○ Percent Payment-Private Insurance	75.1
● Percent Payment-Medicaid	18.9
○ Percent C-Section	36.7

● Denotes a health status indicator which is significantly lower than the state average.

○ Denotes a health status indicator which is significantly higher than the state average.

LNE (Low Number of Events): is used because the rate or percent may be unreliable.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

● ALL CAUSES	589.4
Heart Disease	164.1
Malignant Neoplasms (cancer)	134.5
Trachea, Bronchus, & Lung	49.7
Colon, Rectum, & Anus	16.4
Female Breast	33.8
Pancreas	LNE
Cerebrovascular Disease	26.4
Chronic Lower Respiratory Diseases	54.9
Accidents	28.0
Motor Vehicle Accidents	LNE
Alzheimer's Disease	24.5
Diabetes Mellitus	28.7
Influenza & Pneumonia	13.6
Intentional Self-Harm (suicide)	LNE
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	LNE

Leading Causes of Death

Total Deaths

1. Heart Disease	42
2. Cancer	30
3. Chronic Lower Respiratory Disease	13
4. Alzheimer's Disease	9
T5. Cerebrovascular Disease	7
T5. Accidents	7
T5. Diabetes	7
8. Influenza and Pneumonia	5

Percent of deaths due to tobacco use

Median age at death

● Denotes a health status indicator which is significantly lower than the state average.

○ Denotes a health status indicator which is significantly higher than the state average.


³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Beadle County

Demographic Information



Beadle County is located in the center of eastern South Dakota and averages 13.8 persons per square mile.

2011 Population Information

Subject	Number	Percent
Total population	17,550	100.0
White	16,041	91.4
Black or African American	228	1.3
American Indian and Alaska Native	228	1.3
Asian	772	4.4
Native Hawaiian & Other Pacific Islander	35	0.2
Two or More Races	281	1.6
Under 5 years	1,351	7.7
Under 18 years	4,300	24.5
65 years and over	2,948	16.8

Source: United States Census Bureau, 2011 American Community Survey estimates

Health Status Indicators 2007-2011

Natality

Percent of Low Birth Weight Infants	7.7
● Percent of Mothers Receiving Care in 1st Trimester	58.5
Percent of Mothers Who Used Tobacco While Pregnant ¹	19.9
Percent of Births Less Than 37 Weeks of Gestation	9.3
● Average Age of Mother	26.4
Teenage Birth Rate ²	23.8
Percent White Births	90.3
Percent American Indian Births	1.0
Percent Unmarried	39.4
○ Percent WIC births	48.7
Percent Breastfeeding at discharge	71.8
● Percent Payment-Private Insurance	49.3
Percent Payment-Medicaid	38.7
○ Percent C-Section	32.5

● Denotes a health status indicator which is significantly lower than the state average.
 ○ Denotes a health status indicator which is significantly higher than the state average.
 LNE (Low Number of Events): is used because the rate or percent may be unreliable.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

ALL CAUSES	741.3
Heart Disease	167.8
Malignant Neoplasms (cancer)	178.8
Trachea, Bronchus, & Lung	45.2
Colon, Rectum, & Anus	19.2
Female Breast	27.2
Pancreas	9.8
Cerebrovascular Disease	45.7
Chronic Lower Respiratory Diseases	41.0
Accidents	37.6
Motor Vehicle Accidents	12.5
Alzheimer's Disease	29.3
○ Diabetes Mellitus	40.4
Influenza & Pneumonia	18.9
Intentional Self-Harm (suicide)	8.4
Chronic Liver Disease and Cirrhosis	6.8
Infant Mortality	8.73

Leading Causes of Death

Total Deaths

1. Heart Disease	239
2. Cancer	218
3. Cerebrovascular Disease	67
4. Chronic Lower Respiratory Disease	56
5. Diabetes	52
6. Alzheimer's Disease	48
7. Accidents	33
8. Influenza and Pneumonia	27
9. Urinary Tract Infection, site not specified	15
10. Essential (primary) hypertension and hypertensive renal disease	13

Percent of deaths due to tobacco use 17.8
 Median age at death 82

● Denotes a health status indicator which is significantly lower than the state average.
 ○ Denotes a health status indicator which is significantly higher than the state average.


³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Bennett County

Demographic Information



Bennett County is located on the southern border in western South Dakota and averages 2.9 persons per square mile.

2011 Population Information

Subject	Number	Percent
Total population	3,441	100.0
White	1,197	34.8
Black or African American	7	0.2
American Indian and Alaska Native	2,071	60.2
Asian	17	0.5
Native Hawaiian & Other Pacific Islander	3	0.1
Two or More Races	145	4.2
Under 5 years	358	10.4
Under 18 years	1,180	34.3
65 years and over	379	11.0

Source: United States Census Bureau, 2011 American Community Survey estimates

Health Status Indicators 2007-2011

Nativity

Percent of Low Birth Weight Infants	7.2
● Percent of Mothers Receiving Care in 1st Trimester	41.8
Percent of Mothers Who Used Tobacco While Pregnant ¹	21.9
Percent of Births Less Than 37 Wks. of Gestation	11.0
● Average Age of Mother	25.0
Teenage Birth Rate ²	32.7
Percent White Births	20.5
Percent American Indian Births	74.4
○ Percent Unmarried	72.1
○ Percent WIC births	75.7
● Percent Breastfeeding at discharge	48.6
● Percent Payment-Private Insurance	12.9
○ Percent Payment-Medicaid	65.2
Percent C-Section	24.9

● Denotes a health status indicator which is significantly lower than the state average.
 ○ Denotes a health status indicator which is significantly higher than the state average.
 LNE (Low Number of Events): is used because the rate or percent may be unreliable.
¹Data for mothers who used tobacco are self-reported.
²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

○ ALL CAUSES	971.0
Heart Disease	171.5
Malignant Neoplasms (cancer)	199.2
Trachea, Bronchus, & Lung	44.8
Colon, Rectum, & Anus	37.7
Female Breast	LNE
Pancreas	LNE
Cerebrovascular Disease	43.2
Chronic Lower Respiratory Diseases	47.4
Accidents	70.0
Motor Vehicle Accidents	48.2
Alzheimer's Disease	LNE
Diabetes Mellitus	47.8
○ Influenza & Pneumonia	60.4
Intentional Self-Harm (suicide)	22.2
Chronic Liver Disease and Cirrhosis	36.7
Infant Mortality	15.02

Leading Causes of Death

Total Deaths

1. Cancer	32
2. Heart Disease	29
3. Accidents	12
4. Influenza and Pneumonia	11
T5. Chronic Lower Respiratory Disease	8
T5. Diabetes	8
7. Cerebrovascular Disease	7
8. Chronic Liver Disease and Cirrhosis	5

Percent of deaths due to tobacco use 19.8
 Median age at death 75


● Denotes a health status indicator which is significantly lower than the state average.
 ○ Denotes a health status indicator which is significantly higher than the state average.
³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Bon Homme County

Demographic Information



Bon Homme County is located on the Nebraska border in eastern South Dakota and averages 12.5 persons per square mile.

2011 Population Information

Subject	Number	Percent
Total population	6,983	100.0
White	6,285	90.0
Black or African American	70	1.0
American Indian and Alaska Native	517	7.4
Asian	14	0.2
Native Hawaiian & Other Pacific Islander	0	0.0
Two or More Races	98	1.4
Under 5 years	328	4.7
Under 18 years	1,355	19.0
65 years and over	1,327	19.4

Source: United States Census Bureau, 2011 American Community Survey estimates

Health Status Indicators 2007-2011

Nativity

Percent of Low Birth Weight Infants	6.2
Percent of Mothers Receiving Care in 1st Trimester	79.4
Percent of Mothers Who Used Tobacco While Pregnant ¹	16.1
● Percent of Births Less Than 37 Wks. of Gestation	4.6
○ Average Age of Mother	28.0
Teenage Birth Rate ²	LNE
Percent White Births	95.7
Percent American Indian Births	1.8
● Percent Unmarried	20.9
Percent WIC births	45.2
Percent Breastfeeding at discharge	74.5
○ Percent Payment-Private Insurance	73.2
● Percent Payment-Medicaid	23.4
Percent C-Section	28.6

● Denotes a health status indicator which is significantly lower than the state average.
 ○ Denotes a health status indicator which is significantly higher than the state average.
 LNE (Low Number of Events): is used because the rate or percent may be unreliable.
¹Data for mothers who used tobacco are self-reported.
²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

ALL CAUSES	635.8
Heart Disease	168.8
Malignant Neoplasms (cancer)	158.4
Trachea, Bronchus, & Lung	61.1
Colon, Rectum, & Anus	16.7
Female Breast	15.0
Pancreas	9.6
Cerebrovascular Disease	47.1
Chronic Lower Respiratory Diseases	38.5
Accidents	34.7
Motor Vehicle Accidents	17.5
Alzheimer's Disease	24.0
Diabetes Mellitus	23.4
● Influenza & Pneumonia	7.9
Intentional Self-Harm (suicide)	13.0
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	LNE

Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	109
2. Cancer	90
3. Cerebrovascular Disease	31
4. Chronic Lower Respiratory Disease	25
5. Alzheimer's Disease	21
6. Accidents	18
7. Diabetes	13
8. Neoplasms of uncertain or unknown behavior	10
9. Unspecified dementia	9
T10. Influenza and Pneumonia	5
T10. Essential (primary) hypertension and hypertensive renal disease	5
T10. Parkinson's disease	5

Percent of deaths due to tobacco use 13.0
 Median age at death 84


● Denotes a health status indicator which is significantly lower than the state average.
 ○ Denotes a health status indicator which is significantly higher than the state average.
³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See Technical Notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Brookings County

Demographic Information



Brookings County is located on the eastern border of the state and averages 40.3 persons per square mile.

2011 Population Information

Subject	Number	Percent
Total population	32,226	100.0
White	30,228	93.8
Black or African American	290	0.9
American Indian and Alaska Native	354	1.1
Asian	902	2.8
Native Hawaiian & Other Pacific Islander	32	0.1
Two or More Races	451	1.4
Under 5 years	1,901	5.9
Under 18 years	6,026	18.7
65 years and over	3,255	10.1

Source: United States Census Bureau, 2011 American Community Survey estimates

Health Status Indicators 2007-2011

Natality

- Percent of Low Birth Weight Infants 4.7
- Percent of Mothers Receiving Care in 1st Trimester 75.7
- Percent of Mothers Who Used Tobacco While Pregnant¹ 12.2
- Percent of Births Less Than 37 Wks. of Gestation ... 6.9
- Average Age of Mother 27.7
- Teenage Birth Rate² 8.8
- Percent White Births 90.6
- Percent American Indian Births 1.9
- Percent Unmarried 23.6
- Percent WIC births 28.3
- Percent Breastfeeding at discharge 81.3
- Percent Payment-Private Insurance 73.8
- Percent Payment-Medicaid 20.9
- Percent C-Section 20.6

● Denotes a health status indicator which is significantly lower than the state average.
 ○ Denotes a health status indicator which is significantly higher than the state average.
 LNE (Low Number of Events): is used because the rate or percent may be unreliable.
¹Data for mothers who used tobacco are self-reported.
²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

- ALL CAUSES 634.9
- Heart Disease 136.3
- Malignant Neoplasms (cancer) 152.4
 - Trachea, Bronchus, & Lung 31.9
 - Colon, Rectum, & Anus 15.6
 - Female Breast 19.7
 - Pancreas 9.3
- Cerebrovascular Disease 39.9
- Chronic Lower Respiratory Diseases 36.1
- Accidents 32.0
 - Motor Vehicle Accidents 8.7
- Alzheimer's Disease 36.8
- Diabetes Mellitus 13.3
- Influenza & Pneumonia 16.1
- Intentional Self-Harm (suicide) 6.8
- Chronic Liver Disease and Cirrhosis 2.9
- Infant Mortality 6.75

Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	212
2. Cancer	206
3. Cerebrovascular Disease	64
4. Alzheimer's Disease	61
5. Chronic Lower Respiratory Disease	51
6. Accidents	48
7. Influenza and Pneumonia	26
8. Unspecified Dementia	21
9. Diabetes	19
T10. Suicide	11
T10. Congenital malformations, deformations, and chromosomal abnormalities	11

Percent of deaths due to tobacco use 18.1
 Median age at death 83


● Denotes a health status indicator which is significantly lower than the state average.
 ○ Denotes a health status indicator which is significantly higher than the state average.
³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See Technical Notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Brown County

Demographic Information



Brown County is located on the North Dakota border in the eastern part of the state and averages 21.3 persons per square mile.

2011 Population Information

Subject	Number	Percent
Total population	36,822	100.0
White	34,465	93.6
Black or African American	221	0.6
American Indian and Alaska Native	1,141	3.1
Asian	368	1.0
Native Hawaiian and Other Pacific Islander	37	0.1
Two or More Races	552	1.5
Under 5 years	2,541	6.9
Under 18 years	8,506	23.1
65 years and over	5,855	15.9

Source: United States Census Bureau, 2011 American Community Survey estimates

Health Status Indicators 2007-2011

Natality

Percent of Low Birth Weight Infants	5.9
○ Percent of Mothers Receiving Care in 1st Trimester	77.2
Percent of Mothers Who Used Tobacco While Pregnant ¹	16.2
Percent of Births Less Than 37 Weeks of Gestation	8.1
Average Age of Mother	27.1
● Teenage Birth Rate ²	11.3
Percent White Births	91.3
Percent American Indian Births	5.8
● Percent Unmarried	33.2
● Percent WIC births	32.0
○ Percent Breastfeeding at discharge	77.4
○ Percent Payment-Private Insurance	60.5
● Percent Payment-Medicaid	27.4
Percent C-Section	28.2

● Denotes a health status indicator which is significantly lower than the state average.
 ○ Denotes a health status indicator which is significantly higher than the state average.
 LNE (Low Number of Events): is used because the rate or percent may be unreliable.
¹Data for mothers who used tobacco are self-reported.
²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

ALL CAUSES	685.8
Heart Disease	170.9
Malignant Neoplasms (cancer)	165.5
Trachea, Bronchus, & Lung	44.2
Colon, Rectum, & Anus	16.2
Female Breast	23.9
Pancreas	11.0
Cerebrovascular Disease	38.4
Chronic Lower Respiratory Diseases	40.4
Accidents	29.3
● Motor Vehicle Accidents	8.7
Alzheimer's Disease	37.2
Diabetes Mellitus	20.3
Influenza & Pneumonia	20.2
Intentional Self-Harm (suicide)	10.3
Chronic Liver Disease and Cirrhosis	9.6
Infant Mortality	4.90

Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	481
2. Cancer	390
3. Alzheimer's Disease	112
4. Chronic Lower Respiratory Disease	108
5. Cerebrovascular Disease	107
6. Accidents	62
7. Influenza and Pneumonia	58
8. Diabetes	53
9. Nephritis, nephrotic syndrome, and nephrosis	32
10. Unspecified dementia	27

Percent of deaths due to tobacco use 16.4
 Median age at death 83


● Denotes a health status indicator which is significantly lower than the state average.
 ○ Denotes a health status indicator which is significantly higher than the state average.
³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See Technical Notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Brule County

Demographic Information



Brule County is located in the south central part of the state and averages 6.4 persons per square mile.

2011 Population Information

Subject	Number	Percent
Total population	5,283	100.0
White	4,670	88.4
Black or African American	16	0.3
American Indian and Alaska Native	449	8.5
Asian	11	0.2
Native Hawaiian and Other Pacific Islander	0	0.0
Two or More Races	137	2.6
Under 5 years	365	6.9
Under 18 years	1,337	25.3
65 years and over	888	16.8

Source: United States Census Bureau, 2011 American Community Survey estimates

Health Status Indicators 2007-2011

Natality

Percent of Low Birth Weight Infants	6.2
Percent of Mothers Receiving Care in 1st Trimester	66.4
Percent of Mothers Who Used Tobacco While Pregnant ¹	18.7
Percent of Births Less Than 37 Wks. of Gestation	8.6
Average Age of Mother	27.6
Teenage Birth Rate ²	10.7
Percent White Births	76.4
Percent American Indian Births	20.9
Percent Unmarried	35.3
Percent WIC births	37.7
Percent Breastfeeding at discharge	70.4
o Percent Payment-Private Insurance	63.3
Percent Payment-Medicaid	34.5
Percent C-Section	29.1

Mortality³

ALL CAUSES	652.8
Heart Disease	150.8
Malignant Neoplasms (cancer)	147.8
Trachea, Bronchus, & Lung	37.7
Colon, Rectum, & Anus	20.3
Female Breast	17.3
Pancreas	LNE
• Cerebrovascular Disease	22.1
Chronic Lower Respiratory Diseases	40.5
Accidents	46.9
Motor Vehicle Accidents	9.3
Alzheimer's Disease	32.4
o Diabetes Mellitus	48.9
Influenza & Pneumonia	12.7
Intentional Self-Harm (suicide)	LNE
Chronic Liver Disease and Cirrhosis	10.3
Infant Mortality	13.48

Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	64
2. Cancer	56
T3. Alzheimer's Disease	17
T3. Diabetes	17
5. Chronic Lower Respiratory Diseases	15
6. Accidents	13
7. Cerebrovascular Disease	9
8. Influenza and Pneumonia	6
9. Disorders of lipoprotein metabolism and other lipidemias	5

Percent of deaths due to tobacco use	23.4
Median age at death	82

•Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

LNE (Low Number of Events): is used because the rate or percent may be unreliable.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

•Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.


³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See Technical Notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Buffalo County

Demographic Information



Buffalo County is located in the central portion of the state and averages 4.1 persons per square mile.

2011 Population Information

Subject	Number	Percent
Total population	1,988	100.0
White	350	17.6
Black or African American	6	0.3
American Indian and Alaska Native	1,600	80.5
Asian	4	0.2
Native Hawaiian and Other Pacific Islander	0	0.0
Two or More Races	30	1.5
Under 5 years	249	12.5
Under 18 years	805	40.5
65 years and over	137	6.9

Source: United States Census Bureau, 2011 American Community Survey estimates

Health Status Indicators 2007-2011

Nativity

Percent of Low Birth Weight Infants	6.3
● Percent of Mothers Receiving Care in 1st Trimester	46.2
○ Percent of Mothers Who Used Tobacco While Pregnant ¹	35.7
Percent of Births Less Than 37 Wks. of Gestation	12.0
● Average Age of Mother	25.3
○ Teenage Birth Rate ²	58.0
Percent White Births	13.8
Percent American Indian Births	80.6
○ Percent Unmarried	77.2
○ Percent WIC births	77.6
● Percent Breastfeeding at discharge	44.8
● Percent Payment-Private Insurance	16.4
○ Percent Payment-Medicaid	77.2
Percent C-Section	27.6

● Denotes a health status indicator which is significantly lower than the state average.
 ○ Denotes a health status indicator which is significantly higher than the state average.
 LNE (Low Number of Events): is used because the rate or percent may be unreliable.
¹Data for mothers who used tobacco are self-reported.
²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

○ ALL CAUSES	1329.1
○ Heart Disease	347.7
Malignant Neoplasms (cancer)	277.2
Trachea, Bronchus, & Lung	53.0
Colon, Rectum, & Anus	LNE
Female Breast	LNE
Pancreas	LNE
Cerebrovascular Disease	45.6
Chronic Lower Respiratory Diseases	55.0
○ Accidents	119.1
Motor Vehicle Accidents	57.5
Alzheimer's Disease	LNE
Diabetes Mellitus	60.6
Influenza & Pneumonia	33.1
Intentional Self-Harm (suicide)	63.5
○ Chronic Liver Disease and Cirrhosis	85.5
Infant Mortality	14.93

Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart disease	22
2. Cancer	17
3. Accidents	11
T4. Suicide	6
T4. Chronic Liver Disease and Cirrhosis	6
6. Diabetes	5

Percent of deaths due to tobacco use 30.2
 Median age at death 58


● Denotes a health status indicator which is significantly lower than the state average.
 ○ Denotes a health status indicator which is significantly higher than the state average.
³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See Technical Notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Butte County

Demographic Information



Butte County is located on the western border of the state and averages 4.5 people per square mile.

2011 Population Information

Subject	Number	Percent
Total population	10,259	100.0
White	9,715	94.7
Black or African American	41	0.4
American Indian and Alaska Native	215	2.1
Asian	31	0.3
Native Hawaiian and Other Pacific Islander	10	0.1
Two or More Races	256	2.5
Under 5 years	728	7.1
Under 18 years	2,544	24.8
65 years and over	1,621	15.8

Source: United States Census Bureau, 2011 American Community Survey estimates

Health Status Indicators 2007-2011

Natality

● Percent of Low Birth Weight Infants	4.4
○ Percent of Mothers Receiving Care in 1st Trimester	77.2
○ Percent of Mothers Who Used Tobacco While Pregnant ¹	23.8
Percent of Births Less Than 37 Weeks of Gestation	7.4
Average Age of Mother	26.6
Teenage Birth Rate ²	16.1
Percent White Births	93.2
Percent American Indian Births	1.6
● Percent Unmarried	31.9
○ Percent WIC births	45.4
Percent Breastfeeding at discharge	78.5
Percent Payment-Private Insurance	50.4
○ Percent Payment-Medicaid	40.8
Percent C-Section	23.7

● Denotes a health status indicator which is significantly lower than the state average.
 ○ Denotes a health status indicator which is significantly higher than the state average.
 LNE (Low Number of Events): is used because the rate or percent may be unreliable.
¹Data for mothers who used tobacco are self-reported.
²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

○ ALL CAUSES	853.9
Heart Disease	173.9
○ Malignant Neoplasms (cancer)	224.6
○ Trachea, Bronchus, & Lung	75.8
Colon, Rectum, & Anus	22.8
Female Breast	23.0
Pancreas	13.7
Cerebrovascular Disease	36.5
Chronic Lower Respiratory Diseases	53.3
Accidents	44.7
Motor Vehicle Accidents	23.3
Alzheimer's Disease	28.6
Diabetes Mellitus	25.2
○ Influenza & Pneumonia	34.2
Intentional Self-Harm (suicide)	20.3
Chronic Liver Disease and Cirrhosis	7.4
Infant Mortality	10.16

Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Cancer	141
2. Heart Disease	112
3. Chronic Lower Respiratory Disease	34
4. Cerebrovascular Disease	23
T5. Accidents	22
T5. Influenza and Pneumonia	22
7. Alzheimer's Disease	18
8. Diabetes	16
9. Suicide	10
T10. Unspecified Dementia	9
T10. Essential (primary) hypertension and hypertensive renal disease	9

Percent of deaths due to tobacco use 20.9
 Median age at death 78


● Denotes a health status indicator which is significantly lower than the state average.
 ○ Denotes a health status indicator which is significantly higher than the state average.
³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See Technical Notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Campbell County

Demographic Information



Campbell County is located in the north central portion of the state and averages 2.0 persons per square mile.

2011 Population Information

Subject	Number	Percent
Total population	1,427	100.0
White	1406	98.5
Black or African American	1	0.1
American Indian and Alaska Native	6	0.4
Asian	7	0.5
Native Hawaiian and Other Pacific Islander	0	0.0
Two or More Races	7	0.5
Under 5 years	56	3.9
Under 18 years	247	17.3
65 years and over	375	26.3

Source: United States Census Bureau, 2011 American Community Survey estimates

Health Status Indicators 2007-2011

Nativity

Percent of Low Birth Weight Infants	4.4
Percent of Mothers Receiving Care in 1st Trimester	73.5
Percent of Mothers Who Used Tobacco While Pregnant ¹	LNE
Percent of Births Less Than 37 Weeks of Gestation	10.3
o Average Age of Mother	29.1
Teenage Birth Rate ²	LNE
Percent White Births	98.5
Percent American Indian Births	LNE
• Percent Unmarried	15.9
Percent WIC births	36.2
Percent Breastfeeding at discharge	82.6
Percent Payment-Private Insurance	71.0
• Percent Payment-Medicaid	8.7
Percent C-Section	39.1

•Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

LNE (Low Number of Events): is used because the rate or percent may be unreliable.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

• ALL CAUSES	485.6
Heart Disease	156.0
Malignant Neoplasms (cancer)	107.9
Trachea, Bronchus, & Lung	LNE
Colon, Rectum, & Anus	32.5
Female Breast	LNE
Pancreas	LNE
Cerebrovascular Disease	44.6
Chronic Lower Respiratory Diseases	LNE
Accidents	33.9
Motor Vehicle Accidents	LNE
Alzheimer's Disease	19.1
Diabetes Mellitus	19.9
Influenza & Pneumonia	LNE
Intentional Self-Harm (suicide)	LNE
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	LNE

Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	20
2. Cancer	14
3. Cerebrovascular Disease	7
4. Accidents	5

Percent of deaths due to tobacco use

Median age at death

•Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.


³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See Technical Notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Charles Mix County

Demographic Information



Charles Mix County is located in the south central area of the state and averages 8.3 persons per square mile.

2011 Population Information

Subject	Number	Percent
Total population	9,208	100.0
White	5,994	65.1
Black or African American	18	0.2
American Indian and Alaska Native	2,901	31.5
Asian	18	0.2
Native Hawaiian and Other Pacific Islander	0	0.0
Two or More Races	276	3.0
Under 5 years	792	8.6
Under 18 years	2,735	29.7
65 years and over	1,602	17.4

Source: United States Census Bureau, 2011 American Community Survey estimates

Health Status Indicators 2007-2011

Nativity

Percent of Low Birth Weight Infants	5.0
● Percent of Mothers Receiving Care in 1st Trimester	62.3
○ Percent of Mothers Who Used Tobacco While Pregnant ¹	25.6
Percent of Births Less Than 37 Weeks of Gestation	7.9
● Average Age of Mother	25.6
Teenage Birth Rate ²	24.9
Percent White Births	43.9
Percent American Indian Births	50.9
○ Percent Unmarried	56.2
○ Percent WIC births	62.8
Percent Breastfeeding at discharge	66.8
● Percent Payment-Private Insurance	36.3
○ Percent Payment-Medicaid	56.5
Percent C-Section	24.8

● Denotes a health status indicator which is significantly lower than the state average.

○ Denotes a health status indicator which is significantly higher than the state average.

LNE (Low Number of Events): is used because the rate or percent may be unreliable.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

ALL CAUSES	764.4
Heart Disease	186.8
Malignant Neoplasms (cancer)	180.8
Trachea, Bronchus, & Lung	30.9
Colon, Rectum, & Anus	23.1
Female Breast	9.9
Pancreas	15.7
● Cerebrovascular Disease	23.7
Chronic Lower Respiratory Diseases	49.0
Accidents	58.8
Motor Vehicle Accidents	24.2
Alzheimer's Disease	27.3
○ Diabetes Mellitus	43.7
Influenza & Pneumonia	18.7
Intentional Self-Harm (suicide)	15.4
Chronic Liver Disease and Cirrhosis	22.3
Infant Mortality	7.56

Leading Causes of Death

Total Deaths

1. Heart Disease	133
2. Cancer	115
3. Chronic Lower Respiratory Disease	35
4. Accidents	31
5. Diabetes	27
6. Alzheimer's Disease	22
7. Cerebrovascular Disease	19
8. Influenza and Pneumonia	14
9. Essential (primary) hypertension and hypertensive renal disease	12
10. Chronic Liver Disease and Cirrhosis	11

Percent of deaths due to tobacco use

Median age at death

● Denotes a health status indicator which is significantly lower than the state average.

○ Denotes a health status indicator which is significantly higher than the state average.


³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See Technical Notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Clark County

Demographic Information



Clark County is located in east central South Dakota and averages 3.9 persons per square mile.

2011 Population Information

Subject	Number	Percent
Total population	3,628	100.0
White	3,577	98.6
Black or African American	15	0.4
American Indian and Alaska Native	4	0.1
Asian	4	0.1
Native Hawaiian and Other Pacific Islander	0	0.0
Two or More Races	29	0.8
Under 5 years	268	7.4
Under 18 years	856	23.6
65 years and over	798	22.0

Source: United States Census Bureau, 2011 American Community Survey estimates

Health Status Indicators 2007-2011

Natality

Percent of Low Birth Weight Infants	4.1
• Percent of Mothers Receiving Care in 1st Trimester	57.3
Percent of Mothers Who Used Tobacco While Pregnant ¹	15.0
Percent of Births Less Than 37 Weeks of Gestation	5.8
○ Average Age of Mother	28.5
Teenage Birth Rate ²	LNE
Percent White Births	97.5
Percent American Indian Births	1.2
• Percent Unmarried	19.4
• Percent WIC births	24.0
Percent Breastfeeding at discharge	84.3
○ Percent Payment-Private Insurance	77.3
• Percent Payment-Medicaid	17.4
• Percent C-Section	19.8

• Denotes a health status indicator which is significantly lower than the state average.

○ Denotes a health status indicator which is significantly higher than the state average.

LNE (Low Number of Events): is used because the rate or percent may be unreliable.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

ALL CAUSES	716.9
Heart Disease	187.7
Malignant Neoplasms (cancer)	148.2
Trachea, Bronchus, & Lung	40.4
Colon, Rectum, & Anus	11.0
Female Breast	16.7
Pancreas	6.8
Cerebrovascular Disease	43.4
Chronic Lower Respiratory Diseases	46.1
Accidents	46.8
Motor Vehicle Accidents	LNE
Alzheimer's Disease	27.8
Diabetes Mellitus	22.2
Influenza & Pneumonia	21.4
Intentional Self-Harm (suicide)	LNE
Chronic Liver Disease and Cirrhosis	17.0
Infant Mortality	LNE

Leading Causes of Death

Total Deaths

1. Heart Disease	72
2. Cancer	50
3. Cerebrovascular Disease	16
4. Chronic Lower Respiratory Disease	15
5. Alzheimer's Disease	12
6. Accidents	11
7. Influenza and Pneumonia	8
8. Diabetes	7
T9. Chronic Liver Disease and Cirrhosis	5
T9. Gastrointestinal hemorrhage, unspecified	5

Percent of deaths due to tobacco use

Median age at death

• Denotes a health status indicator which is significantly lower than the state average.

○ Denotes a health status indicator which is significantly higher than the state average.

³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.


See Technical Notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Clay County

Health Status Indicators 2007-2011

Demographic Information



Clay County is located in the southeastern part of the state and averages 33.6 persons per square mile.

2011 Population Information

Subject	Number	Percent
Total population	14,051	100.0
White	12,800	91.1
Black or African American	197	1.4
American Indian and Alaska Native	464	3.3
Asian	267	1.9
Native Hawaiian and Other Pacific Islander	0	0.0
Two or More Races	309	2.2
Under 5 years	759	5.4
Under 18 years	2,431	17.3
65 years and over	1,419	10.1

Source: United States Census Bureau, 2011 American Community Survey estimates

Nativity

Percent of Low Birth Weight Infants	6.3
○ Percent of Mothers Receiving Care in 1st Trimester	77.7
● Percent of Mothers Who Used Tobacco While Pregnant ¹	11.0
Percent of Births Less Than 37 Weeks of Gestation	7.8
○ Average Age of Mother	27.6
Teenage Birth Rate ²	4.9
Percent White Births	91.0
Percent American Indian Births	4.4
● Percent Unmarried	27.1
Percent WIC births	38.3
Percent Breastfeeding at discharge	80.0
Percent Payment-Private Insurance	58.5
Percent Payment-Medicaid	35.7
○ Percent C-Section	32.2

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

LNE (Low Number of Events): is used because the rate or percent may be unreliable.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

ALL CAUSES	701.6
Heart Disease	188.2
Malignant Neoplasms (cancer)	150.5
Trachea, Bronchus, & Lung	33.2
Colon, Rectum, & Anus	20.0
Female Breast	28.5
Pancreas	4.7
Cerebrovascular Disease	54.3
Chronic Lower Respiratory Diseases	53.0
Accidents	39.2
Motor Vehicle Accidents	12.7
Alzheimer's Disease	25.5
Diabetes Mellitus	30.1
Influenza & Pneumonia	15.2
Intentional Self-Harm (suicide)	8.9
Chronic Liver Disease and Cirrhosis	10.1
Infant Mortality	LNE

Leading Causes of Death

Total Deaths

1. Heart Disease	123
2. Cancer	88
3. Cerebrovascular Disease	36
4. Chronic Lower Respiratory Disease	33
5. Accidents	22
T6. Alzheimer's Disease	18
T6. Diabetes	18
8. Influenza and Pneumonia	10
9. Nephritis, nephrotic syndrome, and nephrosis	8
10. Septicemia	6

Percent of deaths due to tobacco use

Median age at death

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See Technical Notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Codington County

Demographic Information



Codington County is located in eastern South Dakota and averages 39.5 persons per square mile.

2011 Population Information

Subject	Number	Percent
Total population	27,442	100.0
White	26,235	95.6
Black or African American	137	0.5
American Indian and Alaska Native	576	2.1
Asian	137	0.5
Native Hawaiian and Other Pacific Islander	0	0.0
Two or More Races	357	1.3
Under 5 years	2,058	7.5
Under 18 years	6,751	24.6
65 years and over	4,061	14.8

Source: United States Census Bureau, 2011 American Community Survey estimates

Health Status Indicators 2007-2011

Natality

Percent of Low Birth Weight Infants	5.5
○ Percent of Mothers Receiving Care in 1st Trimester	77.1
○ Percent of Mothers Who Used Tobacco While Pregnant ¹	24.7
● Percent of Births Less Than 37 Wks. of Gestation	6.8
Average Age of Mother	26.7
Teenage Birth Rate ²	13.5
Percent White Births	93.4
Percent American Indian Births	3.9
Percent Unmarried	38.2
Percent WIC births	40.0
Percent Breastfeeding at discharge	73.8
○ Percent Payment-Private Insurance	64.6
● Percent Payment-Medicaid	30.8
● Percent C-Section	22.8

● Denotes a health status indicator which is significantly lower than the state average.

○ Denotes a health status indicator which is significantly higher than the state average.

LNE (Low Number of Events): is used because the rate or percent may be unreliable.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

ALL CAUSES	703.4
Heart Disease	166.9
Malignant Neoplasms (cancer)	175.7
Trachea, Bronchus, & Lung	51.9
Colon, Rectum, & Anus	14.5
Female Breast	18.6
Pancreas	16.1
Cerebrovascular Disease	38.6
Chronic Lower Respiratory Diseases	40.6
Accidents	39.8
Motor Vehicle Accidents	11.6
Alzheimer's Disease	34.8
Diabetes Mellitus	21.5
Influenza & Pneumonia	18.4
Intentional Self-Harm (suicide)	19.4
Chronic Liver Disease and Cirrhosis	6.6
Infant Mortality	5.10

Leading Causes of Death

Total Deaths

1. Heart Disease	301
2. Cancer	286
3. Cerebrovascular Disease	71
4. Chronic Lower Respiratory Disease	70
5. Alzheimer's Disease	67
6. Accidents	61
7. Diabetes	36
8. Influenza and Pneumonia	34
9. Suicide	24
10. Vascular Dementia	14

Percent of deaths due to tobacco use
 22.8 |

Median age at death
 81 |

● Denotes a health status indicator which is significantly lower than the state average.

○ Denotes a health status indicator which is significantly higher than the state average.


³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See Technical Notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Corson County

Demographic Information



Corson County is located in the north central portion of the state and averages 1.6 persons per square mile.

2011 Population Information

Subject	Number	Percent
Total population	4,022	100.0
White	1,239	30.8
Black or African American	8	0.2
American Indian and Alaska Native	2,655	66.0
Asian	16	0.4
Native Hawaiian and Other Pacific Islander	4	0.1
Two or More Races	101	2.5
Under 5 years	370	9.2
Under 18 years	1,392	34.6
65 years and over	418	10.4

Source: United States Census Bureau, 2011 American Community Survey estimates

Health Status Indicators 2007-2011

Natality

Percent of Low Birth Weight Infants	6.4
● Percent of Mothers Receiving Care in 1st Trimester	48.7
○ Percent of Mothers Who Used Tobacco While Pregnant ¹	24.8
Percent of Births Less Than 37 Wks. of Gestation	11.4
● Average Age of Mother	24.9
○ Teenage Birth Rate ²	42.4
Percent White Births	21.4
Percent American Indian Births	76.0
○ Percent Unmarried	70.9
○ Percent WIC births	75.7
● Percent Breastfeeding at discharge	49.2
● Percent Payment-Private Insurance	17.3
○ Percent Payment-Medicaid	69.7
Percent C-Section	23.2

● Denotes a health status indicator which is significantly lower than the state average.
 ○ Denotes a health status indicator which is significantly higher than the state average.
 LNE (Low Number of Events): is used because the rate or percent may be unreliable.
¹Data for mothers who used tobacco are self-reported.
²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

○ ALL CAUSES	1213.7
○ Heart Disease	279.6
Malignant Neoplasms (cancer)	215.7
Trachea, Bronchus, & Lung	55.3
Colon, Rectum, & Anus	LNE
Female Breast	68.1
Pancreas	LNE
Cerebrovascular Disease	32.8
Chronic Lower Respiratory Diseases	61.6
○ Accidents	138.6
○ Motor Vehicle Accidents	60.4
Alzheimer's Disease	35.6
○ Diabetes Mellitus	97.6
Influenza & Pneumonia	18.1
○ Intentional Self-Harm (suicide)	63.4
Chronic Liver Disease and Cirrhosis	19.1
Infant Mortality	11.82

Leading Causes of Death

Total Deaths

1. Heart Disease	46
2. Cancer	37
3. Accidents	26
4. Diabetes	17
5. Suicide	14
6. Chronic Lower Respiratory Disease	10
T7. Cerebrovascular Disease	5
T7. Alzheimer's Disease	5

Percent of deaths due to tobacco use 17.1
 Median age at death 66


● Denotes a health status indicator which is significantly lower than the state average.
 ○ Denotes a health status indicator which is significantly higher than the state average.
³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See Technical Notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Custer County

Demographic Information



Custer County is located in the southern Black Hills and averages 5.3 persons per square mile.

2011 Population Information

Subject	Number	Percent
Total population	8,338	100.0
White	7,846	94.1
Black or African American	42	0.5
American Indian and Alaska Native	258	3.1
Asian	33	0.4
Native Hawaiian and Other Pacific Islander	0	0.0
Two or More Races	158	1.9
Under 5 years	375	4.5
Under 18 years	1,601	19.2
65 years and over	1,843	22.1

Source: United States Census Bureau, 2011 American Community Survey estimates

Health Status Indicators 2007-2011

Natality

Percent of Low Birth Weight Infants	6.6
Percent of Mothers Receiving Care in 1st Trimester	70.6
Percent of Mothers Who Used Tobacco While Pregnant ¹	18.2
Percent of Births Less Than 37 Weeks of Gestation	9.7
Average Age of Mother	27.5
Teenage Birth Rate ²	11.8
Percent White Births	90.3
Percent American Indian Births	5.1
• Percent Unmarried	29.4
Percent WIC births	37.9
Percent Breastfeeding at discharge	82.6
Percent Payment-Private Insurance	52.7
Percent Payment-Medicaid	37.9
• Percent C-Section	19.2

- Denotes a health status indicator which is significantly lower than the state average.
- Denotes a health status indicator which is significantly higher than the state average.
- LNE (Low Number of Events): is used because the rate or percent may be unreliable.
- ¹Data for mothers who used tobacco are self-reported.
- ²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

ALL CAUSES	662.7
Heart Disease	134.7
Malignant Neoplasms (cancer)	184.0
Trachea, Bronchus, & Lung	49.1
Colon, Rectum, & Anus	14.2
Female Breast	23.1
Pancreas	LNE
Cerebrovascular Disease	45.8
• Chronic Lower Respiratory Diseases	23.4
Accidents	52.1
Motor Vehicle Accidents	9.9
Alzheimer's Disease	29.3
Diabetes Mellitus	13.4
Influenza & Pneumonia	15.3
Intentional Self-Harm (suicide)	9.6
Chronic Liver Disease and Cirrhosis	8.5
Infant Mortality	LNE

Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Cancer	113
2. Heart Disease	84
3. Cerebrovascular Disease	26
4. Accidents	23
5. Alzheimer's Disease	17
6. Chronic Lower Respiratory Disease	15
7. Influenza and Pneumonia	9
8. Diabetes	8
T9. Parkinson's Disease	6
T9. Disorders of lipoprotein metabolism and other lipidemias	6
T9. Vascular disorders of intestine	6

Percent of deaths due to tobacco use20.7
Median age at death 79


- Denotes a health status indicator which is significantly lower than the state average.
- Denotes a health status indicator which is significantly higher than the state average.
- ³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See Technical Notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Davison County

Demographic Information



Davison County is located in eastern South Dakota and averages 44.8 persons per square mile.

2011 Population Information

Subject	Number	Percent
Total population	19,651	100.0
White	18,590	94.6
Black or African American	138	0.7
American Indian and Alaska Native	511	2.6
Asian	98	0.5
Native Hawaiian and Other Pacific Islander	20	0.1
Two or More Races	295	1.5
Under 5 years	1,376	7.0
Under 18 years	4,539	23.1
65 years and over	3,341	17.0

Source: United States Census Bureau, 2011 American Community Survey estimates

Health Status Indicators 2007-2011

Nativity

Percent of Low Birth Weight Infants	6.4
○ Percent of Mothers Receiving Care in 1st Trimester	76.8
○ Percent of Mothers Who Used Tobacco While Pregnant ¹	22.3
Percent of Births Less Than 37 Weeks of Gestation	8.7
● Average Age of Mother	26.4
● Teenage Birth Rate ²	11.5
Percent White Births	90.5
Percent American Indian Births	7.1
Percent Unmarried	40.5
○ Percent WIC births	44.8
● Percent Breastfeeding at discharge	66.5
○ Percent Payment-Private Insurance	60.0
Percent Payment-Medicaid	37.0
○ Percent C-Section	31.2

● Denotes a health status indicator which is significantly lower than the state average.

○ Denotes a health status indicator which is significantly higher than the state average.

LNE (Low Number of Events): is used because the rate or percent may be unreliable.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

ALL CAUSES	687.1
Heart Disease	177.2
Malignant Neoplasms (cancer)	164.9
Trachea, Bronchus, & Lung	46.2
Colon, Rectum, & Anus	18.1
Female Breast	15.4
Pancreas	7.9
Cerebrovascular Disease	46.7
Chronic Lower Respiratory Diseases	45.3
Accidents	24.3
● Motor Vehicle Accidents	5.9
Alzheimer's Disease	29.6
● Diabetes Mellitus	12.2
Influenza & Pneumonia	19.9
Intentional Self-Harm (suicide)	16.2
Chronic Liver Disease and Cirrhosis	9.4
Infant Mortality	5.77

Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	256
2. Cancer	215
3. Cerebrovascular Disease	69
4. Chronic Lower Respiratory Disease	61
5. Alzheimer's Disease	50
6. Influenza and Pneumonia	37
7. Accidents	30
8. Pneumonitis due to solids and liquids	23
9. Parkinson's Disease	17
T10. Diabetes	16
T10. Unspecified Dementia	16

Percent of deaths due to tobacco use 19.6
Median age at death 82

● Denotes a health status indicator which is significantly lower than the state average.

○ Denotes a health status indicator which is significantly higher than the state average.


³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See Technical Notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Day County

Demographic Information



Day County is located in the northeastern part of the state and averages 5.6 persons per square mile.

2011 Population Information

Subject	Number	Percent
Total population	5,741	100.0
White	5,069	88.3
Black or African American	17	0.3
American Indian and Alaska Native	540	9.4
Asian	11	0.2
Native Hawaiian and Other Pacific Islander	0	0.0
Two or More Races	103	1.8
Under 5 years	344	6.0
Under 18 years	1,246	21.7
65 years and over	1,320	23.0

Source: United States Census Bureau, 2011 American Community Survey estimates

Health Status Indicators 2007-2011

Natality

Percent of Low Birth Weight Infants	4.9
Percent of Mothers Receiving Care in 1st Trimester	69.1
○ Percent of Mothers Who Used Tobacco While Pregnant ¹	28.5
● Percent of Births Less Than 37 Wks. of Gestation	5.5
Average Age of Mother	27.4
Teenage Birth Rate ²	9.3
Percent White Births	73.7
Percent American Indian Births	20.4
Percent Unmarried	40.6
Percent WIC births	41.2
Percent Breastfeeding at discharge	69.8
Percent Payment-Private Insurance	53.2
Percent Payment-Medicaid	38.5
Percent C-Section	21.2

● Denotes a health status indicator which is significantly lower than the state average.
 ○ Denotes a health status indicator which is significantly higher than the state average.
 LNE (Low Number of Events): is used because the rate or percent may be unreliable.
¹Data for mothers who used tobacco are self-reported.
²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

ALL CAUSES	729.1
Heart Disease	197.2
Malignant Neoplasms (cancer)	170.2
Trachea, Bronchus, & Lung	46.2
Colon, Rectum, & Anus	14.6
Female Breast	14.7
Pancreas	15.8
○ Cerebrovascular Disease	66.9
Chronic Lower Respiratory Diseases	44.1
Accidents	48.4
Motor Vehicle Accidents	16.8
Alzheimer's Disease	42.5
Diabetes Mellitus	14.6
Influenza & Pneumonia	15.0
Intentional Self-Harm (suicide)	LNE
Chronic Liver Disease and Cirrhosis	9.1
Infant Mortality	LNE

Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	110
2. Cancer	86
3. Cerebrovascular Disease	37
4. Alzheimer's Disease	28
5. Chronic Lower Respiratory Disease	24
6. Accidents	20
7. Influenza and Pneumonia	9
8. Diabetes	7
9. Nephritis, nephrotic syndrome, and nephrosis	5

Percent of deaths due to tobacco use 19.3
 Median age at death 83


● Denotes a health status indicator which is significantly lower than the state average.
 ○ Denotes a health status indicator which is significantly higher than the state average.
³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See Technical Notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Deuel County

Demographic Information



Deuel County borders Minnesota and averages 7.0 persons per square mile.

2011 Population Information

Subject	Number	Percent
Total population	4,359	100.0
White	4,285	98.3
Black or African American	17	0.4
American Indian and Alaska Native	22	0.5
Asian	4	0.1
Native Hawaiian and Other Pacific Islander	0	0.0
Two or More Races	31	0.7
Under 5 years	270	6.2
Under 18 years	1,011	23.2
65 years and over	850	19.5

Source: United States Census Bureau, 2011 American Community Survey estimates

Health Status Indicators 2007-2011

Natality

Percent of Low Birth Weight Infants	6.4
○ Percent of Mothers Receiving Care in 1st Trimester	84.4
Percent of Mothers Who Used Tobacco While Pregnant ¹	17.8
Percent of Births Less Than 37 Weeks of Gestation	8.0
Average Age of Mother	27.4
Teenage Birth Rate ²	10.2
Percent White Births	99.2
Percent American Indian Births	LNE
● Percent Unmarried	24.2
● Percent WIC births	27.4
Percent Breastfeeding at discharge	71.4
○ Percent Payment-Private Insurance	79.4
● Percent Payment-Medicaid	12.7
Percent C-Section	22.2

● Denotes a health status indicator which is significantly lower than the state average.
 ○ Denotes a health status indicator which is significantly higher than the state average.
 LNE (Low Number of Events): is used because the rate or percent may be unreliable.
¹Data for mothers who used tobacco are self-reported.
²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

ALL CAUSES	704.3
Heart Disease	141.4
Malignant Neoplasms (cancer)	129.6
Trachea, Bronchus, & Lung	64.3
Colon, Rectum, & Anus	13.5
Female Breast	16.1
Pancreas	LNE
Cerebrovascular Disease	35.1
Chronic Lower Respiratory Diseases	71.0
Accidents	54.6
Motor Vehicle Accidents	17.6
Alzheimer's Disease	42.7
Diabetes Mellitus	21.8
Influenza & Pneumonia	10.1
Intentional Self-Harm (suicide)	21.9
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	LNE

Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	51
2. Cancer	44
3. Chronic Lower Respiratory Disease	24
4. Alzheimer's Disease	18
T5. Cerebrovascular Disease	13
T5. Accidents	13
7. Septicemia	10
8. Diabetes	7
9. Nephritis, nephrotic syndrome, and nephrosis	5

Percent of deaths due to tobacco use 21.4
 Median age at death 81


● Denotes a health status indicator which is significantly lower than the state average.
 ○ Denotes a health status indicator which is significantly higher than the state average.
³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See Technical Notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Dewey County

Demographic Information



Dewey County is located in the north central region of the state and averages 2.3 persons per square mile.

2011 Population Information

Subject	Number	Percent
Total population	5,421	100.0
White	1,241	22.9
Black or African American	11	0.2
American Indian and Alaska Native	3,963	73.1
Asian	11	0.2
Native Hawaiian and Other Pacific Islander	0	0.0
Two or More Races	201	3.7
Under 5 years	526	9.7
Under 18 years	1,838	33.9
65 years and over	515	9.5

Source: United States Census Bureau, 2011 American Community Survey estimates

Health Status Indicators 2007-2011

Nativity

Percent of Low Birth Weight Infants	6.7
● Percent of Mothers Receiving Care in 1st Trimester	53.4
○ Percent of Mothers Who Used Tobacco While Pregnant ¹	28.9
Percent of Births Less Than 37 Weeks of Gestation	8.6
● Average Age of Mother	25.2
○ Teenage Birth Rate ²	58.1
Percent White Births	9.7
Percent American Indian Births	86.0
○ Percent Unmarried	67.3
○ Percent WIC births	82.1
● Percent Breastfeeding at discharge	49.3
● Percent Payment-Private Insurance	14.3
○ Percent Payment-Medicaid	77.2
○ Percent C-Section	34.5

● Denotes a health status indicator which is significantly lower than the state average.
○ Denotes a health status indicator which is significantly higher than the state average.

LNE (Low Number of Events): is used because the rate or percent may be unreliable.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

○ ALL CAUSES	1176.0
○ Heart Disease	261.3
Malignant Neoplasms (cancer)	207.1
Trachea, Bronchus, & Lung	79.4
Colon, Rectum, & Anus	19.4
Female Breast	LNE
Pancreas	LNE
Cerebrovascular Disease	26.3
Chronic Lower Respiratory Diseases	50.8
○ Accidents	115.2
○ Motor Vehicle Accidents	54.4
○ Alzheimer's Disease	42.4
Diabetes Mellitus	82.7
Influenza & Pneumonia	40.3
○ Intentional Self-Harm (suicide)	60.2
○ Chronic Liver Disease and Cirrhosis	60.4
Infant Mortality	LNE

Leading Causes of Death

Total Deaths

1. Heart Disease	56
2. Cancer	46
3. Accidents	27
4. Diabetes	19
5. Suicide	16
6. Chronic Liver Disease and Cirrhosis	15
7. Chronic Lower Respiratory Disease	10
T8. Alzheimer's Disease	8
T8. Influenza and Pneumonia	8
10. Mental and behavioral disorders due to use of alcohol	6

Percent of deaths due to tobacco use 17.2
Median age at death 65

● Denotes a health status indicator which is significantly lower than the state average.

○ Denotes a health status indicator which is significantly higher than the state average.


³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See Technical Notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Douglas County

Demographic Information



Douglas County is located in south central South Dakota and averages 7.0 persons per square mile.

2011 Population Information

Subject	Number	Percent
Total population	2,972	100.0
White	2,874	96.7
Black or African American	15	0.5
American Indian and Alaska Native	59	2.0
Asian	3	0.1
Native Hawaiian and Other Pacific Islander	0	0.0
Two or More Races	21	0.7
Under 5 years	158	5.3
Under 18 years	669	22.5
65 years and over	713	24.0

Source: United States Census Bureau, 2011 American Community Survey estimates

Health Status Indicators 2007-2011

Nativity

Percent of Low Birth Weight Infants	3.8
Percent of Mothers Receiving Care in 1st Trimester	70.1
Percent of Mothers Who Used Tobacco While Pregnant ¹	15.0
Percent of Births Less Than 37 Weeks of Gestation	5.0
○ Average Age of Mother	28.3
Teenage Birth Rate ²	LNE
Percent White Births	95.0
Percent American Indian Births	2.5
● Percent Unmarried	15.6
● Percent WIC births	26.9
Percent Breastfeeding at discharge	83.1
Percent Payment-Private Insurance	56.3
● Percent Payment-Medicaid	25.0
Percent C-Section	26.3

● Denotes a health status indicator which is significantly lower than the state average.

○ Denotes a health status indicator which is significantly higher than the state average.

LNE (Low Number of Events): is used because the rate or percent may be unreliable.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

ALL CAUSES	691.9
Heart Disease	144.0
Malignant Neoplasms (cancer)	183.7
Trachea, Bronchus, & Lung	25.3
Colon, Rectum, & Anus	10.7
Female Breast	30.6
Pancreas	23.8
Cerebrovascular Disease	43.6
Chronic Lower Respiratory Diseases	42.3
Accidents	37.9
Motor Vehicle Accidents	21.1
Alzheimer's Disease	38.1
Diabetes Mellitus	26.7
Influenza & Pneumonia	27.8
Intentional Self-Harm (suicide)	18.5
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	LNE

Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	57
2. Cancer	45
3. Alzheimer's Disease	20
4. Cerebrovascular Disease	17
5. Chronic Lower Respiratory Disease	15
6. Accidents	11
T7. Diabetes	10
T7. Influenza and Pneumonia	10
9. Pneumonitis due to solids and liquids	5

Percent of deaths due to tobacco use

Median age at death

● Denotes a health status indicator which is significantly lower than the state average.

○ Denotes a health status indicator which is significantly higher than the state average.


³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See Technical Notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Edmunds County

Demographic Information



Edmunds County is located in the north central region of the state and averages 3.6 persons per square mile.

2011 Population Information

Subject	Number	Percent
Total population	4,056	100.0
White	3,971	97.9
Black or African American	4	0.1
American Indian and Alaska Native	24	0.6
Asian	4	0.1
Native Hawaiian and Other Pacific Islander	0	0.0
Two or More Races	49	1.2
Under 5 years	227	5.6
Under 18 years	965	23.8
65 years and over	868	21.4

Source: United States Census Bureau, 2011 American Community Survey estimates

Health Status Indicators 2007-2011

Natality

Percent of Low Birth Weight Infants	4.6
Percent of Mothers Receiving Care in 1st Trimester	71.2
• Percent of Mothers Who Used Tobacco While Pregnant ¹	10.8
Percent of Births Less Than 37 Weeks of Gestation	6.0
Average Age of Mother	27.7
Teenage Birth Rate ²	19.0
Percent White Births	97.7
Percent American Indian Births	LNE
• Percent Unmarried	18.8
• Percent WIC births	22.9
Percent Breastfeeding at discharge	78.0
○ Percent Payment-Private Insurance	71.6
• Percent Payment-Medicaid	15.1
Percent C-Section	29.8

•Denotes a health status indicator which is significantly lower than the state average.
 ○Denotes a health status indicator which is significantly higher than the state average.
 LNE (Low Number of Events): is used because the rate or percent may be unreliable.
¹Data for mothers who used tobacco are self-reported.
²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

ALL CAUSES	661.7
Heart Disease	166.0
Malignant Neoplasms (cancer)	143.5
• Trachea, Bronchus, & Lung	24.6
Colon, Rectum, & Anus	14.9
Female Breast	32.1
Pancreas	12.4
Cerebrovascular Disease	29.0
Chronic Lower Respiratory Diseases	62.4
Accidents	29.2
Motor Vehicle Accidents	16.8
○ Alzheimer's Disease	33.9
Diabetes Mellitus	50.5
Influenza & Pneumonia	8.2
Intentional Self-Harm (suicide)	16.0
Chronic Liver Disease and Cirrhosis	11.9
Infant Mortality	LNE

Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	65
2. Cancer	47
3. Chronic Lower Respiratory Disease	23
4. Diabetes	19
5. Alzheimer's Disease	16
6. Cerebrovascular Disease	11
7. Accidents	8

Percent of deaths due to tobacco use 16.3
 Median age at death..... 82


•Denotes a health status indicator which is significantly lower than the state average.
 ○Denotes a health status indicator which is significantly higher than the state average.
³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See Technical Notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Fall River County

Demographic Information



Fall River County is located in the southwestern corner of the state and averages 4.1 persons per square mile.

2011 Population Information

Subject	Number	Percent
Total population	6,981	100.0
White	6,199	88.8
Black or African American	63	0.9
American Indian and Alaska Native	489	7.0
Asian	28	0.4
Native Hawaiian and Other Pacific Islander	7	0.1
Two or More Races	195	2.8
Under 5 years	265	3.8
Under 18 years	1,201	17.2
65 years and over	1,668	23.9

Source: United States Census Bureau, 2011 American Community Survey estimates

Health Status Indicators 2007-2011

Natality

Percent of Low Birth Weight Infants	7.7
● Percent of Mothers Receiving Care in 1st Trimester	55.7
○ Percent of Mothers Who Used Tobacco While Pregnant ¹	28.7
Percent of Births Less Than 37 Weeks of Gestation	8.5
Average Age of Mother	26.4
● Teenage Birth Rate ²	8.2
Percent White Births	81.9
Percent American Indian Births	11.2
Percent Unmarried	42.5
○ Percent WIC births	51.7
Percent Breastfeeding at discharge	78.8
● Percent Payment-Private Insurance	43.2
○ Percent Payment-Medicaid	51.4
● Percent C-Section	17.8

● Denotes a health status indicator which is significantly lower than the state average.
 ○ Denotes a health status indicator which is significantly higher than the state average.
 LNE (Low Number of Events): is used because the rate or percent may be unreliable.
¹Data for mothers who used tobacco are self-reported.
²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

○ ALL CAUSES	828.2
○ Heart Disease	213.6
Malignant Neoplasms (cancer)	200.2
○ Trachea, Bronchus, & Lung	69.2
● Colon, Rectum, & Anus	3.8
Female Breast	25.5
Pancreas	15.5
Cerebrovascular Disease	29.9
Chronic Lower Respiratory Diseases	62.8
Accidents	60.0
Motor Vehicle Accidents	20.8
● Alzheimer's Disease	21.8
○ Diabetes Mellitus	41.0
Influenza & Pneumonia	13.1
Intentional Self-Harm (suicide)	12.9
Chronic Liver Disease and Cirrhosis	20.1
Infant Mortality	11.58

Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	142
2. Cancer	136
3. Chronic Lower Respiratory Disease	45
4. Diabetes	29
5. Accidents	25
6. Cerebrovascular Disease	21
7. Alzheimer's Disease	17
8. Influenza and Pneumonia	10
T9. Chronic Liver Disease and Cirrhosis	9
T9. Unspecified Dementia	9
T9. Pneumonitis due to solids and liquids	9

Percent of deaths due to tobacco use 19.6
 Median age at death 78


● Denotes a health status indicator which is significantly lower than the state average.
 ○ Denotes a health status indicator which is significantly higher than the state average.
³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See Technical Notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Faulk County

Demographic Information



Faulk County is located in north central South Dakota and averages 2.4 persons per square mile.

2011 Population Information

Subject	Number	Percent
Total population	2,367	100.0
White	2,336	98.7
Black or African American	2	0.1
American Indian and Alaska Native	5	0.2
Asian	2	0.1
Native Hawaiian and Other Pacific Islander	0	0.0
Two or More Races	21	0.9
Under 5 years	142	6.0
Under 18 years	552	23.3
65 years and over	554	23.4

Source: United States Census Bureau, 2011 American Community Survey estimates

Health Status Indicators 2007-2011

Natality

● Percent of Low Birth Weight Infants	2.5
Percent of Mothers Receiving Care in 1st Trimester	56.3
● Percent of Mothers Who Used Tobacco While Pregnant ¹	7.4
● Percent of Births Less Than 37 Wks. of Gestation	4.1
○ Average Age of Mother	29.8
Teenage Birth Rate ²	LNE
Percent White Births	100.0
Percent American Indian Births	LNE
● Percent Unmarried	13.9
● Percent WIC births	21.3
Percent Breastfeeding at discharge	82.0
○ Percent Payment-Private Insurance	75.4
● Percent Payment-Medicaid	12.3
Percent C-Section	23.0

● Denotes a health status indicator which is significantly lower than the state average.
 ○ Denotes a health status indicator which is significantly higher than the state average.
 LNE (Low Number of Events): is used because the rate or percent may be unreliable.
¹Data for mothers who used tobacco are self-reported.
²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

● ALL CAUSES	574.2
● Heart Disease	84.3
Malignant Neoplasms (cancer)	122.0
Trachea, Bronchus, & Lung	23.1
Colon, Rectum, & Anus	23.8
Female Breast	LNE
Pancreas	LNE
Cerebrovascular Disease	35.3
Chronic Lower Respiratory Diseases	41.8
Accidents	28.3
Motor Vehicle Accidents	LNE
Alzheimer's Disease	62.6
Diabetes Mellitus	15.7
Influenza & Pneumonia	41.1
Intentional Self-Harm (suicide)	LNE
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	LNE

Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Cancer	28
2. Heart Disease	20
3. Alzheimer's Disease	19
4. Chronic Lower Respiratory Disease	10
T5. Cerebrovascular Disease	9
T5. Influenza and Pneumonia	9
7. Accidents	5

Percent of deaths due to tobacco use 8.1
 Median age at death 85


● Denotes a health status indicator which is significantly lower than the state average.
 ○ Denotes a health status indicator which is significantly higher than the state average.
³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See Technical Notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Grant County

Demographic Information



Grant County borders Minnesota in northeastern South Dakota and averages 10.8 persons per square mile.

2011 Population Information

Subject	Number	Percent
Total population	7,250	100.0
White	7,112	98.1
Black or African American	15	0.2
American Indian and Alaska Native	44	0.6
Asian	22	0.3
Native Hawaiian and Other Pacific Islander	0	0.0
Two or More Races	51	0.7
Under 5 years	428	5.9
Under 18 years	1,639	22.6
65 years and over	1,370	18.9

Source: United States Census Bureau, 2011 American Community Survey estimates

Health Status Indicators 2007-2011

Natality

● Percent of Low Birth Weight Infants	4.1
Percent of Mothers Receiving Care in 1st Trimester	74.0
Percent of Mothers Who Used Tobacco While Pregnant ¹	16.4
● Percent of Births Less Than 37 Wks. of Gestation	5.0
Average Age of Mother	27.2
Teenage Birth Rate ²	18.7
Percent White Births	96.9
Percent American Indian Births	0.8
● Percent Unmarried	31.4
● Percent WIC births	30.6
● Percent Breastfeeding at discharge	62.2
Percent Payment-Private Insurance	50.9
● Percent Payment-Medicaid	20.6
Percent C-Section	25.4

● Denotes a health status indicator which is significantly lower than the state average.

○ Denotes a health status indicator which is significantly higher than the state average.

LNE (Low Number of Events): is used because the rate or percent may be unreliable.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

ALL CAUSES	663.7
Heart Disease	151.1
Malignant Neoplasms (cancer)	145.8
● Trachea, Bronchus, & Lung	26.3
Colon, Rectum, & Anus	19.9
Female Breast	23.8
Pancreas	8.2
Cerebrovascular Disease	48.2
Chronic Lower Respiratory Diseases	43.2
Accidents	47.1
Motor Vehicle Accidents	27.0
Alzheimer's Disease	41.3
Diabetes Mellitus	27.6
● Influenza & Pneumonia	8.2
Intentional Self-Harm (suicide)	14.0
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	LNE

Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	92
2. Cancer	83
3. Cerebrovascular Disease	32
4. Alzheimer's Disease	31
5. Chronic Lower Respiratory Disease	27
6. Diabetes	16
7. Accidents	15
8. Aortic aneurysm and dissection	9
9. Influenza and Pneumonia	6

Percent of deaths due to tobacco use	15.4
Median age at death	82

● Denotes a health status indicator which is significantly lower than the state average.

○ Denotes a health status indicator which is significantly higher than the state average.


³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See Technical Notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Gregory County

Demographic Information



Gregory County borders the west bank of the Missouri River and the state of Nebraska and averages 4.2 persons per square mile.

2011 Population Information

Subject	Number	Percent
Total population	4,216	100.0
White	3,794	90.0
Black or African American	8	0.2
American Indian and Alaska Native	312	7.4
Asian	13	0.3
Native Hawaiian and Other Pacific Islander	0	0.0
Two or More Races	89	2.1
Under 5 years	249	5.9
Under 18 years	944	22.4
65 years and over	1,003	23.8

Source: United States Census Bureau, 2011 American Community Survey estimates

Health Status Indicators 2007-2011

Nativity

Percent of Low Birth Weight Infants	5.4
Percent of Mothers Receiving Care in 1st Trimester	63.8
Percent of Mothers Who Used Tobacco While Pregnant ¹	19.7
Percent of Births Less Than 37 Weeks of Gestation	9.9
Average Age of Mother	26.9
Teenage Birth Rate ²	12.8
Percent White Births	81.1
Percent American Indian Births	14.4
Percent Unmarried	35.6
o Percent WIC births	50.5
Percent Breastfeeding at discharge	68.9
Percent Payment-Private Insurance	47.3
o Percent Payment-Medicaid	46.4
Percent C-Section	32.4

●Denotes a health status indicator which is significantly lower than the state average.
 ○Denotes a health status indicator which is significantly higher than the state average.
 LNE (Low Number of Events): is used because the rate or percent may be unreliable.
¹Data for mothers who used tobacco are self-reported.
²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

ALL CAUSES	773.6
o Heart Disease	207.2
Malignant Neoplasms (cancer)	182.1
Trachea, Bronchus, & Lung	48.6
Colon, Rectum, & Anus	19.1
Female Breast	12.1
Pancreas	LNE
Cerebrovascular Disease	28.1
Chronic Lower Respiratory Diseases	58.6
Accidents	64.9
Motor Vehicle Accidents	32.0
Alzheimer's Disease	22.0
Diabetes Mellitus	44.9
● Influenza & Pneumonia	4.4
Intentional Self-Harm (suicide)	21.5
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	LNE

Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	99
2. Cancer	74
3. Chronic Lower Respiratory Disease	26
T4. Accidents	19
T4. Diabetes	19
6. Disorders of lipoprotein metabolism and other lipedemias	16
7. Alzheimer's Disease	13
8. Cerebrovascular Disease	11

Percent of deaths due to tobacco use 21.5
 Median age at death 84


●Denotes a health status indicator which is significantly lower than the state average.
 ○Denotes a health status indicator which is significantly higher than the state average.
³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See Technical Notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Haakon County

Demographic Information



Haakon County is located in the west central region of the state and averages 1.1 persons per square mile.

2011 Population Information

Subject	Number	Percent
Total population	1,907	100.0
White	1,800	94.4
Black or African American	4	0.2
American Indian and Alaska Native	38	2.0
Asian	11	0.6
Native Hawaiian and Other Pacific Islander	2	0.1
Two or More Races	53	2.8
Under 5 years	128	6.7
Under 18 years	441	23.1
65 years and over	406	21.3

Source: United States Census Bureau, 2011 American Community Survey estimates

Health Status Indicators 2007-2011

Nativity

Percent of Low Birth Weight Infants	LNE
Percent of Mothers Receiving Care in 1st Trimester	68.7
• Percent of Mothers Who Used Tobacco While Pregnant ¹	10.5
Percent of Births Less Than 37 Weeks of Gestation	7.0
• Average Age of Mother	27.7
Teenage Birth Rate ²	16.7
Percent White Births	93.0
Percent American Indian Births	2.6
• Percent Unmarried	21.7
• Percent WIC births	21.7
Percent Breastfeeding at discharge	90.4
○ Percent Payment-Private Insurance	77.4
• Percent Payment-Medicaid	16.5
Percent C-Section	30.4

• Denotes a health status indicator which is significantly lower than the state average.
 ○ Denotes a health status indicator which is significantly higher than the state average.
 LNE (Low Number of Events): is used because the rate or percent may be unreliable.
¹Data for mothers who used tobacco are self-reported.
²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

ALL CAUSES	648.4
Heart Disease	117.4
Malignant Neoplasms (cancer)	160.4
Trachea, Bronchus, & Lung	41.9
Colon, Rectum, & Anus	24.1
Female Breast	LNE
Pancreas	LNE
Cerebrovascular Disease	36.8
Chronic Lower Respiratory Diseases	26.5
Accidents	66.3
Motor Vehicle Accidents	52.3
Alzheimer's Disease	LNE
Diabetes Mellitus	17.3
○ Influenza & Pneumonia	49.7
Intentional Self-Harm (suicide)	LNE
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	LNE

Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Cancer	29
2. Heart Disease	23
3. Influenza and Pneumonia	11
4. Accidents	8
5. Nephritis, nephrotic syndrome, nephritis	6
T6. Chronic Lower Respiratory Disease	5
T6. Cerebrovascular Disease	5

Percent of deaths due to tobacco use 12.4
 Median age at death 85


• Denotes a health status indicator which is significantly lower than the state average.
 ○ Denotes a health status indicator which is significantly higher than the state average.
³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See Technical Notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Hamlin County

Demographic Information



Hamlin County is located in the northeastern region of the state and averages 11.6 persons per square mile.

2011 Population Information

Subject	Number	Percent
Total population	5,978	100.0
White	5,888	98.5
Black or African American	12	0.2
American Indian and Alaska Native	24	0.4
Asian	18	0.3
Native Hawaiian and Other Pacific Islander	0	0.0
Two or More Races	36	0.6
Under 5 years	538	9.0
Under 18 years	1,793	30.0
65 years and over	1,046	17.5

Source: United States Census Bureau, 2011 American Community Survey estimates

Health Status Indicators 2007-2011

Natality

● Percent of Low Birth Weight Infants	4.2
Percent of Mothers Receiving Care in 1st Trimester	68.5
Percent of Mothers Who Used Tobacco While Pregnant ¹	15.6
● Percent of Births Less Than 37 Wks. of Gestation	5.2
Average Age of Mother	27.2
Teenage Birth Rate ²	18.9
Percent White Births	99.2
Percent American Indian Births	LNE
● Percent Unmarried	11.4
Percent WIC births	40.6
○ Percent Breastfeeding at discharge	86.3
○ Percent Payment-Private Insurance	70.5
● Percent Payment-Medicaid	21.5
● Percent C-Section	15.8

● Denotes a health status indicator which is significantly lower than the state average.
 ○ Denotes a health status indicator which is significantly higher than the state average.
 LNE (Low Number of Events): is used because the rate or percent may be unreliable.
¹Data for mothers who used tobacco are self-reported.
²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

○ ALL CAUSES	814.6
Heart Disease	170.9
Malignant Neoplasms (cancer)	158.5
Trachea, Bronchus, & Lung	37.3
Colon, Rectum, & Anus	22.9
Female Breast	LNE
Pancreas	15.7
○ Cerebrovascular Disease	67.0
Chronic Lower Respiratory Diseases	54.4
Accidents	42.9
Motor Vehicle Accidents	15.2
○ Alzheimer's Disease	101.6
Diabetes Mellitus	30.0
Influenza & Pneumonia	24.2
Intentional Self-Harm (suicide)	LNE
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	LNE

Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	78
2. Cancer	60
3. Alzheimer's Disease	54
4. Cerebrovascular Disease	32
5. Chronic Lower Respiratory Disease	22
6. Accidents	17
7. Diabetes	13
8. Influenza and Pneumonia	11
T9. Parkinson's Disease	7
T9. Vascular Dementia	7

Percent of deaths due to tobacco use 12.7
 Median age at death 84


● Denotes a health status indicator which is significantly lower than the state average.
 ○ Denotes a health status indicator which is significantly higher than the state average.
³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See Technical Notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Hand County

Demographic Information



Hand County is located in central South Dakota and averages 2.4 persons per square mile.

2011 Population Information

Subject	Number	Percent
Total population	3,423	100.0
White	3,372	98.5
Black or African American	3	0.1
American Indian and Alaska Native	10	0.3
Asian	10	0.3
Native Hawaiian and Other Pacific Islander	0	0.0
Two or More Races	27	0.8
Under 5 years	181	5.3
Under 18 years	705	20.6
65 years and over	849	24.8

Source: United States Census Bureau, 2011 American Community Survey estimates

Health Status Indicators 2007-2011

Nativity

Percent of Low Birth Weight Infants	8.6
Percent of Mothers Receiving Care in 1st Trimester	71.9
• Percent of Mothers Who Used Tobacco While Pregnant ¹	8.2
Percent of Births Less Than 37 Weeks of Gestation	8.6
○ Average Age of Mother	28.2
Teenage Birth Rate ²	LNE
Percent White Births	97.8
Percent American Indian Births	1.6
• Percent Unmarried	18.4
• Percent WIC births	24.3
Percent Breastfeeding at discharge	77.3
○ Percent Payment-Private Insurance	76.2
• Percent Payment-Medicaid	15.7
○ Percent C-Section	41.6

• Denotes a health status indicator which is significantly lower than the state average.
 ○ Denotes a health status indicator which is significantly higher than the state average.
 LNE (Low Number of Events): is used because the rate or percent may be unreliable.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

• ALL CAUSES	555.0
• Heart Disease	115.8
Malignant Neoplasms (cancer)	125.9
Trachea, Bronchus, & Lung	38.9
Colon, Rectum, & Anus	8.3
Female Breast	LNE
Pancreas	LNE
Cerebrovascular Disease	31.4
Chronic Lower Respiratory Diseases	34.4
Accidents	45.2
Motor Vehicle Accidents	19.8
• Alzheimer's Disease	10.3
Diabetes Mellitus	17.9
Influenza & Pneumonia	LNE
Intentional Self-Harm (suicide)	LNE
Chronic Liver Disease and Cirrhosis	16.6
Infant Mortality	LNE

Leading Causes of Death

Total Deaths

1. Heart Disease	48
2. Cancer	41
3. Atherosclerosis	14
T4. Chronic Lower Respiratory Disease	12
T4. Cerebrovascular Disease	12
6. Accidents	7
T7. Diabetes	6
T7. Essential (primary) hypertension and hypertensive renal disease	6

Percent of deaths due to tobacco use 19.5
 Median age at death 83

• Denotes a health status indicator which is significantly lower than the state average.
 ○ Denotes a health status indicator which is significantly higher than the state average.


³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See Technical Notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Hanson County

Demographic Information



Hanson County is located in southeastern South Dakota and averages 7.7 people per square mile.

2011 Population Information

Subject	Number	Percent
Total population	3,376	100.0
White	3,322	98.4
Black or African American	3	0.1
American Indian and Alaska Native	20	0.6
Asian	14	0.4
Native Hawaiian and Other Pacific Islander	3	0.1
Two or More Races	17	0.5
Under 5 years	297	8.8
Under 18 years	1,080	32.0
65 years and over	469	13.9

Source: United States Census Bureau, 2011 American Community Survey estimates

Health Status Indicators 2007-2011

Natality

Percent of Low Birth Weight Infants	9.0
Percent of Mothers Receiving Care in 1st Trimester	66.5
• Percent of Mothers Who Used Tobacco While Pregnant ¹	7.8
Percent of Births Less Than 37 Wks. of Gestation	10.0
○ Average Age of Mother	29.5
Teenage Birth Rate ²	LNE
Percent White Births	99.6
Percent American Indian Births	LNE
• Percent Unmarried	10.7
• Percent WIC births	18.9
Percent Breastfeeding at discharge	77.7
○ Percent Payment-Private Insurance	79.8
• Percent Payment-Medicaid	11.2
Percent C-Section	33.5

•Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

LNE (Low Number of Events): is used because the rate or percent may be unreliable.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

ALL CAUSES	672.9
Heart Disease	178.9
Malignant Neoplasms (cancer)	172.0
Trachea, Bronchus, & Lung	60.4
Colon, Rectum, & Anus	LNE
Female Breast	LNE
Pancreas	LNE
Cerebrovascular Disease	62.6
• Chronic Lower Respiratory Diseases	17.7
Accidents	46.5
Motor Vehicle Accidents	LNE
Alzheimer's Disease	LNE
Diabetes Mellitus	LNE
Influenza & Pneumonia	LNE
Intentional Self-Harm (suicide)	LNE
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	LNE

Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Cancer	30
2. Heart Disease	27
3. Cerebrovascular Disease	9
4. Accidents	7

Percent of deaths due to tobacco use

Median age at death

•Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See Technical Notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Harding County

Demographic Information



Harding County is located in the northwestern corner of the state and averages 0.5 persons per square mile.

2011 Population Information

Subject	Number	Percent
Total population	1,269	100.0
White	1,227	96.7
Black or African American	1	0.1
American Indian and Alaska Native	20	1.6
Asian	1	0.1
Native Hawaiian and Other Pacific Islander	0	0.0
Two or More Races	20	1.6
Under 5 years	71	5.6
Under 18 years	291	22.9
65 years and over	183	14.4

Source: United States Census Bureau, 2011 American Community Survey estimates

Health Status Indicators 2007-2011

Natality

Percent of Low Birth Weight Infants	LNE
Percent of Mothers Receiving Care in 1st Trimester	77.4
• Percent of Mothers Who Used Tobacco While Pregnant ¹	8.2
Percent of Births Less Than 37 Wks. of Gestation	LNE
Average Age of Mother	28.2
Teenage Birth Rate ²	LNE
Percent White Births	98.4
Percent American Indian Births	LNE
• Percent Unmarried	15.9
• Percent WIC births	20.6
Percent Breastfeeding at discharge	87.3
Percent Payment-Private Insurance	63.5
• Percent Payment-Medicaid	17.5
Percent C-Section	34.9

- Denotes a health status indicator which is significantly lower than the state average.
- Denotes a health status indicator which is significantly higher than the state average.
- LNE (Low Number of Events): is used because the rate or percent may be unreliable.
- ¹Data for mothers who used tobacco are self-reported.
- ²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

ALL CAUSES	611.0
Heart Disease	192.4
Malignant Neoplasms (cancer)	94.7
Trachea, Bronchus, & Lung	41.5
Colon, Rectum, & Anus	LNE
Female Breast	LNE
Pancreas	LNE
Cerebrovascular Disease	57.3
Chronic Lower Respiratory Diseases	53.6
Accidents	86.4
Motor Vehicle Accidents	LNE
Alzheimer's Disease	LNE
Diabetes Mellitus	LNE
Influenza & Pneumonia	LNE
Intentional Self-Harm (suicide)	LNE
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	LNE

Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	15
2. Cancer	7
3. Accidents	5

Percent of deaths due to tobacco use	17.8
Median age at death	80


- Denotes a health status indicator which is significantly lower than the state average.
- Denotes a health status indicator which is significantly higher than the state average.
- ³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See Technical Notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Hughes County

Demographic Information



Hughes County is located in the center of the state and averages 23.0 persons per square mile.

2011 Population Information

Subject	Number	Percent
Total population	17,292	100.0
White	14,785	85.5
Black or African American	156	0.9
American Indian and Alaska Native	1,850	10.7
Asian	104	0.6
Native Hawaiian and Other Pacific Islander	0	0.0
Two or More Races	398	2.3
Under 5 years	1,193	6.9
Under 18 years	4,115	23.8
65 years and over	2,317	13.4

Source: United States Census Bureau, 2011 American Community Survey estimates

Health Status Indicators 2007-2011

Natality

● Percent of Low Birth Weight Infants	4.5
Percent of Mothers Receiving Care in 1st Trimester	70.7
Percent of Mothers Who Used Tobacco While Pregnant ¹	20.7
Percent of Births Less Than 37 Weeks of Gestation	7.7
Average Age of Mother	26.9
● Teenage Birth Rate ²	10.9
Percent White Births	73.0
Percent American Indian Births	21.9
Percent Unmarried	40.1
Percent WIC births	36.5
Percent Breastfeeding at discharge	71.7
Percent Payment-Private Insurance	57.2
Percent Payment-Medicaid	37.3
○ Percent C-Section	37.4

● Denotes a health status indicator which is significantly lower than the state average.
 ○ Denotes a health status indicator which is significantly higher than the state average.
 LNE (Low Number of Events): is used because the rate or percent may be unreliable.
¹Data for mothers who used tobacco are self-reported.
²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

ALL CAUSES	685.3
Heart Disease	158.6
Malignant Neoplasms (cancer)	160.3
Trachea, Bronchus, & Lung	40.2
Colon, Rectum, & Anus	21.7
Female Breast	10.9
Pancreas	14.3
Cerebrovascular Disease	48.0
Chronic Lower Respiratory Diseases	54.2
Accidents	25.4
Motor Vehicle Accidents	13.5
Alzheimer's Disease	37.9
Diabetes Mellitus	25.5
● Influenza & Pneumonia	8.4
Intentional Self-Harm (suicide)	18.6
Chronic Liver Disease and Cirrhosis	8.9
Infant Mortality	6.81

Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	174
2. Cancer	158
3. Chronic Lower Respiratory Disease	56
4. Cerebrovascular Disease	54
5. Alzheimer's Disease	42
6. Diabetes	27
7. Accidents	26
8. Suicide	15
9. Unspecified Dementia	13
10. Influenza and Pneumonia	11

Percent of deaths due to tobacco use 23.1
 Median age at death 80


● Denotes a health status indicator which is significantly lower than the state average.
 ○ Denotes a health status indicator which is significantly higher than the state average.
³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See Technical Notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Hutchinson County

Demographic Information



Hutchinson County is located in the southeastern region of the state and averages 9.0 persons per square mile.

2011 Population Information

Subject	Number	Percent
Total population	7,257	100.0
White	7,083	97.6
Black or African American	36	0.5
American Indian and Alaska Native	73	1.0
Asian	15	0.2
Native Hawaiian and Other Pacific Islander	0	0.0
Two or More Races	58	0.8
Under 5 years	435	6.0
Under 18 years	1,720	23.7
65 years and over	1,800	24.8

Source: United States Census Bureau, 2011 American Community Survey estimates

Health Status Indicators 2007-2011

Natality

- Percent of Low Birth Weight Infants3.4
- Percent of Mothers Receiving Care in 1st Trimester 62.4
- Percent of Mothers Who Used Tobacco While Pregnant¹8.0
- Percent of Births Less Than 37 Weeks of Gestation 6.7
- Average Age of Mother28.3
- Teenage Birth Rate²4.3
- Percent White Births98.9
- Percent American Indian BirthsLNE
- Percent Unmarried 18.0
- Percent WIC births21.4
- Percent Breastfeeding at discharge79.1
- Percent Payment-Private Insurance78.8
- Percent Payment-Medicaid17.5
- Percent C-Section23.8

●Denotes a health status indicator which is significantly lower than the state average.
 ○Denotes a health status indicator which is significantly higher than the state average.
 LNE (Low Number of Events): is used because the rate or percent may be unreliable.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

ALL CAUSES	652.9
Heart Disease	150.6
Malignant Neoplasms (cancer)	164.1
Trachea, Bronchus, & Lung	43.0
Colon, Rectum, & Anus	12.4
Female Breast	26.2
Pancreas	5.3
Cerebrovascular Disease	32.6
Chronic Lower Respiratory Diseases	38.6
Accidents	46.1
Motor Vehicle Accidents	19.6
Alzheimer's Disease	27.4
Diabetes Mellitus	19.6
Influenza & Pneumonia	16.5
Intentional Self-Harm (suicide)	LNE
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	LNE

Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	146
2. Cancer	112
3. Cerebrovascular Disease	36
4. Alzheimer's Disease	34
5. Chronic Lower Respiratory Disease	28
6. Accidents	24
T7. Influenza and Pneumonia	16
T7. Nephritis, nephrotic syndrome, and nephrosis	16
9. Diabetes	14
10. Pneumonitis due to solids and liquids	11

Percent of deaths due to tobacco use 10.1
 Median age at death 85

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See Technical Notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Hyde County

Demographic Information



Hyde County is located in the central region of the state and averages 1.7 people per square mile.

2011 Population Information

Subject	Number	Percent
Total population	1,394	100.0
White	1,252	89.1
Black or African American	1	0.1
American Indian and Alaska Native	107	8.5
Asian	3	0.2
Native Hawaiian and Other Pacific Islander	1	0.1
Two or More Races	28	2.0
Under 5 years	82	5.9
Under 18 years	309	22.2
65 years and over	308	22.1

Source: United States Census Bureau, 2011 American Community Survey estimates

Health Status Indicators 2007-2011

Natality

Percent of Low Birth Weight Infants	LNE
Percent of Mothers Receiving Care in 1st Trimester	75.9
Percent of Mothers Who Used Tobacco While Pregnant ¹	16.4
Percent of Births Less Than 37 Weeks of Gestation	7.3
Average Age of Mother	26.9
Teenage Birth Rate ²	LNE
Percent White Births	81.8
Percent American Indian Births	14.5
Percent Unmarried	32.7
Percent WIC births	45.5
Percent Breastfeeding at discharge	72.7
Percent Payment-Private Insurance	60.0
Percent Payment-Medicaid	38.2
Percent C-Section	29.1

●Denotes a health status indicator which is significantly lower than the state average.
 ○Denotes a health status indicator which is significantly higher than the state average.
 LNE (Low Number of Events): is used because the rate or percent may be unreliable.
¹Data for mothers who used tobacco are self-reported.
²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

ALL CAUSES	790.8
○ Heart Disease	310.0
Malignant Neoplasms (cancer)	149.2
Trachea, Bronchus, & Lung	38.5
Colon, Rectum, & Anus	LNE
Female Breast	56.9
Pancreas	LNE
Cerebrovascular Disease	25.2
● Chronic Lower Respiratory Diseases	19.2
Accidents	121.8
Motor Vehicle Accidents	91.1
Alzheimer's Disease	20.2
Diabetes Mellitus	23.8
Influenza & Pneumonia	LNE
Intentional Self-Harm (suicide)	LNE
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	LNE

Leading Causes of Death

Total Deaths

1. Heart Disease	43
2. Cancer	18
3. Accidents	7
4. Cerebrovascular Disease	5

Percent of deaths due to tobacco use 10.0
 Median age at death 82


●Denotes a health status indicator which is significantly lower than the state average.
 ○Denotes a health status indicator which is significantly higher than the state average.
³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See Technical Notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Jackson County

Demographic Information



Jackson County is located in western South Dakota and averages 1.6 persons per square mile.

2011 Population Information

Subject	Number	Percent
Total population	3,169	100.0
White	1,360	42.9
Black or African American	16	0.5
American Indian and Alaska Native	1,629	51.4
Asian	3	0.1
Native Hawaiian and Other Pacific Islander	0	0.0
Two or More Races	162	5.1
Under 5 years	292	9.2
Under 18 years	1,033	32.6
65 years and over	421	13.3

Source: United States Census Bureau, 2011 American Community Survey estimates

Health Status Indicators 2007-2011

Nativity

Percent of Low Birth Weight Infants	9.2
● Percent of Mothers Receiving Care in 1st Trimester	47.9
○ Percent of Mothers Who Used Tobacco While Pregnant ¹	25.5
○ Percent of Births Less Than 37 Wks. of Gestation	13.4
● Average Age of Mother	24.8
○ Teenage Birth Rate ²	44.2
Percent White Births	17.4
Percent American Indian Births	76.9
○ Percent Unmarried	76.2
○ Percent WIC births	78.3
● Percent Breastfeeding at discharge	49.1
● Percent Payment-Private Insurance	11.9
○ Percent Payment-Medicaid	68.6
Percent C-Section	23.3

● Denotes a health status indicator which is significantly lower than the state average.
 ○ Denotes a health status indicator which is significantly higher than the state average.
 LNE (Low Number of Events): is used because the rate or percent may be unreliable.
¹Data for mothers who used tobacco are self-reported.
²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

○ ALL CAUSES	1126.9
Heart Disease	205.6
Malignant Neoplasms (cancer)	178.2
Trachea, Bronchus, & Lung	71.2
Colon, Rectum, & Anus	25.7
Female Breast	LNE
Pancreas	LNE
Cerebrovascular Disease	LNE
Chronic Lower Respiratory Diseases	78.7
○ Accidents	164.2
○ Motor Vehicle Accidents	82.6
Alzheimer's Disease	30.5
Diabetes Mellitus	45.0
Influenza & Pneumonia	29.2
Intentional Self-Harm (suicide)	45.0
○ Chronic Liver Disease and Cirrhosis	64.2
Infant Mortality	13.55

Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	29
2. Cancer	26
3. Accidents	22
4. Chronic Lower Respiratory Disease	12
5. Chronic Liver Disease and Cirrhosis	8
6. Diabetes	7
7. Suicide	6
T8. Alzheimer's Disease	5
T8. Influenza and Pneumonia	5

Percent of deaths due to tobacco use 15.7
 Median age at death 72


● Denotes a health status indicator which is significantly lower than the state average.
 ○ Denotes a health status indicator which is significantly higher than the state average.
³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See Technical Notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Jerauld County

Demographic Information



Jerauld County is located in the central region of the state and averages 3.9 persons per square mile.

2011 Population Information

Subject	Number	Percent
Total population	2,085	100.0
White	2,047	98.2
Black or African American	4	0.2
American Indian and Alaska Native	8	0.4
Asian	4	0.2
Native Hawaiian and Other Pacific Islander	2	0.1
Two or More Races	17	0.8
Under 5 years	148	7.1
Under 18 years	450	21.6
65 years and over	517	24.8

Source: United States Census Bureau, 2011 American Community Survey estimates

Health Status Indicators 2007-2011

Natality

Percent of Low Birth Weight Infants	6.4
Percent of Mothers Receiving Care in 1st Trimester	66.4
Percent of Mothers Who Used Tobacco While Pregnant ¹	12.1
• Percent of Births Less Than 37 Wks. of Gestation	3.2
Average Age of Mother	27.9
Teenage Birth Rate ²	17.6
Percent White Births	99.2
Percent American Indian Births	LNE
• Percent Unmarried	20.8
Percent WIC births	32.8
Percent Breastfeeding at discharge	83.2
Percent Payment-Private Insurance	64.0
• Percent Payment-Medicaid	22.4
Percent C-Section	26.4

• Denotes a health status indicator which is significantly lower than the state average.
 ○ Denotes a health status indicator which is significantly higher than the state average.
 LNE (Low Number of Events): is used because the rate or percent may be unreliable.
¹Data for mothers who used tobacco are self-reported.
²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

ALL CAUSES	627.4
• Heart Disease	101.1
Malignant Neoplasms (cancer)	229.3
Trachea, Bronchus, & Lung	36.5
Colon, Rectum, & Anus	32.7
Female Breast	LNE
Pancreas	LNE
Cerebrovascular Disease	50.8
Chronic Lower Respiratory Diseases	48.8
Accidents	30.5
Motor Vehicle Accidents	27.7
○ Alzheimer's Disease	69.6
Diabetes Mellitus	LNE
Influenza & Pneumonia	8.4
Intentional Self-Harm (suicide)	LNE
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	LNE

Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Cancer	38
2. Heart Disease	29
3. Alzheimer's Disease	21
4. Cerebrovascular Disease	13
5. Chronic Lower Respiratory Disease	11

Percent of deaths due to tobacco use 13.7
 Median age at death 84

• Denotes a health status indicator which is significantly lower than the state average.
 ○ Denotes a health status indicator which is significantly higher than the state average.
³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.


See Technical Notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Jones County

Health Status Indicators 2007-2011

Demographic Information



Jones County is located in western South Dakota and averages 1.0 persons per square mile.

2011 Population Information

Subject	Number	Percent
Total population	1,003	100.0
White	949	94.6
Black or African American	2	0.2
American Indian and Alaska Native	24	2.4
Asian	1	0.1
Native Hawaiian and Other Pacific Islander	2	0.2
Two or More Races	25	2.5
Under 5 years	63	6.3
Under 18 years	226	22.5
65 years and over	203	20.2

Source: United States Census Bureau, 2011 American Community Survey estimates

Natality

Percent of Low Birth Weight Infants	8.1
Percent of Mothers Receiving Care in 1st Trimester	80.6
Percent of Mothers Who Used Tobacco While Pregnant ¹	16.1
Percent of Births Less Than 37 Wks. of Gestation	16.1
Average Age of Mother	26.5
Teenage Birth Rate ²	11.3
Percent White Births	95.2
Percent American Indian Births	LNE
• Percent Unmarried	21.0
Percent WIC births	45.2
Percent Breastfeeding at discharge	79.0
Percent Payment-Private Insurance	71.0
Percent Payment-Medicaid	25.8
Percent C-Section	33.9

• Denotes a health status indicator which is significantly lower than the state average.
 ○ Denotes a health status indicator which is significantly higher than the state average.
 LNE (Low Number of Events): is used because the rate or percent may be unreliable.
¹Data for mothers who used tobacco are self-reported.
²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

ALL CAUSES	563.5
• Heart Disease	84.9
Malignant Neoplasms (cancer)	165.5
Trachea, Bronchus, & Lung	53.6
Colon, Rectum, & Anus	LNE
Female Breast	58.9
Pancreas	LNE
Cerebrovascular Disease	LNE
Chronic Lower Respiratory Diseases	56.4
Accidents	LNE
Motor Vehicle Accidents	LNE
Alzheimer's Disease	LNE
Diabetes Mellitus	LNE
Influenza & Pneumonia	26.0
Intentional Self-Harm (suicide)	LNE
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	LNE

Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Cancer	15
2. Heart Disease	8
3. Chronic Lower Respiratory Disease	5
Percent of deaths due to tobacco use	27.1
Median age at death	80


• Denotes a health status indicator which is significantly lower than the state average.
 ○ Denotes a health status indicator which is significantly higher than the state average.
³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See Technical Notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Kingsbury County

Demographic Information



Kingsbury County is located in east central South Dakota and averages 6.2 persons per square mile.

2011 Population Information

Subject	Number	Percent
Total population	5,179	100.0
White	5,070	97.9
Black or African American	10	0.2
American Indian and Alaska Native	31	0.6
Asian	16	0.3
Native Hawaiian and Other Pacific Islander	0	0.0
Two or More Races	52	1.0
Under 5 years	321	6.2
Under 18 years	1,145	22.1
65 years and over	1,113	21.5

Source: United States Census Bureau, 2011 American Community Survey estimates

Health Status Indicators 2007-2011

Nativity

- Percent of Low Birth Weight Infants3.5
- Percent of Mothers Receiving Care in 1st Trimester71.1
- Percent of Mothers Who Used Tobacco While Pregnant¹15.9
- Percent of Births Less Than 37 Wks. of Gestation5.4
- Average Age of Mother27.7
- Teenage Birth Rate²LNE
- Percent White Births96.8
- Percent American Indian Births1.9
- Percent Unmarried23.1
- Percent WIC births26.3
- Percent Breastfeeding at discharge78.5
- Percent Payment-Private Insurance78.5
- Percent Payment-Medicaid18.0
- Percent C-Section23.1

●Denotes a health status indicator which is significantly lower than the state average.
 ○Denotes a health status indicator which is significantly higher than the state average.
 LNE (Low Number of Events): is used because the rate or percent may be unreliable.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

ALL CAUSES	767.4
Heart Disease	180.6
Malignant Neoplasms (cancer)	177.7
Trachea, Bronchus, & Lung	59.0
Colon, Rectum, & Anus	24.0
● Female Breast	5.7
Pancreas	6.7
○ Cerebrovascular Disease	62.8
Chronic Lower Respiratory Diseases	33.3
Accidents	58.4
Motor Vehicle Accidents	31.9
Alzheimer's Disease	37.5
Diabetes Mellitus	17.9
Influenza & Pneumonia	23.7
Intentional Self-Harm (suicide)	17.6
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	9.49

Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	101
2. Cancer	82
3. Cerebrovascular Disease	39
4. Alzheimer's Disease	24
5. Accidents	19
6. Chronic Lower Respiratory Disease	16
7. Influenza and Pneumonia	14
8. Diabetes	9
9. Unspecified Dementia	8
T10. Vascular Dementia	6
T10. Urinary Tract Infection, site not specified	6

Percent of deaths due to tobacco use13.5
 Median age at death83

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See Technical Notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Lake County

Demographic Information



Lake County is located in the east central region of the state and averages 19.9 persons per square mile.

2011 Population Information

Subject	Number	Percent
Total population	11,567	100.0
White	11,162	96.5
Black or African American	81	0.7
American Indian and Alaska Native	93	0.8
Asian	81	0.7
Native Hawaiian and Other Pacific Islander	0	0.0
Two or More Races	150	1.3
Under 5 years	636	5.5
Under 18 years	2,394	20.7
65 years and over	1,978	17.1

Source: United States Census Bureau, 2011 American Community Survey estimates

Health Status Indicators 2007-2011

Nativity

Percent of Low Birth Weight Infants	5.8
Percent of Mothers Receiving Care in 1st Trimester	73.4
Percent of Mothers Who Used Tobacco While Pregnant ¹	17.5
Percent of Births Less Than 37 Weeks of Gestation	7.2
○ Average Age of Mother	27.8
● Teenage Birth Rate ²	7.0
Percent White Births	96.8
Percent American Indian Births	1.7
● Percent Unmarried	23.4
● Percent WIC births	27.4
Percent Breastfeeding at discharge	78.4
Percent Payment-Private Insurance	69.9
● Percent Payment-Medicaid	23.2
Percent C-Section	30.3

●Denotes a health status indicator which is significantly lower than the state average.
 ○Denotes a health status indicator which is significantly higher than the state average.
 LNE (Low Number of Events): is used because the rate or percent may be unreliable.
¹Data for mothers who used tobacco are self-reported.
²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

● ALL CAUSES	619.2
Heart Disease	154.9
Malignant Neoplasms (cancer)	151.9
Trachea, Bronchus, & Lung	33.6
● Colon, Rectum, & Anus	7.8
Female Breast	32.5
Pancreas	13.4
Cerebrovascular Disease	43.0
Chronic Lower Respiratory Diseases	48.2
Accidents	21.5
● Motor Vehicle Accidents	5.7
Alzheimer's Disease	49.3
● Diabetes Mellitus	14.3
● Influenza & Pneumonia	6.4
Intentional Self-Harm (suicide)	12.9
Chronic Liver Disease and Cirrhosis	7.9
Infant Mortality	4.70

Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	132
2. Cancer	124
3. Alzheimer's Disease	47
4. Chronic Lower Respiratory Disease	42
5. Cerebrovascular Disease	38
6. Accidents	17
7. Diabetes	12
8. Essential (primary) hypertension and hypertensive renal disease	9
9. Suicide	7
10. Influenza and Pneumonia	6

Percent of deaths due to tobacco use 19.7
 Median age at death 82


●Denotes a health status indicator which is significantly lower than the state average.
 ○Denotes a health status indicator which is significantly higher than the state average.
³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See Technical Notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Lawrence County

Demographic Information



Lawrence County is located along the Wyoming border and averages 30.1 persons per square mile.

2011 Population Information

Subject	Number	Percent
Total population	24,312	100.0
White	22,902	94.2
Black or African American	170	0.7
American Indian and Alaska Native	559	2.3
Asian	170	0.7
Native Hawaiian and Other Pacific Islander	24	0.1
Two or More Races	486	2.0
Under 5 years	1,337	5.5
Under 18 years	4,644	19.1
65 years and over	4,060	16.7

Source: United States Census Bureau, 2011 American Community Survey estimates

Health Status Indicators 2007-2011

Natality

Percent of Low Birth Weight Infants	7.3
○ Percent of Mothers Receiving Care in 1st Trimester	79.7
Percent of Mothers Who Used Tobacco While Pregnant ¹	20.8
Percent of Births Less Than 37 Weeks of Gestation	8.3
Average Age of Mother	27.1
Teenage Birth Rate ²	13.0
Percent White Births	93.8
Percent American Indian Births	2.5
Percent Unmarried	35.3
Percent WIC births	38.4
○ Percent Breastfeeding at discharge	80.0
Percent Payment-Private Insurance	53.0
Percent Payment-Medicaid	38.1
● Percent C-Section	21.9

● Denotes a health status indicator which is significantly lower than the state average.
 ○ Denotes a health status indicator which is significantly higher than the state average.
 LNE (Low Number of Events): is used because the rate or percent may be unreliable.
¹Data for mothers who used tobacco are self-reported.
²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

● ALL CAUSES	630.3
Heart Disease	138.5
Malignant Neoplasms (cancer)	165.0
Trachea, Bronchus, & Lung	40.4
Colon, Rectum, & Anus	15.0
Female Breast	33.4
Pancreas	12.6
Cerebrovascular Disease	35.4
Chronic Lower Respiratory Diseases	51.4
Accidents	32.0
Motor Vehicle Accidents	11.4
● Alzheimer's Disease	24.3
Diabetes Mellitus	22.9
● Influenza & Pneumonia	7.6
Intentional Self-Harm (suicide)	16.0
Chronic Liver Disease and Cirrhosis	8.3
Infant Mortality	4.73

Leading Causes of Death

Total Deaths

1. Cancer	263
2. Heart Disease	232
3. Chronic Lower Respiratory Disease	84
4. Cerebrovascular Disease	60
5. Alzheimer's Disease	43
6. Accidents	42
7. Diabetes	38
8. Suicide	21
T9. Influenza and Pneumonia	14
T9. Unspecified dementia	14

Percent of deaths due to tobacco use 18.9
 Median age at death 80


● Denotes a health status indicator which is significantly lower than the state average.
 ○ Denotes a health status indicator which is significantly higher than the state average.
³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See Technical Notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Lincoln County

Demographic Information



Lincoln County is located in southeastern South Dakota and averages 77.7 persons per square mile.

2011 Population Information

Subject	Number	Percent
Total population	46,793	100.0
White	44,921	96.0
Black or African American	374	0.8
American Indian and Alaska Native	328	0.7
Asian	515	1.1
Native Hawaiian and Other Pacific Islander	0	0.0
Two or More Races	655	1.4
Under 5 years	4,399	9.4
Under 18 years	13,664	29.2
65 years and over	4,305	9.2

Source: United States Census Bureau, 2011 American Community Survey estimates

Health Status Indicators 2007-2011

Natality

Percent of Low Birth Weight Infants	6.3
○ Percent of Mothers Receiving Care in 1st Trimester	78.3
● Percent of Mothers Who Used Tobacco While Pregnant ¹	8.1
Percent of Births Less Than 37 Weeks of Gestation	8.7
○ Average Age of Mother	28.7
● Teenage Birth Rate ²	6.8
Percent White Births	97.6
Percent American Indian Births	0.5
● Percent Unmarried	15.3
● Percent WIC births	10.8
○ Percent Breastfeeding at discharge	78.6
○ Percent Payment-Private Insurance	85.3
● Percent Payment-Medicaid	11.8
Percent C-Section	27.5

- Denotes a health status indicator which is significantly lower than the state average.
- Denotes a health status indicator which is significantly higher than the state average.
- LNE (Low Number of Events): is used because the rate or percent may be unreliable.
- ¹Data for mothers who used tobacco are self-reported.
- ²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

● ALL CAUSES	631.3
Heart Disease	140.4
Malignant Neoplasms (cancer)	148.6
Trachea, Bronchus, & Lung	45.7
Colon, Rectum, & Anus	13.4
Female Breast	18.4
Pancreas	6.2
Cerebrovascular Disease	37.2
Chronic Lower Respiratory Diseases	40.8
Accidents	33.0
● Motor Vehicle Accidents	9.1
Alzheimer's Disease	39.4
Diabetes Mellitus	26.4
● Influenza & Pneumonia	8.8
● Intentional Self-Harm (suicide)	7.6
● Chronic Liver Disease and Cirrhosis	2.9
Infant Mortality	4.29

Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Cancer	225
2. Heart Disease	209
3. Alzheimer's Disease	58
T4. Chronic Lower Respiratory Disease	56
T4. Accidents	56
6. Cerebrovascular Disease	53
7. Diabetes	37
8. Unspecified Dementia	18
9. Essential (primary) hypertension and hypertensive renal disease	17
10. Suicide	16

Percent of deaths due to tobacco use 15.7
 Median age at death 82


- Denotes a health status indicator which is significantly lower than the state average.
- Denotes a health status indicator which is significantly higher than the state average.
- ³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See Technical Notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics.

Lyman County

Demographic Information



Lyman County is located in central South Dakota and averages 2.3 persons per square mile.

2011 Population Information

Subject	Number	Percent
Total population	3,806	100.0
White	2,253	59.2
Black or African American	8	0.2
American Indian and Alaska Native	1,427	37.5
Asian	11	0.3
Native Hawaiian and Other Pacific Islander	0	0.0
Two or More Races	107	0.8
Under 5 years	263	6.9
Under 18 years	1,085	28.5
65 years and over	563	14.8

Source: United States Census Bureau, 2011 American Community Survey estimates

Health Status Indicators 2007-2011

Natality

Percent of Low Birth Weight Infants	5.3
● Percent of Mothers Receiving Care in 1st Trimester	52.0
○ Percent of Mothers Who Used Tobacco While Pregnant ¹	24.8
Percent of Births Less Than 37 Weeks of Gestation	7.2
● Average Age of Mother	25.8
○ Teenage Birth Rate ²	62.1
Percent White Births	36.6
Percent American Indian Births	60.0
○ Percent Unmarried	62.5
○ Percent WIC births	65.9
● Percent Breastfeeding at discharge	62.5
● Percent Payment-Private Insurance	35.9
○ Percent Payment-Medicaid	59.7
○ Percent C-Section	33.1

● Denotes a health status indicator which is significantly lower than the state average.
 ○ Denotes a health status indicator which is significantly higher than the state average.
 LNE (Low Number of Events): is used because the rate or percent may be unreliable.
¹Data for mothers who used tobacco are self-reported.
²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

ALL CAUSES	757.6
Heart Disease	173.9
Malignant Neoplasms (cancer)	133.6
Trachea, Bronchus, & Lung	76.3
Colon, Rectum, & Anus	LNE
Female Breast	LNE
Pancreas	LNE
Cerebrovascular Disease	28.6
Chronic Lower Respiratory Diseases	52.4
○ Accidents	138.4
○ Motor Vehicle Accidents	82.3
Alzheimer's Disease	43.5
Diabetes Mellitus	13.2
Influenza & Pneumonia	18.0
Intentional Self-Harm (suicide)	LNE
Chronic Liver Disease and Cirrhosis	18.0
Infant Mortality	LNE

Leading Causes of Death

Total Deaths

1. Heart Disease	35
2. Cancer	29
3. Accidents	25
4. Chronic Lower Respiratory Disease	11
5. Alzheimer's Disease	9
6. Cerebrovascular Disease	6

Percent of deaths due to tobacco use 24.0
 Median age at death 73


● Denotes a health status indicator which is significantly lower than the state average.
 ○ Denotes a health status indicator which is significantly higher than the state average.
³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See Technical Notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

McCook County

Demographic Information



McCook County is located in eastern South Dakota and averages 9.8 persons per square mile.

2011 Population Information

Subject	Number	Percent
Total population	5,556	100.0
White	5,456	98.2
Black or African American	17	0.3
American Indian and Alaska Native	33	0.6
Asian	11	0.2
Native Hawaiian and Other Pacific Islander	6	0.1
Two or More Races	39	0.7
Under 5 years	406	7.3
Under 18 years	1,422	25.6
65 years and over	1,056	19.0

Source: United States Census Bureau, 2011 American Community Survey estimates

Health Status Indicators 2007-2011

Natality

Percent of Low Birth Weight Infants	6.7
Percent of Mothers Receiving Care in 1st Trimester	69.1
● Percent of Mothers Who Used Tobacco While Pregnant ¹	11.6
Percent of Births Less Than 37 Weeks of Gestation	9.7
○ Average Age of Mother	28.6
Teenage Birth Rate ²	LNE
Percent White Births	98.4
Percent American Indian Births	LNE
● Percent Unmarried	18.8
● Percent WIC births	31.1
Percent Breastfeeding at discharge	76.7
○ Percent Payment-Private Insurance	75.3
● Percent Payment-Medicaid	20.1
Percent C-Section	24.4

● Denotes a health status indicator which is significantly lower than the state average.
 ○ Denotes a health status indicator which is significantly higher than the state average.
 LNE (Low Number of Events): is used because the rate or percent may be unreliable.
¹Data for mothers who used tobacco are self-reported.
²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

○ ALL CAUSES	1001.9
○ Heart Disease	258.0
○ Malignant Neoplasms (cancer)	252.2
Trachea, Bronchus, & Lung	71.9
Colon, Rectum, & Anus	10.0
Female Breast	25.8
Pancreas	LNE
○ Cerebrovascular Disease	74.8
Chronic Lower Respiratory Diseases	76.1
Accidents	25.0
Motor Vehicle Accidents	12.5
Alzheimer's Disease	52.2
Diabetes Mellitus	22.8
Influenza & Pneumonia	32.3
Intentional Self-Harm (suicide)	13.0
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	LNE

Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	102
2. Cancer	89
3. Cerebrovascular Disease	29
4. Chronic Lower Respiratory Disease	28
5. Alzheimer's Disease	23
6. Influenza and Pneumonia	13
7. Diabetes	9
8. Accidents	6
T9. Nephritis, nephrotic syndrome, and nephrosis	5
T9. Parkinson's Disease	5

Percent of deaths due to tobacco use 17.2
 Median age at death 82


● Denotes a health status indicator which is significantly lower than the state average.
 ○ Denotes a health status indicator which is significantly higher than the state average.
³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See Technical Notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

McPherson County

Demographic Information



McPherson County is located in the north central region of the state and averages 2.2 persons per square mile.

2011 Population Information

Subject	Number	Percent
Total population	2,452	100.0
White	2,420	98.7
Black or African American	0	0.0
American Indian and Alaska Native	2	0.1
Asian	5	0.2
Native Hawaiian and Other Pacific Islander	2	0.1
Two or More Races	20	0.8
Under 5 years	130	5.3
Under 18 years	535	21.8
65 years and over	733	29.9

Source: United States Census Bureau, 2011 American Community Survey estimates

Health Status Indicators 2007-2011

Nativity

Percent of Low Birth Weight Infants	8.7
Percent of Mothers Receiving Care in 1st Trimester	65.2
Percent of Mothers Who Used Tobacco While Pregnant ¹	12.1
Percent of Births Less Than 37 Weeks of Gestation	7.6
○ Average Age of Mother	30.4
Teenage Birth Rate ²	LNE
Percent White Births	96.7
Percent American Indian Births	0.0
● Percent Unmarried	17.4
● Percent WIC births	19.6
Percent Breastfeeding at discharge	80.4
○ Percent Payment-Private Insurance	76.1
● Percent Payment-Medicaid	15.2
Percent C-Section	25.0

●Denotes a health status indicator which is significantly lower than the state average.
 ○Denotes a health status indicator which is significantly higher than the state average.
 LNE (Low Number of Events): is used because the rate or percent may be unreliable.
¹Data for mothers who used tobacco are self-reported.
²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

● ALL CAUSES	339.0
● Heart Disease	108.9
● Malignant Neoplasms (cancer)	44.5
● Trachea, Bronchus, & Lung	19.3
Colon, Rectum, & Anus	LNE
Female Breast	LNE
Pancreas	LNE
Cerebrovascular Disease	40.7
● Chronic Lower Respiratory Diseases	9.9
Accidents	21.9
Motor Vehicle Accidents	13.4
● Alzheimer's Disease	13.0
Diabetes Mellitus	16.0
Influenza & Pneumonia	8.0
Intentional Self-Harm (suicide)	LNE
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	32.61

Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	56
2. Cerebrovascular Disease	20
3. Cancer	19
4. Essential (primary) hypertension and hypertensive renal disease	10
T5. Alzheimer's Disease	8
T5. Accidents	8
T5. Diabetes	8
8. Chronic Lower Respiratory Disease	6

Percent of deaths due to tobacco use 7.9
 Median age at death 85


●Denotes a health status indicator which is significantly lower than the state average.
 ○Denotes a health status indicator which is significantly higher than the state average.
³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See Technical Notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Marshall County

Demographic Information



Marshall County is located in the northeastern part of the state and averages 5.6 persons per square mile.

2011 Population Information

Subject	Number	Percent
Total population	4,597	100.0
White	4,064	88.4
Black or African American	18	0.4
American Indian and Alaska Native	437	9.5
Asian	14	0.3
Native Hawaiian and Other Pacific Islander	0	0.0
Two or More Races	60	1.3
Under 5 years	285	6.2
Under 18 years	1,011	22.0
65 years and over	878	19.1

Source: United States Census Bureau, 2011 American Community Survey estimates

Health Status Indicators 2007-2011

Nativity

Percent of Low Birth Weight Infants	5.9
Percent of Mothers Receiving Care in 1st Trimester	61.8
Percent of Mothers Who Used Tobacco While Pregnant ¹	18.4
Percent of Births Less Than 37 Weeks of Gestation	6.3
Average Age of Mother	26.8
Teenage Birth Rate ²	15.6
Percent White Births	79.2
Percent American Indian Births	20.0
Percent Unmarried	33.0
Percent WIC births	31.9
Percent Breastfeeding at discharge	73.3
Percent Payment-Private Insurance	58.9
Percent Payment-Medicaid	29.3
Percent C-Section	23.7

● Denotes a health status indicator which is significantly lower than the state average.

○ Denotes a health status indicator which is significantly higher than the state average.

LNE (Low Number of Events): is used because the rate or percent may be unreliable.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

○ ALL CAUSES	1070.1
Heart Disease	201.0
○ Malignant Neoplasms (cancer)	242.8
Trachea, Bronchus, & Lung	53.5
Colon, Rectum, & Anus	33.6
Female Breast	27.8
Pancreas	30.7
○ Cerebrovascular Disease	75.3
Chronic Lower Respiratory Diseases	69.1
Accidents	86.3
Motor Vehicle Accidents	36.4
Alzheimer's Disease	48.5
Diabetes Mellitus	51.4
Influenza & Pneumonia	14.0
Intentional Self-Harm (suicide)	LNE
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	18.52

Leading Causes of Death

Total Deaths

1. Cancer	64
2. Heart Disease	61
3. Cerebrovascular Disease	27
4. Chronic Lower Respiratory Disease	25
5. Alzheimer's Disease	21
T6. Accidents	9
T6. Diabetes	9
8. Influenza and Pneumonia	6

Percent of deaths due to tobacco use	15.2
Median age at death	83

● Denotes a health status indicator which is significantly lower than the state average.

○ Denotes a health status indicator which is significantly higher than the state average.


³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See Technical Notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Meade County

Demographic Information



Meade County is located in west central South Dakota and averages 7.3 persons per square mile.

2011 Population Information

Subject	Number	Percent
Total population	25,546	100.0
White	23,553	92.2
Black or African American	434	1.7
American Indian and Alaska Native	639	2.5
Asian	204	0.8
Native Hawaiian and Other Pacific Islander	26	0.1
Two or More Races	715	2.8
Under 5 years	1,865	7.3
Under 18 years	6,208	24.3
65 years and over	3,168	12.4

Source: United States Census Bureau, 2011 American Community Survey estimates

Health Status Indicators 2007-2011

Nativity

Percent of Low Birth Weight Infants	6.7
○ Percent of Mothers Receiving Care in 1st Trimester	77.7
Percent of Mothers Who Used Tobacco While Pregnant ¹	15.9
Percent of Births Less Than 37 Weeks of Gestation	8.0
Average Age of Mother	27.4
● Teenage Birth Rate ²	12.0
Percent White Births	91.3
Percent American Indian Births	3.9
● Percent Unmarried	26.4
● Percent WIC births	35.3
○ Percent Breastfeeding at discharge	82.5
Percent Payment-Private Insurance	50.7
● Percent Payment-Medicaid	29.8
● Percent C-Section	23.5

●Denotes a health status indicator which is significantly lower than the state average.
 ○Denotes a health status indicator which is significantly higher than the state average.
 LNE (Low Number of Events): is used because the rate or percent may be unreliable.
¹Data for mothers who used tobacco are self-reported.
²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

○ ALL CAUSES	778.9
Heart Disease	169.9
Malignant Neoplasms (cancer)	178.7
Trachea, Bronchus, & Lung	55.0
● Colon, Rectum, & Anus	8.2
Female Breast	20.2
Pancreas	13.5
Cerebrovascular Disease	47.0
Chronic Lower Respiratory Diseases	57.1
Accidents	56.0
Motor Vehicle Accidents	22.5
Alzheimer's Disease	26.2
Diabetes Mellitus	27.1
Influenza & Pneumonia	20.1
Intentional Self-Harm (suicide)	20.9
Chronic Liver Disease and Cirrhosis	10.2
Infant Mortality	8.92

Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Cancer	221
2. Heart Disease	202
T3. Chronic Lower Respiratory Disease	69
T3. Accidents	69
5. Cerebrovascular Disease	54
6. Diabetes	33
7. Alzheimer's Disease	29
8. Suicide	25
9. Influenza and Pneumonia	23
10. Chronic Liver Disease and Cirrhosis	15

Percent of deaths due to tobacco use 20.3
 Median age at death 76


●Denotes a health status indicator which is significantly lower than the state average.
 ○Denotes a health status indicator which is significantly higher than the state average.
³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See Technical Notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Mellette County

Demographic Information



Mellette County is located in the south central region of the state and averages 1.6 persons per square mile.

2011 Population Information

Subject	Number	Percent
Total population	2,067	100.0
White	821	39.7
Black or African American	4	0.2
American Indian and Alaska Native	1,110	53.7
Asian	6	0.3
Native Hawaiian and Other Pacific Islander	0	0.0
Two or More Races	126	6.1
Under 5 years	192	9.3
Under 18 years	666	32.2
65 years and over	539	12.9

Source: United States Census Bureau, 2011 American Community Survey estimates

Health Status Indicators 2007-2011

Natality

Percent of Low Birth Weight Infants	7.0
● Percent of Mothers Receiving Care in 1st Trimester	44.4
Percent of Mothers Who Used Tobacco While Pregnant ¹	21.5
Percent of Births Less Than 37 Weeks of Gestation	8.6
● Average Age of Mother	24.1
○ Teenage Birth Rate ²	70.0
Percent White Births	24.3
Percent American Indian Births	62.7
○ Percent Unmarried	69.9
○ Percent WIC births	66.7
Percent Breastfeeding at discharge	69.4
● Percent Payment-Private Insurance	14.5
○ Percent Payment-Medicaid	57.0
○ Percent C-Section	35.5

● Denotes a health status indicator which is significantly lower than the state average.

○ Denotes a health status indicator which is significantly higher than the state average.

LNE (Low Number of Events): is used because the rate or percent may be unreliable.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

○ ALL CAUSES	1067.0
Heart Disease	172.5
Malignant Neoplasms (cancer)	222.8
Trachea, Bronchus, & Lung	61.0
Colon, Rectum, & Anus	24.3
Female Breast	LNE
Pancreas	LNE
Cerebrovascular Disease	58.7
Chronic Lower Respiratory Diseases	92.4
○ Accidents	133.6
○ Motor Vehicle Accidents	73.7
Alzheimer's Disease	23.8
Diabetes Mellitus	34.8
Influenza & Pneumonia	LNE
Intentional Self-Harm (suicide)	LNE
○ Chronic Liver Disease and Cirrhosis	79.8
Infant Mortality	LNE

Leading Causes of Death

Total Deaths

1. Cancer	25
2. Heart Disease	20
3. Accidents	14
4. Chronic Lower Respiratory Disease	10
T5. Cerebrovascular Disease	7
T5. Chronic Liver Disease and Cirrhosis	7

Percent of deaths due to tobacco use	22.0
Median age at death	74

● Denotes a health status indicator which is significantly lower than the state average.

○ Denotes a health status indicator which is significantly higher than the state average.


³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See Technical Notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Miner County

Demographic Information



Miner County is located in the east central region of the state and averages 4.2 persons per square mile.

2011 Population Information

Subject	Number	Percent
Total population	2,359	100.0
White	2,319	98.3
Black or African American	2	0.1
American Indian and Alaska Native	5	0.2
Asian	9	0.4
Native Hawaiian and Other Pacific Islander	0	0.0
Two or More Races	24	1.0
Under 5 years	151	6.4
Under 18 years	552	23.4
65 years and over	526	22.3

Source: United States Census Bureau, 2011 American Community Survey estimates

Health Status Indicators 2007-2011

Natality

Percent of Low Birth Weight Infants	LNE
Percent of Mothers Receiving Care in 1st Trimester	71.4
Percent of Mothers Who Used Tobacco While Pregnant ¹	12.6
Percent of Births Less Than 37 Weeks of Gestation	4.7
○ Average Age of Mother	28.6
Teenage Birth Rate ²	12.0
Percent White Births	95.3
Percent American Indian Births	3.1
● Percent Unmarried	16.4
● Percent WIC births	24.2
Percent Breastfeeding at discharge	75.8
○ Percent Payment-Private Insurance	78.1
● Percent Payment-Medicaid	18.8
Percent C-Section	31.3

● Denotes a health status indicator which is significantly lower than the state average.
 ○ Denotes a health status indicator which is significantly higher than the state average.
 LNE (Low Number of Events): is used because the rate or percent may be unreliable.
¹Data for mothers who used tobacco are self-reported.
²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

ALL CAUSES	678.4
Heart Disease	174.2
Malignant Neoplasms (cancer)	194.6
Trachea, Bronchus, & Lung	62.0
Colon, Rectum, & Anus	22.1
Female Breast	LNE
Pancreas	17.4
Cerebrovascular Disease	39.8
Chronic Lower Respiratory Diseases	49.8
Accidents	28.4
Motor Vehicle Accidents	24.1
○ Alzheimer's Disease	70.8
Diabetes Mellitus	LNE
Influenza & Pneumonia	9.8
Intentional Self-Harm (suicide)	LNE
Chronic Liver Disease and Cirrhosis	13.9
Infant Mortality	LNE

Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	49
2. Cancer	47
3. Alzheimer's Disease	23
4. Cerebrovascular Disease	13
5. Chronic Lower Respiratory Disease	12
6. Parkinson's Disease	5

Percent of deaths due to tobacco use 18.1
 Median age at death 86


● Denotes a health status indicator which is significantly lower than the state average.
 ○ Denotes a health status indicator which is significantly higher than the state average.
³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See Technical Notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Minnehaha County

Demographic Information



Minnehaha County is located in southeastern South Dakota and averages 210.0 persons per square mile.

2011 Population Information

Subject	Number	Percent
Total population	171,752	100.0
White	154,062	89.7
Black or African American	6,527	3.8
American Indian and Alaska Native	4,809	2.8
Asian	2,576	1.5
Native Hawaiian and Other Pacific Islander	172	0.1
Two or More Races	3,607	2.1
Under 5 years	12,881	7.5
Under 18 years	42,766	24.9
65 years and over	19,408	11.3

Source: United States Census Bureau, 2011 American Community Survey estimates

Health Status Indicators 2007-2011

Natality

Percent of Low Birth Weight Infants	7.1
Percent of Mothers Receiving Care in 1st Trimester	68.5
• Percent of Mothers Who Used Tobacco While Pregnant ¹	16.3
Percent of Births Less Than 37 Weeks of Gestation	9.1
○ Average Age of Mother	27.5
Teenage Birth Rate ²	16.5
Percent White Births	85.0
Percent American Indian Births	4.5
Percent Unmarried	36.9
• Percent WIC births	30.6
Percent Breastfeeding at discharge	73.6
○ Percent Payment-Private Insurance	62.4
• Percent Payment-Medicaid	32.5
Percent C-Section	27.1

•Denotes a health status indicator which is significantly lower than the state average.
 ○Denotes a health status indicator which is significantly higher than the state average.
 LNE (Low Number of Events): is used because the rate or percent may be unreliable.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

ALL CAUSES	686.0
• Heart Disease	146.4
Malignant Neoplasms (cancer)	170.6
Trachea, Bronchus, & Lung	47.5
Colon, Rectum, & Anus	14.6
Female Breast	19.8
Pancreas	10.1
Cerebrovascular Disease	40.7
Chronic Lower Respiratory Diseases	50.7
Accidents	31.5
• Motor Vehicle Accidents	8.6
Alzheimer's Disease	39.8
• Diabetes Mellitus	15.9
Influenza & Pneumonia	12.7
Intentional Self-Harm (suicide)	13.5
Chronic Liver Disease and Cirrhosis	9.8
Infant Mortality	6.34

Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Cancer	1,499
2. Heart Disease	1,347
3. Chronic Lower Respiratory Disease	444
4. Alzheimer's Disease	373
5. Cerebrovascular Disease	368
6. Accidents	282
7. Diabetes	145
8. Suicide	122
9. Influenza and Pneumonia	117
10. Chronic Liver Disease and Cirrhosis	89

Percent of deaths due to tobacco use 18.1
 Median age at death 80

•Denotes a health status indicator which is significantly lower than the state average.
 ○Denotes a health status indicator which is significantly higher than the state average.


³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See Technical Notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Moody County

Demographic Information



Moody County is located on the Minnesota border and averages 12.5 persons per square mile.

2011 Population Information

Subject	Number	Percent
Total population	6,475	100.0
White	5,303	81.9
Black or African American	39	0.6
American Indian and Alaska Native	900	13.9
Asian	71	1.1
Native Hawaiian and Other Pacific Islander	0	0.0
Two or More Races	162	2.5
Under 5 years	473	7.3
Under 18 years	1,671	25.8
65 years and over	991	15.3

Source: United States Census Bureau, 2011 American Community Survey estimates

Health Status Indicators 2007-2011

Nativity

Percent of Low Birth Weight Infants	6.7
• Percent of Mothers Receiving Care in 1st Trimester	60.6
Percent of Mothers Who Used Tobacco While Pregnant ¹	20.8
Percent of Births Less Than 37 Weeks of Gestation	10.7
Average Age of Mother	27.2
Teenage Birth Rate ²	13.2
Percent White Births	78.4
Percent American Indian Births	16.0
Percent Unmarried	38.0
Percent WIC births	40.0
Percent Breastfeeding at discharge	70.8
Percent Payment-Private Insurance	57.9
Percent Payment-Medicaid	32.2
Percent C-Section	27.1

•Denotes a health status indicator which is significantly lower than the state average.
 ○Denotes a health status indicator which is significantly higher than the state average.
 LNE (Low Number of Events): is used because the rate or percent may be unreliable.
¹Data for mothers who used tobacco are self-reported.
²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

ALL CAUSES	645.7
Heart Disease	177.0
Malignant Neoplasms (cancer)	160.9
Trachea, Bronchus, & Lung	32.1
Colon, Rectum, & Anus	24.1
Female Breast	26.8
Pancreas	12.3
Cerebrovascular Disease	37.5
• Chronic Lower Respiratory Diseases	26.0
Accidents	36.4
Motor Vehicle Accidents	12.2
○ Alzheimer's Disease	64.0
Diabetes Mellitus	30.5
Influenza & Pneumonia	LNE
Intentional Self-Harm (suicide)	14.3
Chronic Liver Disease and Cirrhosis	10.7
Infant Mortality	6.94

Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	74
2. Cancer	64
3. Alzheimer's Disease	29
4. Cerebrovascular Disease	15
5. Accidents	14
6. Diabetes	12
7. Chronic Lower Respiratory Disease	10
8. Suicide	5

Percent of deaths due to tobacco use 19.8
 Median age at death..... 80


•Denotes a health status indicator which is significantly lower than the state average.
 ○Denotes a health status indicator which is significantly higher than the state average.
³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See Technical Notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Pennington County

Demographic Information



Pennington County is located on the Wyoming border and averages 36.4 persons per square mile.

2011 Population Information

Subject	Number	Percent
Total population	102,815	100.0
White	86,673	84.3
Black or African American	1,337	1.3
American Indian and Alaska Native	9,973	9.7
Asian	1,131	1.1
Native Hawaiian and Other Pacific Islander	103	0.1
Two or More Races	3,599	3.5
Under 5 years	7,505	7.3
Under 18 years	24,984	24.3
65 years and over	14,291	13.9

Source: United States Census Bureau, 2011 American Community Survey estimates

Health Status Indicators 2007-2011

Nativity

Percent of Low Birth Weight Infants	7.1
○ Percent of Mothers Receiving Care in 1st Trimester	72.2
○ Percent of Mothers Who Used Tobacco While Pregnant ¹	19.9
Percent of Births Less Than 37 Weeks of Gestation	8.5
● Average Age of Mother	26.3
● Teenage Birth Rate ²	21.2
● Percent White Births	72.5
● Percent American Indian Births	19.0
○ Percent Unmarried	43.6
○ Percent WIC births	46.1
○ Percent Breastfeeding at discharge	78.2
● Percent Payment-Private Insurance	37.3
○ Percent Payment-Medicaid	43.8
● Percent C-Section	20.2

●Denotes a health status indicator which is significantly lower than the state average.
 ○Denotes a health status indicator which is significantly higher than the state average.
 LNE (Low Number of Events): is used because the rate or percent may be unreliable.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

● ALL CAUSES	642.8
● Heart Disease	143.7
Malignant Neoplasms (cancer)	162.9
Trachea, Bronchus, & Lung	45.4
Colon, Rectum, & Anus	12.9
Female Breast	24.4
Pancreas	10.4
● Cerebrovascular Disease	27.7
Chronic Lower Respiratory Diseases	44.1
Accidents	40.6
● Motor Vehicle Accidents	9.7
Alzheimer's Disease	30.2
● Diabetes Mellitus	11.7
Influenza & Pneumonia	16.3
Intentional Self-Harm (suicide)	17.8
● Chronic Liver Disease and Cirrhosis	6.0
Infant Mortality	7.07

Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Cancer	903
2. Heart Disease	814
3. Chronic Lower Respiratory Disease	246
4. Accidents	214
5. Alzheimer's Disease	175
6. Cerebrovascular Disease	155
7. Influenza and Pneumonia	92
8. Suicide	91
9. Unspecified dementia	85
10. Diabetes	66

Percent of deaths due to tobacco use 16.3
 Median age at death 78

●Denotes a health status indicator which is significantly lower than the state average.
 ○Denotes a health status indicator which is significantly higher than the state average.


³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See Technical Notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Perkins County

Demographic Information



Perkins County is located in northwestern South Dakota and averages 1.0 persons per square mile.

2011 Population Information

Subject	Number	Percent
Total population	3,001	100.0
White	2,905	96.8
Black or African American	9	0.3
American Indian and Alaska Native	45	1.5
Asian	6	0.2
Native Hawaiian and Other Pacific Islander	0	0.0
Two or More Races	36	1.2
Under 5 years	159	5.3
Under 18 years	636	21.2
65 years and over	672	22.4

Source: United States Census Bureau, 2011 American Community Survey estimates

Health Status Indicators 2007-2011

Natality

• Percent of Low Birth Weight Infants	3.3
Percent of Mothers Receiving Care in 1st Trimester	66.4
• Percent of Mothers Who Used Tobacco While Pregnant ¹	10.2
Percent of Births Less Than 37 Weeks of Gestation	5.9
○ Average Age of Mother	28.2
Teenage Birth Rate ²	LNE
Percent White Births	97.4
Percent American Indian Births	2.0
• Percent Unmarried	14.4
Percent WIC births	39.9
Percent Breastfeeding at discharge	79.7
Percent Payment-Private Insurance	58.8
• Percent Payment-Medicaid	22.2
Percent C-Section	30.1

- Denotes a health status indicator which is significantly lower than the state average.
- Denotes a health status indicator which is significantly higher than the state average.
- LNE (Low Number of Events): is used because the rate or percent may be unreliable.
- ¹Data for mothers who used tobacco are self-reported.
- ²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

ALL CAUSES	782.0
Heart Disease	151.2
Malignant Neoplasms (cancer)	192.3
Trachea, Bronchus, & Lung	41.2
Colon, Rectum, & Anus	29.9
Female Breast	33.5
Pancreas	13.7
Cerebrovascular Disease	33.4
Chronic Lower Respiratory Diseases	56.4
Accidents	72.0
Motor Vehicle Accidents	20.5
Alzheimer's Disease	33.8
Diabetes Mellitus	20.8
Influenza & Pneumonia	LNE
Intentional Self-Harm (suicide)	LNE
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	LNE

Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Cancer	56
2. Heart Disease	51
3. Chronic Lower Respiratory Disease	19
4. Accidents	15
5. Cerebrovascular Disease	13
6. Alzheimer's Disease	12
7. Diabetes	7
8. Unspecified dementia	6
9. Septicemia	5

Percent of deaths due to tobacco use 16.8
 Median age at death 83


- Denotes a health status indicator which is significantly lower than the state average.
- Denotes a health status indicator which is significantly higher than the state average.
- ³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See Technical Notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Potter County

Demographic Information



Potter County is located in north central South Dakota and averages 2.7 persons per square mile.

2011 Population Information

Subject	Number	Percent
Total population	2,364	100.0
White	2,303	97.4
Black or African American	5	0.2
American Indian and Alaska Native	21	0.9
Asian	9	0.4
Native Hawaiian and Other Pacific Islander	0	0.0
Two or More Races	26	1.1
Under 5 years	132	5.6
Under 18 years	478	20.2
65 years and over	610	25.8

Source: United States Census Bureau, 2011 American Community Survey estimates

Health Status Indicators 2007-2011

Natality

Percent of Low Birth Weight Infants	3.9
Percent of Mothers Receiving Care in 1st Trimester	74.2
• Percent of Mothers Who Used Tobacco While Pregnant ¹	9.4
Percent of Births Less Than 37 Weeks of Gestation	5.4
Average Age of Mother	27.6
Teenage Birth Rate ²	16.2
Percent White Births	89.9
Percent American Indian Births	7.8
• Percent Unmarried	24.0
• Percent WIC births	27.1
Percent Breastfeeding at discharge	66.7
Percent Payment-Private Insurance	69.0
• Percent Payment-Medicaid	24.8
Percent C-Section	31.8

• Denotes a health status indicator which is significantly lower than the state average.
 ○ Denotes a health status indicator which is significantly higher than the state average.
 LNE (Low Number of Events): is used because the rate or percent may be unreliable.
¹Data for mothers who used tobacco are self-reported.
²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

ALL CAUSES	690.6
Heart Disease	119.2
Malignant Neoplasms (cancer)	198.5
Trachea, Bronchus, & Lung	45.7
Colon, Rectum, & Anus	18.3
Female Breast	37.6
Pancreas	13.1
Cerebrovascular Disease	28.8
Chronic Lower Respiratory Diseases	37.8
Accidents	42.2
Motor Vehicle Accidents	LNE
Alzheimer's Disease	28.5
Diabetes Mellitus	40.3
Influenza & Pneumonia	30.8
Intentional Self-Harm (suicide)	13.9
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	LNE

Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Cancer	49
2. Heart Disease	36
3. Chronic Lower Respiratory Disease	12
4. Influenza and Pneumonia	11
T5. Alzheimer's Disease	10
T5. Diabetes	10
7. Accidents	8
T8. Cerebrovascular Disease	7
T8. Parkinson's Disease	7

Percent of deaths due to tobacco use 19.0
 Median age at death 83


• Denotes a health status indicator which is significantly lower than the state average.
 ○ Denotes a health status indicator which is significantly higher than the state average.
³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See Technical Notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Roberts County

Demographic Information



Roberts County is located in the extreme northeastern corner of the state and averages 9.2 persons per square mile.

2011 Population Information

Subject	Number	Percent
Total population	10,286	100.0
White	6,346	61.7
Black or African American	31	0.3
American Indian and Alaska Native	3,559	34.6
Asian	31	0.3
Native Hawaiian and Other Pacific Islander	0	0.0
Two or More Races	319	3.1
Under 5 years	854	8.3
Under 18 years	2,942	28.6
65 years and over	1,759	17.1

Source: United States Census Bureau, 2011 American Community Survey estimates

Health Status Indicators 2007-2011

Natality

Percent of Low Birth Weight Infants	6.8
● Percent of Mothers Receiving Care in 1st Trimester	52.7
○ Percent of Mothers Who Used Tobacco While Pregnant ¹	28.8
Percent of Births Less Than 37 Weeks of Gestation	9.5
● Average Age of Mother	25.2
○ Teenage Birth Rate ²	33.9
Percent White Births	37.7
Percent American Indian Births	55.7
○ Percent Unmarried	60.4
○ Percent WIC births	57.6
● Percent Breastfeeding at discharge	62.0
● Percent Payment-Private Insurance	30.9
○ Percent Payment-Medicaid	57.3
Percent C-Section	24.3

● Denotes a health status indicator which is significantly lower than the state average.
 ○ Denotes a health status indicator which is significantly higher than the state average.
 LNE (Low Number of Events): is used because the rate or percent may be unreliable.
¹Data for mothers who used tobacco are self-reported.
²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

○ ALL CAUSES	805.1
○ Heart Disease	198.3
Malignant Neoplasms (cancer)	197.2
Trachea, Bronchus, & Lung	44.1
Colon, Rectum, & Anus	28.3
Female Breast	25.7
Pancreas	7.5
Cerebrovascular Disease	44.5
● Chronic Lower Respiratory Diseases	30.0
Accidents	56.6
Motor Vehicle Accidents	30.4
● Alzheimer's Disease	19.6
Diabetes Mellitus	29.8
● Influenza & Pneumonia	7.8
Intentional Self-Harm (suicide)	22.5
Chronic Liver Disease and Cirrhosis	22.7
○ Infant Mortality	16.34

Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	147
2. Cancer	132
3. Cerebrovascular Disease	34
4. Accidents	28
5. Chronic Lower Respiratory Disease	22
6. Diabetes	19
7. Alzheimer's Disease	16
8. Chronic Liver Disease and Cirrhosis	11
9. Suicide	10
10. Unspecified dementia	9

Percent of deaths due to tobacco use 17.6
 Median age at death 80


● Denotes a health status indicator which is significantly lower than the state average.
 ○ Denotes a health status indicator which is significantly higher than the state average.
³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See Technical Notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Sanborn County

Demographic Information



Sanborn County is located in east central South Dakota and averages 4.1 persons per square mile.

2011 Population Information

Subject	Number	Percent
Total population	2,392	100.0
White	2,344	98.0
Black or African American	0	0.0
American Indian and Alaska Native	12	0.5
Asian	7	0.3
Native Hawaiian and Other Pacific Islander	0	0.0
Two or More Races	31	1.3
Under 5 years	134	5.6
Under 18 years	500	20.9
65 years and over	500	20.9

Source: United States Census Bureau, 2011 American Community Survey estimates

Health Status Indicators 2007-2011

Natality

Percent of Low Birth Weight Infants	6.5
Percent of Mothers Receiving Care in 1st Trimester	72.6
Percent of Mothers Who Used Tobacco While Pregnant ¹	13.9
Percent of Births Less Than 37 Weeks of Gestation	5.8
○ Average Age of Mother	28.5
Teenage Birth Rate ²	LNE
Percent White Births	99.3
Percent American Indian Births	0.0
● Percent Unmarried	23.2
● Percent WIC births	29.0
Percent Breastfeeding at discharge	78.3
○ Percent Payment-Private Insurance	75.4
Percent Payment-Medicaid	18.1
Percent C-Section	31.2

● Denotes a health status indicator which is significantly lower than the state average.
 ○ Denotes a health status indicator which is significantly higher than the state average.
 LNE (Low Number of Events): is used because the rate or percent may be unreliable.
¹Data for mothers who used tobacco are self-reported.
²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

ALL CAUSES	607.4
Heart Disease	122.5
Malignant Neoplasms (cancer)	115.4
● Trachea, Bronchus, & Lung	19.2
Colon, Rectum, & Anus	22.6
Female Breast	LNE
Pancreas	LNE
Cerebrovascular Disease	33.0
● Chronic Lower Respiratory Diseases	20.6
Accidents	82.7
Motor Vehicle Accidents	49.7
Alzheimer's Disease	50.6
Diabetes Mellitus	26.9
Influenza & Pneumonia	18.4
Intentional Self-Harm (suicide)	LNE
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	LNE

Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	24
2. Cancer	22
3. Accidents	12
4. Alzheimer's Disease	11
5. Cerebrovascular Disease	7
6. Diabetes	5

Percent of deaths due to tobacco use 17.1
 Median age at death 81


● Denotes a health status indicator which is significantly lower than the state average.
 ○ Denotes a health status indicator which is significantly higher than the state average.
³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See Technical Notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Shannon County

Demographic Information



Shannon County is located in the southwestern part of the state, along the Nebraska border and averages 6.5 persons per square mile.

2011 Population Information

Subject	Number	Percent
Total population	13,928	100.0
White	752	5.4
Black or African American	42	0.3
American Indian and Alaska Native	12,869	92.4
Asian	56	0.4
Native Hawaiian and Other Pacific Islander	14	0.1
Two or More Races	195	1.4
Under 5 years	1,588	11.4
Under 18 years	5,446	39.1
65 years and over	864	6.2

Source: United States Census Bureau, 2011 American Community Survey estimates

Health Status Indicators 2007-2011

Nativity

Percent of Low Birth Weight Infants	6.2
● Percent of Mothers Receiving Care in 1st Trimester	51.6
○ Percent of Mothers Who Used Tobacco While Pregnant ¹	24.0
Percent of Births Less Than 37 Weeks of Gestation	10.3
● Average Age of Mother	24.1
○ Teenage Birth Rate ²	73.3
Percent White Births	2.3
Percent American Indian Births	96.8
○ Percent Unmarried	85.6
○ Percent WIC births	75.2
● Percent Breastfeeding at discharge	49.9
● Percent Payment-Private Insurance	3.8
○ Percent Payment-Medicaid	68.1
● Percent C-Section	23.3

● Denotes a health status indicator which is significantly lower than the state average.
 ○ Denotes a health status indicator which is significantly higher than the state average.
 LNE (Low Number of Events): is used because the rate or percent may be unreliable.
¹Data for mothers who used tobacco are self-reported.
²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

○ ALL CAUSES	1471.1
○ Heart Disease	222.1
Malignant Neoplasms (cancer)	221.3
Trachea, Bronchus, & Lung	58.9
Colon, Rectum, & Anus	34.8
Female Breast	19.7
Pancreas	8.2
○ Cerebrovascular Disease	74.7
○ Chronic Lower Respiratory Diseases	91.4
○ Accidents	153.3
○ Motor Vehicle Accidents	77.1
Alzheimer's Disease	LNE
○ Diabetes Mellitus	124.2
○ Influenza & Pneumonia	52.8
○ Intentional Self-Harm (suicide)	28.2
○ Chronic Liver Disease and Cirrhosis	93.5
○ Infant Mortality	14.83

Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Accidents	90
T2. Heart Disease	72
T2. Cancer	72
4. Chronic Liver Disease and Cirrhosis	46
5. Diabetes	41
6. Chronic Lower Respiratory Disease	24
7. Suicide	23
8. Cerebrovascular Disease	20
9. Influenza and Pneumonia	18
10. Nephritis, nephrotic syndrome, and nephrosis	14

Percent of deaths due to tobacco use 11.6
 Median age at death 57


● Denotes a health status indicator which is significantly lower than the state average.
 ○ Denotes a health status indicator which is significantly higher than the state average.
³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See Technical Notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Spink County

Demographic Information



Spink County is located in the center of eastern South Dakota and averages 4.3 persons per square mile.

2011 Population Information

Subject	Number	Percent
Total population	6,470	100.0
White	6,302	97.4
Black or African American	19	0.3
American Indian and Alaska Native	91	1.4
Asian	6	0.1
Native Hawaiian and Other Pacific Islander	0	0.0
Two or More Races	45	0.7
Under 5 years	382	5.9
Under 18 years	1,598	24.7
65 years and over	1,275	19.7

Source: United States Census Bureau, 2011 American Community Survey estimates

Health Status Indicators 2007-2011

Nativity

○ Percent of Low Birth Weight Infants	10.8
Percent of Mothers Receiving Care in 1st Trimester	68.5
Percent of Mothers Who Used Tobacco While Pregnant ¹	16.1
Percent of Births Less Than 37 Weeks of Gestation	10.1
Average Age of Mother	27.4
● Teenage Birth Rate ²	5.6
Percent White Births	98.1
Percent American Indian Births	0.8
● Percent Unmarried	24.1
● Percent WIC births	32.3
Percent Breastfeeding at discharge	74.9
○ Percent Payment-Private Insurance	66.7
● Percent Payment-Medicaid	23.5
Percent C-Section	28.6

●Denotes a health status indicator which is significantly lower than the state average.
 ○Denotes a health status indicator which is significantly higher than the state average.
 LNE (Low Number of Events): is used because the rate or percent may be unreliable.
¹Data for mothers who used tobacco are self-reported.
²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

ALL CAUSES	736.4
Heart Disease	167.8
Malignant Neoplasms (cancer)	138.3
Trachea, Bronchus, & Lung	30.8
Colon, Rectum, & Anus	19.5
Female Breast	LNE
Pancreas	7.3
Cerebrovascular Disease	39.4
Chronic Lower Respiratory Diseases	60.4
○ Accidents	70.6
Motor Vehicle Accidents	20.5
Alzheimer's Disease	44.8
Diabetes Mellitus	26.8
Influenza & Pneumonia	17.5
Intentional Self-Harm (suicide)	9.5
Chronic Liver Disease and Cirrhosis	15.8
Infant Mortality	10.58

Leading Causes of Death

Total Deaths

1. Heart Disease	98
2. Cancer	74
3. Chronic Lower Respiratory Disease	32
4. Alzheimer's Disease	31
5. Accidents	28
6. Cerebrovascular Disease	23
T7. Diabetes	12
T7. Influenza and Pneumonia	12
9. Chronic Liver Disease and Cirrhosis	7
T10. Parkinson's Disease	5
T10. Pneumonitis due to solids and liquids	5
T10. Congenital Malformations, deformations, and chromosomal abnormalities	5

Percent of deaths due to tobacco use 14.4
 Median age at death 82


●Denotes a health status indicator which is significantly lower than the state average.
 ○Denotes a health status indicator which is significantly higher than the state average.
³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See Technical Notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Stanley County

Demographic Information



Stanley County is located in the center of the state and averages 2.1 persons per square mile.

2011 Population Information

Subject	Number	Percent
Total population	3,002	100.0
White	2,705	90.1
Black or African American	9	0.3
American Indian and Alaska Native	198	6.6
Asian	3	0.1
Native Hawaiian and Other Pacific Islander	0	0.0
Two or More Races	84	2.8
Under 5 years	183	6.1
Under 18 years	693	23.1
65 years and over	498	16.6

Source: United States Census Bureau, 2011 American Community Survey estimates

Health Status Indicators 2007-2011

Natality

Percent of Low Birth Weight Infants	7.7
Percent of Mothers Receiving Care in 1st Trimester	72.3
○ Percent of Mothers Who Used Tobacco While Pregnant ¹	26.0
Percent of Births Less Than 37 Weeks of Gestation	8.2
Average Age of Mother	27.3
Teenage Birth Rate ²	17.1
Percent White Births	80.6
Percent American Indian Births	14.8
Percent Unmarried	45.4
Percent WIC births	47.4
Percent Breastfeeding at discharge	75.0
Percent Payment-Private Insurance	61.2
Percent Payment-Medicaid	38.3
○ Percent C-Section	40.8

● Denotes a health status indicator which is significantly lower than the state average.
 ○ Denotes a health status indicator which is significantly higher than the state average.
 LNE (Low Number of Events): is used because the rate or percent may be unreliable.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

● ALL CAUSES	536.5
● Heart Disease	81.5
Malignant Neoplasms (cancer)	172.3
Trachea, Bronchus, & Lung	57.1
Colon, Rectum, & Anus	19.0
Female Breast	LNE
Pancreas	LNE
Cerebrovascular Disease	42.5
Chronic Lower Respiratory Diseases	53.0
Accidents	31.9
Motor Vehicle Accidents	LNE
Alzheimer's Disease	LNE
Diabetes Mellitus	25.5
Influenza & Pneumonia	LNE
Intentional Self-Harm (suicide)	LNE
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	LNE

Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Cancer	29
2. Heart Disease	14
3. Chronic Lower Respiratory Disease	10
4. Cerebrovascular Disease	6
5. Accidents	5

Percent of deaths due to tobacco use 36.0

Median age at death 69

● Denotes a health status indicator which is significantly lower than the state average.

○ Denotes a health status indicator which is significantly higher than the state average.


³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See Technical Notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Sully County

Demographic Information



Sully County is located in the central region of the state and averages 1.4 persons per square mile.

2011 Population Information

Subject	Number	Percent
Total population	1,375	100.0
White	1,330	96.7
Black or African American	3	0.2
American Indian and Alaska Native	17	1.2
Asian	0	0.0
Native Hawaiian and Other Pacific Islander	0	0.0
Two or More Races	26	1.9
Under 5 years	87	6.3
Under 18 years	298	21.7
65 years and over	254	18.5

Source: United States Census Bureau, 2011 American Community Survey estimates

Health Status Indicators 2007-2011

Natality

Percent of Low Birth Weight Infants	7.8
Percent of Mothers Receiving Care in 1st Trimester	73.0
Percent of Mothers Who Used Tobacco While Pregnant ¹	11.2
Percent of Births Less Than 37 Weeks of Gestation	8.9
Average Age of Mother	28.1
Teenage Birth Rate ²	LNE
Percent White Births	92.2
Percent American Indian Births	6.7
• Percent Unmarried	21.1
• Percent WIC births	25.6
Percent Breastfeeding at discharge	75.6
Percent Payment-Private Insurance	70.0
Percent Payment-Medicaid	25.6
Percent C-Section	31.1

•Denotes a health status indicator which is significantly lower than the state average.
 ○Denotes a health status indicator which is significantly higher than the state average.
 LNE (Low Number of Events): is used because the rate or percent may be unreliable.
¹Data for mothers who used tobacco are self-reported.
²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

• ALL CAUSES	487.9
Heart Disease	124.1
Malignant Neoplasms (cancer)	125.7
Trachea, Bronchus, & Lung	43.0
Colon, Rectum, & Anus	LNE
Female Breast	LNE
Pancreas	LNE
Cerebrovascular Disease	28.4
Chronic Lower Respiratory Diseases	LNE
Accidents	54.2
Motor Vehicle Accidents	42.6
Alzheimer's Disease	LNE
Diabetes Mellitus	28.4
Influenza & Pneumonia	LNE
Intentional Self-Harm (suicide)	LNE
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	LNE

Leading Causes of Death

Total Deaths

T1. Heart Disease	12
T1. Cancer	12
Percent of deaths due to tobacco use	27.3
Median age at death	75


•Denotes a health status indicator which is significantly lower than the state average.
 ○Denotes a health status indicator which is significantly higher than the state average.
³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See Technical Notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Todd County

Demographic Information



Todd County is located in south central South Dakota, along the Nebraska border and averages 6.9 persons per square mile.

2011 Population Information

Subject	Number	Percent
Total population	9,822	100.0
White	1,080	11.0
Black or African American	29	0.3
American Indian and Alaska Native	8,506	86.6
Asian	29	0.3
Native Hawaiian and Other Pacific Islander	0	0.0
Two or More Races	187	1.9
Under 5 years	1,306	13.3
Under 18 years	4,027	41.0
65 years and over	599	6.1

Source: United States Census Bureau, 2011 American Community Survey estimates

Health Status Indicators 2007-2011

Natality

Percent of Low Birth Weight Infants	8.0
● Percent of Mothers Receiving Care in 1st Trimester	39.1
○ Percent of Mothers Who Used Tobacco While Pregnant ¹	26.4
Percent of Births Less Than 37 Weeks of Gestation	9.8
● Average Age of Mother	24.4
○ Teenage Birth Rate ²	77.0
Percent White Births	3.9
Percent American Indian Births	93.6
○ Percent Unmarried	83.0
○ Percent WIC births	73.3
● Percent Breastfeeding at discharge	55.6
● Percent Payment-Private Insurance	4.0
○ Percent Payment-Medicaid	47.0
○ Percent C-Section	33.7

●Denotes a health status indicator which is significantly lower than the state average.
 ○Denotes a health status indicator which is significantly higher than the state average.
 LNE (Low Number of Events): is used because the rate or percent may be unreliable.
¹Data for mothers who used tobacco are self-reported.
²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

○ ALL CAUSES	1204.2
Heart Disease	182.9
Malignant Neoplasms (cancer)	208.8
Trachea, Bronchus, & Lung	68.3
Colon, Rectum, & Anus	17.6
Female Breast	LNE
Pancreas	LNE
Cerebrovascular Disease	59.3
Chronic Lower Respiratory Diseases	74.7
○ Accidents	137.3
○ Motor Vehicle Accidents	72.4
Alzheimer's Disease	18.5
○ Diabetes Mellitus	101.9
Influenza & Pneumonia	17.5
○ Intentional Self-Harm (suicide)	76.4
○ Chronic Liver Disease and Cirrhosis	50.2
○ Infant Mortality	14.98

Leading Causes of Death

Total Deaths

T1. Heart Disease	57
T1. Accidents	57
3. Cancer	54
4. Suicide	38
5. Diabetes	31
6. Chronic Lower Respiratory Disease	20
7. Chronic Liver Disease and Cirrhosis	17
8. Cerebrovascular Disease	15
9. Assault (homicide)	10
T10. Nephritis, nephrotic syndrome, and nephrosis	9
T10. Congenital malformations, deformations, and chromosomal abnormalities	9
Percent of deaths due to tobacco use	15.0
Median age at death	58


●Denotes a health status indicator which is significantly lower than the state average.
 ○Denotes a health status indicator which is significantly higher than the state average.
³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See Technical Notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Tripp County

Demographic Information



Tripp County is located along the Nebraska border in south central South Dakota and averages 3.5 persons per square mile.

2011 Population Information

Subject	Number	Percent
Total population	5,615	100.0
White	4,683	83.4
Black or African American	11	0.2
American Indian and Alaska Native	780	13.9
Asian	11	0.2
Native Hawaiian and Other Pacific Islander	0	0.0
Two or More Races	129	2.3
Under 5 years	331	5.9
Under 18 years	1,297	23.1
65 years and over	1,179	21.0

Source: United States Census Bureau, 2011 American Community Survey estimates

Health Status Indicators 2007-2011

Nativity

Percent of Low Birth Weight Infants	8.1
Percent of Mothers Receiving Care in 1st Trimester	65.1
○ Percent of Mothers Who Used Tobacco While Pregnant ¹	25.6
Percent of Births Less Than 37 Weeks of Gestation	8.4
Average Age of Mother	26.5
Teenage Birth Rate ²	21.3
Percent White Births	62.9
Percent American Indian Births	32.0
○ Percent Unmarried	48.5
○ Percent WIC births	53.9
Percent Breastfeeding at discharge	65.9
● Percent Payment-Private Insurance	40.7
○ Percent Payment-Medicaid	51.5
○ Percent C-Section	33.5

● Denotes a health status indicator which is significantly lower than the state average.

○ Denotes a health status indicator which is significantly higher than the state average.

LNE (Low Number of Events): is used because the rate or percent may be unreliable.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

ALL CAUSES	704.3
Heart Disease	176.3
Malignant Neoplasms (cancer)	135.7
● Trachea, Bronchus, & Lung	27.4
Colon, Rectum, & Anus	19.4
Female Breast	23.1
Pancreas	8.1
Cerebrovascular Disease	42.3
Chronic Lower Respiratory Diseases	46.5
Accidents	47.5
Motor Vehicle Accidents	27.0
Alzheimer's Disease	38.2
Diabetes Mellitus	32.0
Influenza & Pneumonia	22.3
Intentional Self-Harm (suicide)	11.5
Chronic Liver Disease and Cirrhosis	11.7
Infant Mortality	LNE

Leading Causes of Death

Total Deaths

1. Heart Disease	101
2. Cancer	63
3. Alzheimer's Disease	27
4. Chronic Lower Respiratory Disease	24
5. Cerebrovascular Disease	22
6. Diabetes	17
7. Accidents	15
8. Influenza and Pneumonia	13
9. Parkinson's Disease	6

Percent of deaths due to tobacco use

Median age at death

● Denotes a health status indicator which is significantly lower than the state average.

○ Denotes a health status indicator which is significantly higher than the state average.


³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See Technical Notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Turner County

Demographic Information



Turner County is located in southeastern South Dakota and averages 13.5 persons per square mile.

2011 Population Information

Subject	Number	Percent
Total population	8,332	100.0
White	8,124	97.5
Black or African American	25	0.3
American Indian and Alaska Native	92	1.1
Asian	17	0.2
Native Hawaiian and Other Pacific Islander	0	0.0
Two or More Races	75	0.9
Under 5 years	508	6.1
Under 18 years	1,966	23.6
65 years and over	1,566	18.8

Source: United States Census Bureau, 2011 American Community Survey estimates

Health Status Indicators 2007-2011

Nativity

Percent of Low Birth Weight Infants	6.8
Percent of Mothers Receiving Care in 1st Trimester	73.1
● Percent of Mothers Who Used Tobacco While Pregnant ¹	13.0
Percent of Births Less Than 37 Weeks of Gestation	10.0
○ Average Age of Mother	27.9
● Teenage Birth Rate ²	3.3
Percent White Births	98.7
Percent American Indian Births	0.6
● Percent Unmarried	23.7
● Percent WIC births	26.0
Percent Breastfeeding at discharge	76.3
○ Percent Payment-Private Insurance	72.9
● Percent Payment-Medicaid	21.1
Percent C-Section	28.6

● Denotes a health status indicator which is significantly lower than the state average.

○ Denotes a health status indicator which is significantly higher than the state average.

LNE (Low Number of Events): is used because the rate or percent may be unreliable.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

ALL CAUSES	673.6
Heart Disease	176.4
Malignant Neoplasms (cancer)	146.8
Trachea, Bronchus, & Lung	36.6
Colon, Rectum, & Anus	17.9
Female Breast	15.5
Pancreas	10.0
Cerebrovascular Disease	35.2
Chronic Lower Respiratory Diseases	41.6
Accidents	40.1
Motor Vehicle Accidents	14.9
Alzheimer's Disease	51.0
Diabetes Mellitus	21.9
Influenza & Pneumonia	16.5
Intentional Self-Harm (suicide)	8.3
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	LNE

Leading Causes of Death

Total Deaths

1. Heart Disease	134
2. Cancer	98
3. Alzheimer's Disease	45
4. Chronic Lower Respiratory Disease	30
5. Cerebrovascular Disease	26
6. Accidents	19
7. Diabetes	16
8. Influenza and Pneumonia	15
9. Unspecified Dementia	8
10. Nephritis, nephrotic syndrome, and nephrosis	7

Percent of deaths due to tobacco use 13.1
Median age at death 84

● Denotes a health status indicator which is significantly lower than the state average.

○ Denotes a health status indicator which is significantly higher than the state average.


³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See Technical Notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Union County

Demographic Information



Union County is located in the southeastern corner of the state and averages 31.3 persons per square mile.

2011 Population Information

Subject	Number	Percent
Total population	14,651	100.0
White	14,094	96.2
Black or African American	117	0.8
American Indian and Alaska Native	103	0.7
Asian	147	1.0
Native Hawaiian and Other Pacific Islander	15	0.1
Two or More Races	176	1.2
Under 5 years	952	6.5
Under 18 years	3,736	25.5
65 years and over	2,110	14.4

Source: United States Census Bureau, 2011 American Community Survey estimates

Health Status Indicators 2007-2011

Natality

Percent of Low Birth Weight Infants	8.1
○ Percent of Mothers Receiving Care in 1st Trimester	83.7
● Percent of Mothers Who Used Tobacco While Pregnant ¹	13.8
○ Percent of Births Less Than 37 Wks. of Gestation	11.7
○ Average Age of Mother	28.6
● Teenage Birth Rate ²	6.4
Percent White Births	96.2
Percent American Indian Births	0.6
● Percent Unmarried	21.2
● Percent WIC births	20.1
Percent Breastfeeding at discharge	77.2
○ Percent Payment-Private Insurance	76.0
● Percent Payment-Medicaid	20.6
○ Percent C-Section	31.6

● Denotes a health status indicator which is significantly lower than the state average.
 ○ Denotes a health status indicator which is significantly higher than the state average.
 LNE (Low Number of Events): is used because the rate or percent may be unreliable.
¹Data for mothers who used tobacco are self-reported.
²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

ALL CAUSES	646.8
Heart Disease	144.1
Malignant Neoplasms (cancer)	183.8
Trachea, Bronchus, & Lung	55.8
Colon, Rectum, & Anus	19.0
Female Breast	17.8
Pancreas	8.7
Cerebrovascular Disease	30.3
Chronic Lower Respiratory Diseases	50.4
Accidents	40.7
Motor Vehicle Accidents	10.3
Alzheimer's Disease	28.0
Diabetes Mellitus	22.7
Influenza & Pneumonia	11.1
● Intentional Self-Harm (suicide)	6.9
Chronic Liver Disease and Cirrhosis	6.4
Infant Mortality	LNE

Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Cancer	153
2. Heart Disease	125
3. Chronic Lower Respiratory Disease	42
4. Accidents	33
5. Cerebrovascular Disease	27
6. Alzheimer's Disease	25
7. Diabetes	18
8. Unspecified Dementia	11
9. Influenza and Pneumonia	9
10. Other interstitial pulmonary diseases with fibrosis	7

Percent of deaths due to tobacco use 17.9
 Median age at death 81


● Denotes a health status indicator which is significantly lower than the state average.
 ○ Denotes a health status indicator which is significantly higher than the state average.
³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See Technical Notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Walworth County

Demographic Information



Walworth County is located in north central South Dakota, near the North Dakota border and averages 7.7 persons per square mile.

2011 Population Information

Subject	Number	Percent
Total population	5,575	100.0
White	4,622	82.9
Black or African American	11	0.2
American Indian and Alaska Native	775	13.9
Asian	17	0.3
Native Hawaiian and Other Pacific Islander	0	0.0
Two or More Races	151	2.7
Under 5 years	351	6.3
Under 18 years	1,232	22.1
65 years and over	1,310	23.5

Source: United States Census Bureau, 2011 American Community Survey estimates

Health Status Indicators 2007-2011

Natality

- Percent of Low Birth Weight Infants3.8
- Percent of Mothers Receiving
Care in 1st Trimester64.4
- Percent of Mothers Who
Used Tobacco While Pregnant¹23.2
- Percent of Births Less Than 37 Weeks of Gestation7.3
- Average Age of Mother25.7
- Teenage Birth Rate²32.4
- Percent White Births61.7
- Percent American Indian Births35.1
- Percent Unmarried49.0
- Percent WIC births47.1
- Percent Breastfeeding at discharge63.4
- Percent Payment-Private Insurance38.2
- Percent Payment-Medicaid51.3
- Percent C-Section23.2

●Denotes a health status indicator which is significantly lower than the state average.
○Denotes a health status indicator which is significantly higher than the state average.
LNE (Low Number of Events): is used because the rate or percent may be unreliable.
¹Data for mothers who used tobacco are self-reported.
²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

- ALL CAUSES699.1
- Heart Disease119.6
- Malignant Neoplasms (cancer).....134.8
 - Trachea, Bronchus, & Lung33.7
 - Colon, Rectum, & Anus25.6
 - Female Breast.....9.4
 - Pancreas5.4
- Cerebrovascular Disease40.4
- Chronic Lower Respiratory Diseases49.5
- Accidents43.7
 - Motor Vehicle Accidents24.6
- Alzheimer's Disease58.7
- Diabetes Mellitus28.6
- Influenza & Pneumonia.....30.6
- Intentional Self-Harm (suicide)LNE
- Chronic Liver Disease and Cirrhosis18.0
- Infant Mortality9.55

Leading Causes of Death

Total Deaths

1. Heart Disease	70
2. Cancer	69
3. Alzheimer's Disease	38
4. Chronic Lower Respiratory Disease	26
5. Cerebrovascular Disease	23
6. Diabetes	14
7. Influenza and Pneumonia	13
8. Accidents	11
T9. Chronic Liver Disease and Cirrhosis	5
T9. Septicemia	5

Percent of deaths due to tobacco use20.5
Median age at death82


●Denotes a health status indicator which is significantly lower than the state average.
○Denotes a health status indicator which is significantly higher than the state average.
³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See Technical Notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Yankton County

Demographic Information



Yankton County is located in southeastern South Dakota on the Nebraska border and averages 43.1 persons per square mile.

2011 Population Information

Subject	Number	Percent
Total population	22,612	100.0
White	21,165	93.6
Black or African American	384	1.7
American Indian and Alaska Native	633	2.8
Asian	136	0.6
Native Hawaiian and Other Pacific Islander	0	0.0
Two or More Races	294	1.3
Under 5 years	1,289	5.7
Under 18 years	4,862	21.5
65 years and over	3,708	16.4

Source: United States Census Bureau, 2011 American Community Survey estimates

Health Status Indicators 2007-2011

Natality

Percent of Low Birth Weight Infants	6.1
○ Percent of Mothers Receiving Care in 1st Trimester	83.4
○ Percent of Mothers Who Used Tobacco While Pregnant ¹	22.2
Percent of Births Less Than 37 Weeks of Gestation	7.4
○ Average Age of Mother	27.2
● Teenage Birth Rate ²	10.2
Percent White Births	92.8
Percent American Indian Births	3.5
Percent Unmarried	36.7
Percent WIC births	38.3
Percent Breastfeeding at discharge	68.7
○ Percent Payment-Private Insurance	64.8
Percent Payment-Medicaid	32.3
Percent C-Section	26.2

● Denotes a health status indicator which is significantly lower than the state average.
 ○ Denotes a health status indicator which is significantly higher than the state average.
 LNE (Low Number of Events): is used because the rate or percent may be unreliable.
¹Data for mothers who used tobacco are self-reported.
²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

● ALL CAUSES	655.0
Heart Disease	155.0
Malignant Neoplasms (cancer)	145.1
Trachea, Bronchus, & Lung	48.8
Colon, Rectum, & Anus	12.4
Female Breast	14.1
Pancreas	7.8
Cerebrovascular Disease	34.4
Chronic Lower Respiratory Diseases	48.1
Accidents	39.4
Motor Vehicle Accidents	14.9
Alzheimer's Disease	35.8
Diabetes Mellitus	34.6
Influenza & Pneumonia	13.9
Intentional Self-Harm (suicide)	15.3
Chronic Liver Disease and Cirrhosis	7.2
Infant Mortality	6.91

Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	256
2. Cancer	209
3. Chronic Lower Respiratory Disease	70
4. Alzheimer's Disease	66
5. Cerebrovascular Disease	59
6. Diabetes	52
7. Accidents	51
8. Influenza and Pneumonia	26
9. Essential (primary) hypertension and hypertensive renal disease	19
10. Suicide	18

Percent of deaths due to tobacco use 18.7
 Median age at death 82


● Denotes a health status indicator which is significantly lower than the state average.
 ○ Denotes a health status indicator which is significantly higher than the state average.
³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See Technical Notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Ziebach County

Demographic Information



Ziebach County is located in north central South Dakota and averages 1.4 persons per square mile.

2011 Population Information

Subject	Number	Percent
Total population	2,852	100.0
White	673	23.6
Black or African American	9	0.3
American Indian and Alaska Native	2,085	73.1
Asian	9	0.3
Native Hawaiian and Other Pacific Islander	0	0.0
Two or More Races	77	2.7
Under 5 years	317	11.1
Under 18 years	1,104	38.7
65 years and over	197	6.9

Source: United States Census Bureau, 2011 American Community Survey estimates

Health Status Indicators 2007-2011

Nativity

Percent of Low Birth Weight Infants	6.7
● Percent of Mothers Receiving Care in 1st Trimester	46.0
Percent of Mothers Who Used Tobacco While Pregnant ¹	22.3
Percent of Births Less Than 37 Weeks of Gestation	9.9
● Average Age of Mother	25.2
○ Teenage Birth Rate ²	41.2
Percent White Births	14.8
Percent American Indian Births	80.3
○ Percent Unmarried	68.6
○ Percent WIC births	82.5
● Percent Breastfeeding at discharge	52.5
● Percent Payment-Private Insurance	13.9
○ Percent Payment-Medicaid	78.9
○ Percent C-Section	35.4

● Denotes a health status indicator which is significantly lower than the state average.

○ Denotes a health status indicator which is significantly higher than the state average.

LNE (Low Number of Events): is used because the rate or percent may be unreliable.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

○ ALL CAUSES	1028.7
Heart Disease	211.1
Malignant Neoplasms (cancer)	178.9
Trachea, Bronchus, & Lung	62.9
Colon, Rectum, & Anus	LNE
Female Breast	LNE
Pancreas	LNE
Cerebrovascular Disease	35.7
Chronic Lower Respiratory Diseases	63.4
○ Accidents	124.6
Motor Vehicle Accidents	53.3
Alzheimer's Disease	LNE
Diabetes Mellitus	88.0
Influenza & Pneumonia	LNE
Intentional Self-Harm (suicide)	LNE
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	17.94

Leading Causes of Death

Total Deaths

1. Heart Disease	16
2. Accidents	14
3. Cancer	13
T4. Diabetes	6
T4. Nephritis, nephrotic syndrome, and nephrosis	6
6. Chronic Lower Respiratory Disease	5

Percent of deaths due to tobacco use	15.3
Median age at death	72

● Denotes a health status indicator which is significantly lower than the state average.

○ Denotes a health status indicator which is significantly higher than the state average.

³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See Technical Notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics