

# Alcohol Use

## DRANK IN PAST 30 DAYS

**Definition:** Respondents who report drinking alcohol in the past 30 days.


### Prevalence of Drinking in Past 30 Days

- South Dakota 58%
- Nationwide median 55%

### Healthy People 2020 Objective


There was no comparable Healthy People 2020 Objective for drinking alcohol in the past 30 days.

**Figure 75**  
**Percent of Respondents Who Drank Alcohol in the Past 30 Days, 1995, 1997, 1999, and 2001-2010**



Source: The Behavioral Risk Factor Surveillance System, South Dakota Department of Health, 1995, 1997, 1999, and 2001-2010

**Figure 76**  
**Percent of Respondents Who Drank Alcohol in the Past 30 Days, 2011-2013**



Note: Beginning in 2011, the CDC began using a different methodology to weight the data, therefore current data estimates cannot be compared to previous years.

Source: The Behavioral Risk Factor Surveillance System, South Dakota Department of Health, 2011-2013

**Table 73**  
**Respondents Who Drank Alcohol in Past 30 Days, 2011-2013**

		2011-2013	2011	2012	2013
<b>Total</b>		<b>58%</b>	<b>59%</b>	<b>58%</b>	<b>58%</b>
<b>Gender</b>	Male	64%	64%	64%	65%
	Female	52%	54%	53%	51%
<b>Age</b>	18-34	64%	*	*	*
	35-44	66%	*	*	*
	45-54	64%	*	*	*
	55-64	58%	55%	61%	*
	65-74	47%	*	*	*
	75+	31%	*	*	*
<b>Race</b>	White	60%	60%	60%	60%
	American Indian	41%	*	*	*
<b>Ethnicity</b>	Hispanic	52%	*	*	*
	Non-Hispanic	58%	59%	58%	58%
<b>Household Income</b>	Less than \$15,000	44%	*	*	*
	\$15,000-\$24,999	47%	*	*	*
	\$25,000-\$34,999	55%	*	*	*
	\$35,000-\$49,999	59%	*	*	*
	\$50,000-\$74,999	69%	*	*	*
	\$75,000+	76%	76%	75%	*
<b>Education</b>	Less than High School, G.E.D.	37%	*	*	*
	High School, G.E.D.	53%	51%	54%	53%
	Some Post-High School	64%	66%	63%	62%
	College Graduate	68%	69%	68%	68%
<b>Employment Status</b>	Employed for Wages	66%	67%	65%	66%
	Self-employed	67%	*	*	*
	Unemployed	48%	*	*	*
	Homemaker	47%	*	*	*
	Student	58%	*	*	*
	Retired	43%	41%	45%	44%
	Unable to Work	30%	*	*	*
<b>Marital Status</b>	Married/Unmarried Couple	62%	63%	63%	61%
	Divorced/Separated	56%	*	*	*
	Widowed	32%	*	*	*
	Never Married	58%	*	*	*
<b>Home Ownership Status</b>	Own Home	60%	61%	60%	60%
	Rent Home	54%	55%	54%	*
<b>Children Status</b>	Children in Household (Ages 18-44)	64%	*	64%	*
	No Children in Household (Ages 18-44)	66%	*	*	*
<b>Phone Status</b>	Landline	55%	57%	55%	51%
	Cell Phone	65%	*	64%	65%
<b>Pregnancy Status</b>	Pregnant (Ages 18-44)	5%	*	*	*
	Not Pregnant (Ages 18-44)	62%	*	*	*


Note: \*Results based on small sample sizes have been suppressed.

Source: The Behavioral Risk Factor Surveillance System, South Dakota Department of Health, 2011-2013

### **National Statistics**

The national median for respondents who drank in the past 30 days was 54.5 percent, as shown in Figure 77 on the following page. South Dakota had 57.8 percent of respondents who drank in the past 30 days. Utah had the lowest percent of respondents who drank in the past 30 days with 30.7 percent, while Wisconsin had the highest with 64.6 percent.

**Figure 77**  
**Nationally, Respondents Who Drank Alcohol in Past 30 Days, 2013**



Source: The Behavioral Risk Factor Surveillance System, South Dakota Department of Health, 2013

**Further Analysis**

The following data show the percent of those who drank alcohol in the past 30 days for various health behaviors and conditions. For example, 52 percent of respondents who have high blood pressure drank alcohol in the past 30 days, while 61 percent of respondents who do not have high blood pressure drank alcohol in the past 30 days.

<b>Health Behavior or Condition</b>	<b>% Drank Alcohol in Past 30 Days</b>
Underweight (BMI < 18.5)	47%
Recommended (BMI = 18.5-24.9)	61%
Overweight (BMI = 25.0-29.9)	60%
Obese - Class I (BMI = 30.0-34.9)	59%
Obese - Classes II & III (BMI = 35.0+)	55%
Leisure Time Exercise	61%
No Leisure Time Exercise	50%
Met Physical Activity Recommendations	62%
Did Not Meet Physical Activity Recommendations	55%
2+ Servings of Fruit per Day	52%
< 2 Servings of Fruit per Day	61%
3+ Servings of Vegetables per Day	57%
< 3 Servings of Vegetables per Day	59%
5+ Servings of Fruits and Vegetables per Day	52%
< 5 Servings of Fruits and Vegetables per Day	59%

**Table 74 (continued)  
Drank Alcohol in Past 30 Days for Selected Health Behaviors and  
Conditions, 2011-2013**

<b>Health Behavior or Condition</b>	<b>% Drank Alcohol in Past 30 Days</b>
Current Smoker	63%
Former Smoker	59%
Never Smoked	56%
Smokeless Tobacco	69%
No Smokeless Tobacco	58%
Heard of "South Dakota QuitLine"	61%
Not Heard of "South Dakota QuitLine"	44%
Diabetes	38%
No Diabetes	60%
High Blood Pressure	52%
No High Blood Pressure	61%
High Blood Cholesterol	54%
No High Blood Cholesterol	61%
Health Insurance (18-64)	64%
No Health Insurance (18-64)	57%
Employer Based Coverage (18-64)	69%
Private Health Plan (18-64)	65%
Medicare (18-64)	32%
Medicaid or Medical Assistance (18-64)	33%
The Military, CHAMPUS, TriCare, or the VA (18-64)	64%
The Indian Health Service (18-64)	46%
Flu Shot (65+)	40%
No Flu Shot (65+)	38%
Pneumonia Shot (65+)	38%
No Pneumonia Shot (65+)	42%
Previously Had Heart Attack	39%
Never Had Heart Attack	59%
Have Angina or Coronary Heart Disease	43%
Do Not Have Angina or Coronary Heart Disease	59%
Previously Had a Stroke	38%
Never Had a Stroke	59%
Cancer	48%
No Cancer	60%
Skin Cancer	49%
No Skin Cancer	59%
Current Asthma	50%
Former Asthma	57%
Never Had Asthma	59%
Arthritis	48%
No Arthritis	62%
COPD	42%
No COPD	59%
Depression	53%
No Depression	59%
Kidney Disease	38%
No Kidney Disease	59%
Severe Vision Impairment	37%
No Severe Vision Impairment	58%
Recent Increased Confusion/Memory Loss	55%
No Recent Increased Confusion/Memory Loss	58%
< 6 Hours of Sleep	51%
6+ Hours of Sleep	59%

**Table 74 (continued)  
Drank Alcohol in Past 30 Days for Selected Health Behaviors and  
Conditions, 2011-2013**

<b>Health Behavior or Condition</b>	<b>% Drank Alcohol in Past 30 Days</b>
Always/Almost Always Wear Seat Belt	58%
Sometimes/Seldom/Never Wear Seat Belt	61%
Binge Drinker	100%
Not a Binge Drinker	47%
Heavy Drinker	100%
Not a Heavy Drinker	55%
Good/Very Good/Excellent Health Status	61%
Fair/Poor Health Status	38%
Physical Health Not Good for 0-29 Days of the Past 30	60%
Physical Health Not Good for 30 Days of the Past 30	34%
Mental Health Not Good for 0-19 Days of the Past 30	59%
Mental Health Not Good for 20-30 Days of the Past 30	52%
Usual Activities Unattainable for 0-9 Days of the Past 30	60%
Usual Activities Unattainable for 10-30 Days of the Past 30	42%
Physical, Mental, or Emotional Problems	44%
No Physical, Mental, or Emotional Problems	62%
Disability with Special Equipment Needed	35%
No Disability with Special Equipment Needed	60%
< 2 Hours of TV per Day	61%
2+ Hours of TV per Day	57%
Been Tested for HIV (18-64)	61%
Never Been Tested for HIV (18-64)	58%

Source: The Behavioral Risk Factor Surveillance System, South Dakota Department of Health, 2011-2013

## **BINGE DRINKING**

**Definition:** Male respondents who report having five or more alcoholic drinks on one occasion or female respondents who have four or more alcoholic drinks on one occasion, one or more times in the past month.


### **Prevalence of Binge Drinking**

- South Dakota 19%
- Nationwide median 17%

### **Healthy People 2020 Objective**


There was no comparable Healthy People 2020 Objective for binge drinking.

**Figure 78**  
**Percent of Respondents Who Engage in Binge Drinking, 1987-1993, 1995, 1997, 1999, and 2001-2010**



Source: The Behavioral Risk Factor Surveillance System, South Dakota Department of Health, 1987-1993, 1995, 1997, 1999, and 2001-2010

**Figure 79**  
**Percent of Respondents Who Engage in Binge Drinking, 2011-2013**



Note: Beginning in 2011, the CDC began using a different methodology to weight the data, therefore current data estimates cannot be compared to previous years.

Source: The Behavioral Risk Factor Surveillance System, South Dakota Department of Health, 2011-2013

**Table 75  
Respondents Who Engage in Binge Drinking, 2011-2013**

		2011-2013	2011	2012	2013
<b>Total</b>		<b>21%</b>	<b>22%</b>	<b>21%</b>	<b>19%</b>
<b>Gender</b>	Male	27%	29%	27%	25%
	Female	14%	16%	14%	14%
<b>Age</b>	18-34	35%	*	*	*
	35-44	25%	*	*	*
	45-54	20%	*	*	*
	55-64	11%	10%	12%	*
	65-74	5%	*	*	*
	75+	2%	*	*	*
<b>Race</b>	White	20%	22%	20%	18%
	American Indian	26%	*	*	*
<b>Ethnicity</b>	Hispanic	22%	*	*	*
	Non-Hispanic	21%	22%	20%	19%
<b>Household Income</b>	Less than \$15,000	22%	*	*	*
	\$15,000-\$24,999	21%	*	*	*
	\$25,000-\$34,999	21%	*	*	*
	\$35,000-\$49,999	19%	*	*	*
	\$50,000-\$74,999	23%	*	*	*
	\$75,000+	24%	27%	26%	*
<b>Education</b>	Less than High School, G.E.D.	15%	*	*	*
	High School, G.E.D.	20%	20%	19%	19%
	Some Post-High School	23%	26%	22%	21%
	College Graduate	21%	23%	22%	18%
<b>Employment Status</b>	Employed for Wages	26%	27%	25%	25%
	Self-employed	23%	*	*	*
	Unemployed	26%	*	*	*
	Homemaker	12%	*	*	*
	Student	32%	*	*	*
	Retired	4%	5%	6%	3%
	Unable to Work	10%	*	*	*
<b>Marital Status</b>	Married/Unmarried Couple	18%	20%	18%	17%
	Divorced/Separated	21%	*	*	*
	Widowed	4%	*	*	*
	Never Married	33%	*	*	*
<b>Home Ownership Status</b>	Own Home	18%	19%	18%	15%
	Rent Home	29%	30%	27%	*
<b>Children Status</b>	Children in Household (Ages 18-44)	29%	*	27%	*
	No Children in Household (Ages 18-44)	37%	*	*	*
<b>Phone Status</b>	Landline	15%	19%	13%	11%
	Cell Phone	31%	*	32%	28%
<b>Pregnancy Status</b>	Pregnant (Ages 18-44)	0.4%	*	*	*
	Not Pregnant (Ages 18-44)	25%	*	*	*


Note: \*Results based on small sample sizes have been suppressed.

Source: The Behavioral Risk Factor Surveillance System, South Dakota Department of Health, 2011-2013

### **National Statistics**

The national median for respondents who binge drink is 16.8 percent, as shown in Figure 80 on the next page. South Dakota had 19.2 percent of respondents who binge drink. Tennessee had the lowest percent of respondents who binge drink with 9.6 percent, while North Dakota had the highest with 23.2 percent.

**Figure 80**  
**Nationally, Respondents Who Engage in Binge Drinking, 2013**



Source: The Behavioral Risk Factor Surveillance System, South Dakota Department of Health, 2013

**Further Analysis**

The following data show the percent of those who engage in binge drinking for various health behaviors and conditions. For example, 42 percent of respondents who use smokeless tobacco also engage in binge drinking, while 19 percent of respondents who do not use smokeless tobacco engage in binge drinking.

<b>Health Behavior or Condition</b>	<b>% Binge Drinkers</b>
Underweight (BMI < 18.5)	13%
Recommended (BMI = 18.5-24.9)	23%
Overweight (BMI = 25.0-29.9)	22%
Obese - Class I (BMI = 30.0-34.9)	19%
Obese - Classes II & III (BMI = 35.0+)	19%
Leisure Time Exercise	21%
No Leisure Time Exercise	18%
Met Physical Activity Recommendations	20%
Did Not Meet Physical Activity Recommendations	21%
2+ Servings of Fruit per Day	12%
< 2 Servings of Fruit per Day	24%
3+ Servings of Vegetables per Day	14%
< 3 Servings of Vegetables per Day	21%

**Table 76 (continued)  
Binge Drinking for Selected Health Behaviors and Conditions,  
2011-2013**

<b>Health Behavior or Condition</b>	<b>% Binge Drinkers</b>
5+ Servings of Fruits and Vegetables per Day	12%
< 5 Servings of Fruits and Vegetables per Day	22%
Current Smoker	36%
Former Smoker	17%
Never Smoked	16%
Smokeless Tobacco	42%
No Smokeless Tobacco	19%
Heard of "South Dakota QuitLine"	21%
Not Heard of "South Dakota QuitLine"	11%
Diabetes	10%
No Diabetes	22%
High Blood Pressure	15%
No High Blood Pressure	23%
High Blood Cholesterol	13%
No High Blood Cholesterol	19%
Health Insurance (18-64)	23%
No Health Insurance (18-64)	28%
Employer Based Coverage (18-64)	24%
Private Health Plan (18-64)	22%
Medicare (18-64)	13%
Medicaid or Medical Assistance (18-64)	18%
The Military, CHAMPUS, TriCare, or the VA (18-64)	23%
The Indian Health Service (18-64)	33%
Flu Shot (65+)	3%
No Flu Shot (65+)	4%
Pneumonia Shot (65+)	2%
No Pneumonia Shot (65+)	5%
Previously Had Heart Attack	8%
Never Had Heart Attack	21%
Have Angina or Coronary Heart Disease	10%
Do Not Have Angina or Coronary Heart Disease	21%
Previously Had a Stroke	8%
Never Had a Stroke	21%
Cancer	10%
No Cancer	22%
Skin Cancer	8%
No Skin Cancer	21%
Current Asthma	17%
Former Asthma	24%
Never Had Asthma	21%
Arthritis	11%
No Arthritis	24%
COPD	14%
No COPD	21%
Depression	20%
No Depression	21%
Kidney Disease	9%
No Kidney Disease	21%
Severe Vision Impairment	12%
No Severe Vision Impairment	19%

**Table 76 (continued)  
Binge Drinking for Selected Health Behaviors and Conditions,  
2011-2013**

<b>Health Behavior or Condition</b>	<b>% Binge Drinkers</b>
Recent Increased Confusion/Memory Loss	19%
No Recent Increased Confusion/Memory Loss	19%
< 6 Hours of Sleep	22%
6+ Hours of Sleep	19%
Always/Almost Always Wear Seat Belt	18%
Sometimes/Seldom/Never Wear Seat Belt	32%
Drank Alcohol in Past 30 Days	36%
Did Not Drink Alcohol in Past 30 Days	0%
Heavy Drinker	85%
Not a Heavy Drinker	16%
Good/Very Good/Excellent Health Status	22%
Fair/Poor Health Status	13%
Physical Health Not Good for 0-29 Days of the Past 30	21%
Physical Health Not Good for 30 Days of the Past 30	11%
Mental Health Not Good for 0-19 Days of the Past 30	20%
Mental Health Not Good for 20-30 Days of the Past 30	25%
Usual Activities Unattainable for 0-9 Days of the Past 30	21%
Usual Activities Unattainable for 10-30 Days of the Past 30	17%
Physical, Mental, or Emotional Problems	12%
No Physical, Mental, or Emotional Problems	23%
Disability with Special Equipment Needed	8%
No Disability with Special Equipment Needed	22%
< 2 Hours of TV per Day	21%
2+ Hours of TV per Day	19%
Been Tested for HIV (18-64)	27%
Never Been Tested for HIV (18-64)	19%

Source: The Behavioral Risk Factor Surveillance System, South Dakota Department of Health, 2011-2013

## HEAVY DRINKING

**Definition:** Male respondents who report having more than 2 drinks per day, or female respondents who report having more than 1 drink per day.


### Prevalence of Heavy Drinking

- South Dakota 5%
- Nationwide median 6%

### Healthy People 2020 Objective


There was no comparable Healthy People 2020 Objective for heavy drinking.

**Figure 81**  
**Percent of Respondents Who Engage in Heavy Drinking, 1987-1993, 1995, 1997, 1999, and 2001-2010**



Source: The Behavioral Risk Factor Surveillance System, South Dakota Department of Health, 1987-1993, 1995, 1997, 1999, and 2001-2010

**Figure 82**  
**Percent of Respondents Who Engage in Heavy Drinking, 2011-2013**



Note: Beginning in 2011, the CDC began using a different methodology to weight the data, therefore current data estimates cannot be compared to previous years.

Source: The Behavioral Risk Factor Surveillance System, South Dakota Department of Health, 2011-2013

**Table 77**  
**Respondents Who Engage in Heavy Drinking, 2011-2013**

		2011-2013	2011	2012	2013
<b>Total</b>		<b>6%</b>	<b>6%</b>	<b>6%</b>	<b>5%</b>
<b>Gender</b>	Male	7%	7%	7%	6%
	Female	4%	5%	5%	4%
<b>Age</b>	18-34	8%	*	*	*
	35-44	6%	*	*	*
	45-54	7%	*	*	*
	55-64	4%	4%	6%	*
	65-74	4%	*	*	*
	75+	1%	*	*	*
<b>Race</b>	White	6%	6%	6%	5%
	American Indian	6%	*	*	*
<b>Ethnicity</b>	Hispanic	6%	*	*	*
	Non-Hispanic	6%	6%	6%	5%
<b>Household Income</b>	Less than \$15,000	7%	*	*	*
	\$15,000-\$24,999	6%	*	*	*
	\$25,000-\$34,999	7%	*	*	*
	\$35,000-\$49,999	5%	*	*	*
	\$50,000-\$74,999	5%	*	*	*
	\$75,000+	6%	7%	5%	*
<b>Education</b>	Less than High School, G.E.D.	4%	*	*	*
	High School, G.E.D.	6%	6%	6%	6%
	Some Post-High School	6%	7%	6%	6%
	College Graduate	5%	5%	6%	3%
<b>Employment Status</b>	Employed for Wages	6%	6%	6%	6%
	Self-employed	6%	*	*	*
	Unemployed	7%	*	*	*
	Homemaker	2%	*	*	*
	Student	10%	*	*	*
	Retired	3%	3%	5%	3%
	Unable to Work	5%	*	*	*
<b>Marital Status</b>	Married/Unmarried Couple	4%	5%	4%	4%
	Divorced/Separated	7%	*	*	*
	Widowed	2%	*	*	*
	Never Married	10%	*	*	*
<b>Home Ownership Status</b>	Own Home	5%	6%	5%	5%
	Rent Home	7%	6%	8%	*
<b>Children Status</b>	Children in Household (Ages 18-44)	6%	*	5%	*
	No Children in Household (Ages 18-44)	10%	*	*	*
<b>Phone Status</b>	Landline	4%	5%	4%	4%
	Cell Phone	9%	*	9%	7%
<b>Pregnancy Status</b>	Pregnant (Ages 18-44)	1%	*	*	*
	Not Pregnant (Ages 18-44)	6%	*	*	*


Note: \*Results based on small sample sizes have been suppressed.

Source: The Behavioral Risk Factor Surveillance System, South Dakota Department of Health, 2011-2013

### **National Statistics**

The national median for respondents who engage in heavy drinking is 6.2 percent, as shown in Figure 83 on the next page. South Dakota had 5.2 percent of respondents who engage in heavy drinking. Tennessee had the lowest percent of respondents who engage in heavy drinking with 3.3 percent, while Oregon had the highest with 8.6 percent.

**Figure 83**  
**Nationally, Respondents Who Engage in Heavy Drinking, 2013**



Source: The Behavioral Risk Factor Surveillance System, South Dakota Department of Health, 2013

**Further Analysis**

The following data show the percent of those who engage in heavy drinking for various health behaviors and conditions. For example, 12 percent of respondents who currently smoke are heavy drinkers, while three percent of respondents who have never smoked are heavy drinkers.

Health Behavior or Condition	% Heavy Drinkers
Underweight (BMI < 18.5)	5%
Recommended (BMI = 18.5-24.9)	7%
Overweight (BMI = 25.0-29.9)	6%
Obese - Class I (BMI = 30.0-34.9)	5%
Obese - Classes II & III (BMI = 35.0+)	5%
Leisure Time Exercise	5%
No Leisure Time Exercise	6%
Met Physical Activity Recommendations	5%
Did Not Meet Physical Activity Recommendations	6%
2+ Servings of Fruit per Day	4%
< 2 Servings of Fruit per Day	6%
3+ Servings of Vegetables per Day	4%
< 3 Servings of Vegetables per Day	6%
5+ Servings of Fruits and Vegetables per Day	3%
< 5 Servings of Fruits and Vegetables per Day	6%

**Table 78 (continued)  
Heavy Drinking for Selected Health Behaviors and Conditions,  
2011-2013**

<b>Health Behavior or Condition</b>	<b>% Heavy Drinkers</b>
Current Smoker	12%
Former Smoker	5%
Never Smoked	3%
Smokeless Tobacco	12%
No Smokeless Tobacco	5%
Heard of "South Dakota QuitLine"	5%
Not Heard of "South Dakota QuitLine"	4%
Diabetes	3%
No Diabetes	6%
High Blood Pressure	5%
No High Blood Pressure	6%
High Blood Cholesterol	4%
No High Blood Cholesterol	5%
Health Insurance (18-64)	6%
No Health Insurance (18-64)	8%
Employer Based Coverage (18-64)	5%
Private Health Plan (18-64)	7%
Medicare (18-64)	7%
Medicaid or Medical Assistance (18-64)	3%
The Military, CHAMPUS, TriCare, or the VA (18-64)	6%
The Indian Health Service (18-64)	9%
Flu Shot (65+)	2%
No Flu Shot (65+)	4%
Pneumonia Shot (65+)	2%
No Pneumonia Shot (65+)	3%
Previously Had Heart Attack	4%
Never Had Heart Attack	6%
Have Angina or Coronary Heart Disease	5%
Do Not Have Angina or Coronary Heart Disease	6%
Previously Had a Stroke	2%
Never Had a Stroke	6%
Cancer	4%
No Cancer	6%
Skin Cancer	4%
No Skin Cancer	6%
Current Asthma	7%
Former Asthma	7%
Never Had Asthma	6%
Arthritis	4%
No Arthritis	6%
COPD	7%
No COPD	6%
Depression	7%
No Depression	5%
Kidney Disease	4%
No Kidney Disease	6%
Severe Vision Impairment	9%
No Severe Vision Impairment	5%
Recent Increased Confusion/Memory Loss	8%
No Recent Increased Confusion/Memory Loss	5%

**Table 78 (continued)  
Heavy Drinking for Selected Health Behaviors and Conditions,  
2011-2013**

<b>Health Behavior or Condition</b>	<b>% Heavy Drinkers</b>
< 6 Hours of Sleep	8%
6+ Hours of Sleep	5%
Always/Almost Always Wear Seat Belt	5%
Sometimes/Seldom/Never Wear Seat Belt	10%
Drank Alcohol in Past 30 Days	10%
Did Not Drink Alcohol in Past 30 Days	0%
Binge Drinker	23%
Not a Binge Drinker	1%
Good/Very Good/Excellent Health Status	6%
Fair/Poor Health Status	5%
Physical Health Not Good for 0-29 Days of the Past 30	6%
Physical Health Not Good for 30 Days of the Past 30	5%
Mental Health Not Good for 0-19 Days of the Past 30	5%
Mental Health Not Good for 20-30 Days of the Past 30	10%
Usual Activities Unattainable for 0-9 Days of the Past 30	6%
Usual Activities Unattainable for 10-30 Days of the Past 30	7%
Physical, Mental, or Emotional Problems	5%
No Physical, Mental, or Emotional Problems	6%
Disability with Special Equipment Needed	4%
No Disability with Special Equipment Needed	6%
< 2 Hours of TV per Day	4%
2+ Hours of TV per Day	5%
Been Tested for HIV (18-64)	7%
Never Been Tested for HIV (18-64)	5%

Source: The Behavioral Risk Factor Surveillance System, South Dakota Department of Health, 2011-2013