


Map 1

Percent of Low Birth Weight Infants by County, 2007-2011

U.S. = 8.1% South Dakota = 6.5%




• Denotes that the county's percent is significantly lower than the state percent
 ○ Denotes that the county's percent is significantly higher than the state percent
 Note: "Significantly" refers to statistical significance at the 0.05 level. Although a county's actual percent may be higher or lower than the state percent, the difference may not be statistically significant due to the small number of people in the county. The percent of low birth weight infants is calculated based on the first weight of the newborn obtained after birth. Low birth weight infants are those born alive who weigh less than 2,500 grams (about 5 pounds 9 ounces). The U.S. percent of low birth weight infants is from 2010. See technical notes for more complete explanations.
 Source: South Dakota Department of Health, Office of Health Statistics.

Map 2

Percent of Mothers Receiving Prenatal Care in the 1st Trimester by County, 2007-2011

U.S. = 71.0%* **South Dakota = 69.4%**



- Denotes that the county's percent is significantly lower than the state percent.
- Denotes that the county's percent is significantly higher than the state percent.

Note: "Significantly" refers to statistical significance at the 0.05 level. Although a county's actual percent may be higher or lower than the state percent, the difference may not be statistically significant due to the small number of people in the county. The U.S. percent of mothers receiving prenatal care in the 1st trimester is based on 27 states that were using the 2003 standard birth certificate, including South Dakota. See technical notes for more complete explanations.


*Due to lack of comparable data from all 50 states, U.S. and South Dakota cannot be compared. The U.S. percent of first trimester prenatal care is from 2010.

Source: South Dakota Department of Health, Office of Health Statistics.

Map 3

Percent of Mothers Who Used Tobacco While Pregnant by County, 2007-2011

U.S. = 9.7%* **South Dakota = 18.1%**




• Denotes that the county's percent is significantly lower than the state percent.
 ○ Denotes that the county's percent is significantly higher than the state percent.
 Note: "Significantly" refers to statistical significance at the 0.05 level. Although a county's actual percent may be higher or lower than the state percent, the difference may not be statistically significant due to the small number of people in the county. The data for mothers who used tobacco while pregnant are self-reported on the birth certificates. The U.S. percent of mothers who used tobacco while pregnant is from 2008 and is based on 24 states using the 2003 standard birth certificate, including South Dakota. See technical notes for more complete explanations. *Due to lack of comparable data from all 50 states, U.S. and South Dakota cannot be compared.
 Source: South Dakota Department of Health, Office of Health Statistics.

Map 4

Percent of Births Less Than 37 Weeks Gestation by County, 2007-2011

U.S. = 12.0% **South Dakota = 8.5%**




● Denotes that the county's rate is significantly lower than the state rate.
 ○ Denotes that the county's rate is significantly higher than the state rate.
 Note: "Significantly" refers to statistical significance at the 0.05 level. Although a county's actual percent may be higher or lower than the state percent, the difference may not be statistically significant due to the small number of people in the county. The U.S. percent of births less than 37 weeks gestation is from 2010. See technical notes for more complete explanations.
 Source: South Dakota Department of Health, Office of Health Statistics.

Map 5

Average Age of Mother by Resident County, 2007-2011

U.S. = 27.7* **South Dakota = 27.0**




● Denotes that the county's rate is significantly lower than the state rate.
○ Denotes that the county's rate is significantly higher than the state rate.
Note: "Significantly" refers to statistical significance at the 0.05 level. Although a county's actual percent may be higher or lower than the state percent, the difference may not be statistically significant due to the small number of people in the county. The U.S. average age of mothers is from 2010. *Due to lack of comparable data from all 50 states, U.S. and South Dakota can not be compared. See technical notes for more complete explanations.
Source: South Dakota Department of Health, Office of Health Statistics.

Map 7

Death Rate Due to All Causes by County, 2007-2011

U.S. = 741.1 **South Dakota = 704.9**




• Denotes that the county's rate is significantly lower than the state rate.
 ○ Denotes that the county's rate is significantly higher than the state rate.
 Note: "Significantly" refers to statistical significance at the 0.05 level. Although a county's actual percent may be higher or lower than the state percent, the difference may not be statistically significant due to the small number of people in the county. The death rate is age-adjusted per 100,000 population. This eliminates age difference between populations, making them easier to compare. The U.S. age-adjusted death rate is from 2009. See technical notes for more complete explanations.
 Source: South Dakota Department of Health, Office of Health Statistics.

Map 9

Death Rate Due to All Malignant Neoplasms (Cancer) by County, 2007-2011

U.S. = 173.2 **South Dakota = 166.1**




• Denotes that the county's rate is significantly lower than the state rate.
 ○ Denotes that the county's rate is significantly higher than the state rate.
 Note: "Significantly" refers to statistical significance at the 0.05 level. Although a county's actual percent may be higher or lower than the state percent, the difference may not be statistically significant due to the small number of people in the county. The death rate is age-adjusted per 100,000 population. This eliminates age difference between populations, making them easier to compare. Malignant Neoplasms (Cancer) are defined as ICD-10 codes C00-C97. The U.S. age-adjusted Malignant Neoplasms (Cancer) death rate is from 2009. See technical notes for more complete explanations.
 Source: South Dakota Department of Health, Office of Health Statistics.

Map 11

Death Rate Due to Colorectal Cancer by County, 2007-2011

U.S. = 15.9 **South Dakota = 15.9**




Less than or equal to 14.5
 14.6-20.0
 Greater than or equal to 20.1
 Low Number Events

● Denotes that the county's rate is significantly lower than the state rate.
 ○ Denotes that the county's rate is significantly higher than the state rate.
 Note: "Significantly" refers to statistical significance at the 0.05 level. Although a county's actual percent may be higher or lower than the state percent, the difference may not be statistically significant due to the small number of people in the county. The death rate is age-adjusted per 100,000 population. This eliminates age difference between populations, making them easier to compare. Colorectal Cancer are defined as ICD-10 codes C18-C21. The U.S. age-adjusted Colorectal Cancer death rate is from 2008. See technical notes for more complete explanations.
 Source: South Dakota Department of Health, Office of Health Statistics.

Map 12

Death Rate Due to Female Breast Cancer by County, 2007-2011

U.S. = 22.3 **South Dakota = 21.0**




● Denotes that the county's rate is significantly lower than the state rate.
 ○ Denotes that the county's rate is significantly higher than the state rate.
 Note: "Significantly" refers to statistical significance at the 0.05 level. Although a county's actual percent may be higher or lower than the state percent, the difference may not be statistically significant due to the small number of people in the county. The death rate is age-adjusted per 100,000 population. This eliminates age difference between populations, making them easier to compare. Female Breast Cancer is defined as ICD-10 code C50. The U.S. age-adjusted Female Breast Cancer death rate is from 2009. See technical notes for more complete explanations.
 Source: South Dakota Department of Health, Office of Health Statistics.

Map 13

Death Rate Due to Pancreatic Cancer by County, 2007-2011

U.S. = 10.8 **South Dakota = 9.9**




● Denotes that the county's rate is significantly lower than the state rate.
 ○ Denotes that the county's rate is significantly higher than the state rate.
 Note: "Significantly" refers to statistical significance at the 0.05 level. Although a county's actual percent may be higher or lower than the state percent, the difference may not be statistically significant due to the small number of people in the county. The death rate is age-adjusted per 100,000 population. This eliminates age difference between populations, making them easier to compare. Pancreatic Cancer is defined as ICD-10 code C25. The U.S. age-adjusted Pancreatic Cancer death rate is from 2009. See technical notes for more complete explanations.
 Source: South Dakota Department of Health, Office of Health Statistics.

Map 14

Death Rate Due to Cerebrovascular Disease by County, 2007-2011

U.S. = 38.9 **South Dakota = 39.3**



● Denotes that the county's rate is significantly lower than the state rate.
 ○ Denotes that the county's rate is significantly higher than the state rate.
 Note: "Significantly" refers to statistical significance at the 0.05 level. Although a county's actual percent may be higher or lower than the state percent, the difference may not be statistically significant due to the small number of people in the county. The death rate is age-adjusted per 100,000 population. This eliminates age difference between populations, making them easier to compare. Cerebrovascular disease is defined as ICD-10 code I60-I69. The U.S. age-adjusted Cerebrovascular disease death rate is from 2009. See technical notes for more complete explanations.
 Source: South Dakota Department of Health, Office of Health Statistics.

