

The following table shows the percent of those who have previously had a heart attack for various health behaviors and conditions. For example, 21 percent of respondents who have been diagnosed with kidney disease have previously had a heart attack, while five percent of respondents who have not been diagnosed with kidney disease have previously had a heart attack.

**Previously Had a Heart Attack for Selected Health Behaviors and Conditions,  
2011-2014**

	<b>Previously Had Heart Attack</b>
Underweight (BMI < 18.5)	4%
Recommended Weight (BMI = 18.5-24.9)	3%
Overweight (BMI = 25.0-29.9)	6%
Obese - Class I (BMI = 30.0-34.9)	6%
Obese - Classes II & III (BMI = 35.0+)	6%
Participates in Leisure Time Exercise	4%
Does Not Participate in Leisure Time Exercise	7%
Watching Salt Intake	9%
Not Watching Salt Intake	2%
Current Smoker	5%
Former Smoker	9%
Never Smoked	3%
Uses Smokeless Tobacco	3%
Does Not Use Smokeless Tobacco	5%
Heard of "South Dakota Quitline"	5%
Not Heard of "South Dakota Quitline"	7%
Ever Diagnosed with Diabetes	16%
Never Diagnosed with Diabetes	4%
Has High Blood Pressure	11%
Does Not Have High Blood Pressure	2%
Has Health Insurance (18-64)	2%
Does Not Have Health Insurance (18-64)	2%
Employer Based Coverage (18-64)	2%
Private Health Plan (18-64)	1%
Medicare (18-64)	10%
Medicaid or Medical Assistance (18-64)	4%
The military, CHAMPUS, TriCare, or the VA (18-64)	5%
The Indian Health Service (18-64)	3%
Breast Exam Within Past Two Years	2%
No Breast Exam Within Past Two Years	5%
Mammogram Within Past Two Years (40+)	4%
No Mammogram Within Past Two Years (40+)	5%
Pap Smear Within Past Three Years	1%
No Pap Smear Within Past Three Years	4%
Blood Stool Test Within Past Two Years (50+)	10%
No Blood Stool Test Within Past Two Years (50+)	9%
Ever Had a Sigmoidoscopy or Colonoscopy (50+)	10%
Never Had a Sigmoidoscopy or Colonoscopy (50+)	7%
PSA Test Within Past Two Years (40+)	14%
No PSA Test Within Past Two Years (40+)	8%
Had a Flu Shot in Past 12 Months (65+)	16%
Did Not Have a Flu Shot in Past 12 Months (65+)	13%
Ever Had a Pneumonia Shot (65+)	18%
Never Had a Pneumonia Shot (65+)	10%
Ever Had a Shingles Shot (50+)	13%
Never Had a Shingles Shot (50+)	8%
Have Angina or Coronary Heart Disease	50%
Does Not Have Angina or Coronary Heart Disease	3%

**Previously Had a Heart Attack for Selected Health Behaviors and Conditions,  
2011-2014**

	<b>Previously Had Heart Attack</b>
Previously Had a Stroke	27%
Never Had a Stroke	4%
Ever Diagnosed with Cancer (Excluding Skin Cancer)	12%
Never Diagnosed with Cancer	4%
Ever Diagnosed with Skin Cancer	14%
Never Diagnosed with Skin Cancer	4%
Uses Sun Block Most of the Time	4%
Does Not Use Sun Block Most of the Time	5%
Current Asthma	8%
Former Asthma	6%
Never Had Asthma	5%
Ever Diagnosed with Arthritis	10%
Never Diagnosed with Arthritis	3%
Ever Diagnosed with COPD	17%
Never Diagnosed with COPD	4%
Ever Diagnosed with Depression	6%
Never Diagnosed with Depression	5%
Ever Diagnosed with Kidney Disease	21%
Never Diagnosed with Kidney Disease	5%
Has Severe Vision Impairment	18%
Does Not Have Severe Vision Impairment	5%
Recent Increased Confusion/Memory Loss	15%
No Recent Increased Confusion/Memory Loss	5%
< 6 hours of sleep	6%
6+ hours of sleep	5%
Been to the Dentist in the Past Year	4%
Hasn't Been to the Dentist in the Past Year	7%
Always/Almost Always Wears Seat Belt	5%
Sometimes/Seldom/Never Wears Seat Belt	6%
Drank Alcohol in Past 30 Days	3%
Did Not Drink Alcohol in Past 30 Days	7%
Binge Drinker	2%
Not a Binge Drinker	6%
Heavy Drinker	4%
Not a Heavy Drinker	5%
Good/Very Good/Excellent Health Status	3%
Fair/Poor Health Status	15%
Physical Health Not Good for 0-29 Days of the Past 30	4%
Physical Health Not Good for 30 Days of the Past 30	16%
Mental Health Not Good for 0-19 Days of the Past 30	5%
Mental Health Not Good for 20-30 Days of the Past 30	6%
Usual Activities Unattainable for 0-9 Days of the Past 30	4%
Usual Activities Unattainable for 10-30 Days of the Past 30	13%
Has Physical, Mental, or Emotional Problems	11%
Does Not Have Physical, Mental, or Emotional Problems	3%
Has a Disability with Special Equipment Needed	17%
Does Not Have a Disability with Special Equipment Needed	4%
Injured in a Fall in Past 12 Months (45+)	12%
Not Injured in a Fall in Past 12 Months (45+)	8%
Ever Been Tested for HIV	4%
Never Been Tested for HIV	5%

Source: South Dakota Behavioral Risk Factor Surveillance System, 2011-2014