

The following table shows the percent of those who have previously had a stroke for various health behaviors and conditions. For example, 11 percent of respondents who have been diagnosed with kidney disease have previously had a stroke, while three percent of respondents who have not been diagnosed with kidney disease have previously had a stroke.

**Previously Had a Stroke for Selected Health Behaviors and Conditions, 2011-2014**

|                                                   | <b>Previously Had a Stroke</b> |
|---------------------------------------------------|--------------------------------|
| Underweight (BMI < 18.5)                          | 3%                             |
| Recommended Weight (BMI = 18.5-24.9)              | 2%                             |
| Overweight (BMI = 25.0-29.9)                      | 3%                             |
| Obese - Class I (BMI = 30.0-34.9)                 | 3%                             |
| Obese - Classes II & III (BMI = 35.0+)            | 3%                             |
| Participates in Leisure Time Exercise             | 2%                             |
| Does Not Participate in Leisure Time Exercise     | 4%                             |
| Watching Salt Intake                              | 4%                             |
| Not Watching Salt Intake                          | 2%                             |
| Current Smoker                                    | 3%                             |
| Former Smoker                                     | 4%                             |
| Never Smoked                                      | 2%                             |
| Uses Smokeless Tobacco                            | 1%                             |
| Does Not Use Smokeless Tobacco                    | 3%                             |
| Heard of "South Dakota Quitline"                  | 2%                             |
| Not Heard of "South Dakota Quitline"              | 6%                             |
| Ever Diagnosed with Diabetes                      | 9%                             |
| Never Diagnosed with Diabetes                     | 2%                             |
| Has High Blood Pressure                           | 6%                             |
| Does Not Have High Blood Pressure                 | 1%                             |
| Has Health Insurance (18-64)                      | 2%                             |
| Does Not Have Health Insurance (18-64)            | 1%                             |
| Employer Based Coverage (18-64)                   | 1%                             |
| Private Health Plan (18-64)                       | 0%                             |
| Medicare (18-64)                                  | 10%                            |
| Medicaid or Medical Assistance (18-64)            | 5%                             |
| The military, CHAMPUS, TriCare, or the VA (18-64) | 2%                             |
| The Indian Health Service (18-64)                 | 2%                             |
| Breast Exam Within Past Two Years                 | 2%                             |
| No Breast Exam Within Past Two Years              | 4%                             |
| Mammogram Within Past Two Years (40+)             | 4%                             |
| No Mammogram Within Past Two Years (40+)          | 4%                             |
| Pap Smear Within Past Three Years                 | 1%                             |
| No Pap Smear Within Past Three Years              | 4%                             |
| Blood Stool Test Within Past Two Years (50+)      | 5%                             |
| No Blood Stool Test Within Past Two Years (50+)   | 5%                             |
| Ever Had a Sigmoidoscopy or Colonoscopy (50+)     | 5%                             |
| Never Had a Sigmoidoscopy or Colonoscopy (50+)    | 3%                             |
| PSA Test Within Past Two Years (40+)              | 5%                             |
| No PSA Test Within Past Two Years (40+)           | 3%                             |
| Had a Flu Shot in Past 12 Months (65+)            | 9%                             |
| Did Not Have a Flu Shot in Past 12 Months (65+)   | 6%                             |
| Ever Had a Pneumonia Shot (65+)                   | 9%                             |
| Never Had a Pneumonia Shot (65+)                  | 5%                             |
| Ever Had a Shingles Shot (50+)                    | 7%                             |
| Never Had a Shingles Shot (50+)                   | 4%                             |
| Previously Had Heart Attack                       | 15%                            |
| Never Had Heart Attack                            | 2%                             |
| Have Angina or Coronary Heart Disease             | 11%                            |
| Does Not Have Angina or Coronary Heart Disease    | 2%                             |

**Previously Had a Stroke for Selected Health Behaviors and Conditions, 2011-2014**

|                                                             | <b>Previously Had a Stroke</b> |
|-------------------------------------------------------------|--------------------------------|
| Ever Diagnosed with Cancer (Excluding Skin Cancer)          | 7%                             |
| Never Diagnosed with Cancer                                 | 2%                             |
| Ever Diagnosed with Skin Cancer                             | 7%                             |
| Never Diagnosed with Skin Cancer                            | 2%                             |
| Uses Sun Block Most of the Time                             | 2%                             |
| Does Not Use Sun Block Most of the Time                     | 3%                             |
| Current Asthma                                              | 5%                             |
| Former Asthma                                               | 4%                             |
| Never Had Asthma                                            | 2%                             |
| Ever Diagnosed with Arthritis                               | 6%                             |
| Never Diagnosed with Arthritis                              | 2%                             |
| Ever Diagnosed with COPD                                    | 10%                            |
| Never Diagnosed with COPD                                   | 2%                             |
| Ever Diagnosed with Depression                              | 5%                             |
| Never Diagnosed with Depression                             | 2%                             |
| Ever Diagnosed with Kidney Disease                          | 11%                            |
| Never Diagnosed with Kidney Disease                         | 3%                             |
| Has Severe Vision Impairment                                | 15%                            |
| Does Not Have Severe Vision Impairment                      | 2%                             |
| Recent Increased Confusion/Memory Loss                      | 9%                             |
| No Recent Increased Confusion/Memory Loss                   | 3%                             |
| < 6 hours of sleep                                          | 4%                             |
| 6+ hours of sleep                                           | 3%                             |
| Been to the Dentist in the Past Year                        | 2%                             |
| Hasn't Been to the Dentist in the Past Year                 | 4%                             |
| Always/Almost Always Wears Seat Belt                        | 3%                             |
| Sometimes/Seldom/Never Wears Seat Belt                      | 2%                             |
| Drank Alcohol in Past 30 Days                               | 2%                             |
| Did Not Drink Alcohol in Past 30 Days                       | 4%                             |
| Binge Drinker                                               | 1%                             |
| Not a Binge Drinker                                         | 3%                             |
| Heavy Drinker                                               | 1%                             |
| Not a Heavy Drinker                                         | 3%                             |
| Good/Very Good/Excellent Health Status                      | 2%                             |
| Fair/Poor Health Status                                     | 9%                             |
| Physical Health Not Good for 0-29 Days of the Past 30       | 2%                             |
| Physical Health Not Good for 30 Days of the Past 30         | 10%                            |
| Mental Health Not Good for 0-19 Days of the Past 30         | 3%                             |
| Mental Health Not Good for 20-30 Days of the Past 30        | 6%                             |
| Usual Activities Unattainable for 0-9 Days of the Past 30   | 2%                             |
| Usual Activities Unattainable for 10-30 Days of the Past 30 | 9%                             |
| Has Physical, Mental, or Emotional Problems                 | 7%                             |
| Does Not Have Physical, Mental, or Emotional Problems       | 2%                             |
| Has a Disability with Special Equipment Needed              | 11%                            |
| Does Not Have a Disability with Special Equipment Needed    | 2%                             |
| Injured in a Fall in Past 12 Months (45+)                   | 7%                             |
| Not Injured in a Fall in Past 12 Months (45+)               | 4%                             |
| Ever Been Tested for HIV                                    | 2%                             |
| Never Been Tested for HIV                                   | 3%                             |

Source: South Dakota Behavioral Risk Factor Surveillance System, 2011-2014