The following table shows the percent of those who have diabetes for various health behaviors and conditions. For example, 26 percent of respondents who have had a stroke have diabetes, while eight percent of respondents who have not had a stroke have diabetes

Prevalence of Diabetes for Selected Health Behaviors and Conditions, 2011-2015

	Ever Diagnosed with Diabetes
Underweight (BMI < 18.5)	4%
Recommended (BMI = 18.5-24.9)	3%
Overweight (BMI = 25.0-29.9)	8%
Obese - Class I (BMI = 30.0-34.9)	14%
Obese - Classes II & III (BMI = 35.0+)	22%
Participates in Leisure Time Exercise	8%
Does Not Participate in Leisure Time Exercise	13%
Meets Physical Activity Recommendations	8%
Does Not Meet Physical Activity Recommendations	10%
7+ Exercise Trips Per Week	7%
Less Than 7 Exercise Trips Per Week	10%
12+ Hours Sitting Per Day	15%
< 12 Hours Sitting Per Day	9%
Has 2+ Servings of Fruit per Day	11%
Has Less Than 2 Servings of Fruit per Day	9%
Has 3+ Servings of Vegetables per Day	9%
Has Less Than 3 Servings of Vegetables per Day	9%
Has 5+ Servings of Fruits and Vegetables per Day	10%
Has Less Than 5 Servings of Fruits and Vegetables per Day	9%
Current Smoker	8%
Former Smoker	13%
Never Smoked	7%
Uses Smokeless Tobacco	6%
Does Not Use Smokeless Tobacco	9%
Current E-Cigarette User	10%
Not E-Cigarette User	9%
Heard of "South Dakota Quitline"	9%
Not Heard of "South Dakota Quitline"	12%
Has High Blood Pressure	21%
Does Not Have High Blood Pressure	4%
Has High Blood Cholesterol	21%
Does Not Have High Blood Cholesterol	7%
Has Health Insurance (18-64)	7%
Does Not Have Health Insurance (18-64)	6%
Employer Based Coverage (18-64)	5%
Private Health Plan (18-64)	4%
Medicare (18-64)	21%
Medicaid or Medical Assistance (18-64)	7%
The military, CHAMPUS, TriCare, or the VA (18-64)	10%
The Indian Health Service (18-64)	14%
Had a Flu Shot in Past 12 Months (65+)	21%
Did Not Have a Flu Shot in Past 12 Months (65+)	15%
Ever Had a Pneumonia Shot (65+)	22%
Never Had a Pneumonia Shot (65+)	14%
Previously Had Heart Attack	30%
Never Had Heart Attack	8%

Prevalence of Diabetes for Selected Health Behaviors and Conditions, 2011-2015

	Ever Diagnosed with Diabetes
Knows Symptoms of Heart Attack	12%
Does Not Know Symptoms of Heart Attack	9%
Has Angina or Coronary Heart Disease	32%
Does Not Have Angina or Coronary Heart Disease	8%
Previously Had a Stroke	26%
Never Had a Stroke	8%
Ever Diagnosed with Cancer (Excluding Skin Cancer)	17%
Never Diagnosed with Cancer	8%
Ever Diagnosed with Skin Cancer	15%
Never Diagnosed with Skin Cancer	9%
Current Asthma	15%
Former Asthma	10%
Never Had Asthma	8%
Ever Diagnosed with Arthritis	18%
Never Diagnosed with Arthritis	6%
Ever Diagnosed with COPD	21%
Never Diagnosed with COPD	8%
Ever Diagnosed with Depression	13%
Never Diagnosed with Depression	8%
Ever Diagnosed with Kidney Disease	31%
Never Diagnosed with Kidney Disease	8%
Has Severe Vision Impairment	25%
Does Not Have Severe Vision Impairment	9%
Recent Increased Confusion/Memory Loss (45+)	22%
No Recent Increased Confusion/Memory Loss (45+)	14%
Always Wears Seat Belt	9%
Does Not Always Wear Seat Belt	9%
Drank Alcohol in Past 30 Days	6%
Did Not Drink Alcohol in Past 30 Days	13%
Binge Drinker	4%
Not a Binge Drinker	10%
Heavy Drinker	5%
Not a Heavy Drinker	9%
Has an Advance Directive	12%
Does Not Have an Advance Directive	8%
Good/Very Good/Excellent Health Status	7%
Fair/Poor Health Status	24%
Physical Health Not Good for 0-29 Days of the Past 30	8%
Physical Health Not Good for 30 Days of the Past 30	23%
Mental Health Not Good for 0-19 Days of the Past 30	9%
Mental Health Not Good for 20-30 Days of the Past 30	14%
Usual Activities Unattainable for 0-9 Days of the Past 30	8%
Usual Activities Unattainable for 10-30 Days of the Past 30	19%
Has Physical, Mental, or Emotional Problems	17%
Does Not Have Physical, Mental, or Emotional Problems	7%
Has a Disability with Special Equipment Needed	25%
Does Not Have a Disability with Special Equipment Needed	8%
Ever Been Tested for HIV	8%
Never Been Tested for HIV	9%

Source: South Dakota Behavioral Risk Factor Surveillance System, 2011-2015