


# Measles Vaccine Recommendations



SOUTH DAKOTA  
DEPARTMENT OF HEALTH

May 6, 2019



Note: There is no recommendation for an adult catch-up or vaccination campaign among adults or individuals in non-affected areas to prevent measles.

† Population groups recommended to receive 2 doses of MMR (or have presumptive evidence of immunity\*)

- International travelers
  - Adults and children aged 12 months or older: 2 doses of MMR
  - Children aged 6 to 11 months: 1 dose of MMR
- Healthcare personnel (even if born before 1957)
- Student at post-high school educational institutions
- People with HIV infection (who are not severely immunocompromised)
- Household or close contacts to immunocompromised persons

\* Presumptive evidence of immunity includes any of the following (documented rather than self-reported):

- 2 doses of MMR
- Laboratory evidence of immunity (positive measles IgG in serum)
- Laboratory confirmation of measles infection
- Born before 1957

‡ For the full recommendation visit:

<https://www.cdc.gov/vaccines/schedules/index.html>

Contraindications to giving a dose of MMR:

- Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to vaccine component
- Severe immunodeficiency (e.g., hematologic and solid tumors, receipt of chemotherapy, long-term immunosuppressive therapy, or patient with HIV infection who are severely immunocompromised)
- Currently pregnant
- History of anaphylactic reactions to neomycin