

REDUCING RISING SYPHILIS RATES

Doh.sd.gov/diseases/infectious/std/




SD PREVALENCE

874 Adult Syphilis Cases in 2021.

This represents a **1050%** increase from the 5-year median.

2021 Provisional Data, as of 03.01.2022



COURSE OF INFECTION

Infection
No symptoms
(1-90 days)




Primary Syphilis
Chancres at site of infection
(disappears after some weeks)



Secondary Syphilis
Disseminated rash
(after 1-6 months; lasts for 3 months)



Latent Syphilis
Serological proof of infection but no/minor signs and symptoms



Tertiary Syphilis
Benign gummas, cardiovascular, or neurosyphilis

RISK FACTORS

Pregnant Women 	MSM 	Multiple Sex Partners 	HIV Infected
Unstable Housing 	Incarceration 	IDU & Substance Abuse 	History of STI

Now Available 2021 Treatment Guidelines, with app available: <https://www.cdc.gov/std/treatment-guidelines/provider-resources.htm>
 CDC's Sexually Transmitted Infections Treatment Guidelines, 2021 provides current evidence-based prevention, diagnostic and treatment recommendations that replace the 2015 guidance. The recommendations are intended to be a source for clinical guidance.



This publication was supported by Centers for Disease Control and Prevention (CDC) grant number, 6 NH25PS005161-03-04 as a part of an award totaling \$1,000,000.00 with \$0.00 state funding. The views expressed in this written material or publication do not necessarily reflect the official policies of the Centers for Disease Control and Prevention (CDC), Department of Health and Human Services.