




SMOKING DURING PREGNANCY

South Dakota PRAMS, 2018


28% of South Dakota mothers smoked the two years before pregnancy and **10%** smoked the last three months of pregnancy.

97% of South Dakota mothers reported that their baby was not around anyone who was smoking.


Mothers who smoked the 3 months before pregnancy



Quit status among women who smoked the 3 months before getting pregnant



Number of cigarettes smoked the last 3 months of pregnancy among women who smoked in the last 2 years



WOMEN WHO WERE MORE LIKELY TO SMOKE DURING THE 3 MONTHS BEFORE PREGNANCY WERE:

- ▶ American Indian mothers
- ▶ Younger mothers
- ▶ Mothers with less years of education
- ▶ Unmarried mothers
- ▶ Mothers who had less household income

WOMEN WHO SMOKED THE LAST 3 MONTHS OF PREGNANCY WERE MORE LIKELY, COMPARED TO THOSE WHO DID NOT SMOKE, TO:

- ▶ Not be insured before pregnancy
- ▶ Use illicit drugs before pregnancy
- ▶ Attend less than 80% of prenatal care visits
- ▶ Not have their teeth cleaned during pregnancy
- ▶ Be emotionally abused by their husband or partner
- ▶ Have diabetes, hypertension or depression during pregnancy
- ▶ Never breastfeed
- ▶ High higher adverse childhood experiences

