
Map 1
Percent of Low Birth Weight Infants by County, 2014-2018
U.S. = 8.3%
South Dakota = 6.6%




X Denotes that the county's percent is significantly lower than the state percent

Note: "Significantly" refers to statistical significance at the 0.05 level. Although a county's actual percent may be higher or lower than the state percent, the difference may not be statistically significant due to the small number of people in the county. The percent of low birth weight infants is calculated based on the first weight of the newborn obtained after birth. Low birth weight infants are those born alive who weigh less than 2,500 grams (about 5 pounds 9 ounces). The U.S. percent of low birth weight infants is from 2018. See technical notes for more complete explanations.

Source: South Dakota Department of Health. Office of Health Statistics.

O Denotes that the county's percent is significantly higher than the state percent

Map 2
Percent of Mothers Receiving Prenatal Care in the 1st Trimester by County, 2014-2018
U.S. = 77.5%*
South Dakota = 75.1%




X Denotes that the county's percent is significantly lower than the state percent.

Note: "Significantly" refers to statistical significance at the 0.05 level. Although a county's actual percent may be higher or lower than the state percent, the difference may not be statistically significant due to the small number of people in the county. See technical notes for more complete explanations.

O Denotes that the county's percent is significantly higher than the state percent.

^{*}The U.S. percent of first trimester prenatal care is from 2018.

Map 3
Percent of Mothers Who Smoked Cigarettes While Pregnant by County, 2014-2018
U.S. = 6.5%*
South Dakota = 13.4%




X Denotes that the county's percent is significantly lower than the state percent.

Note: "Significantly" refers to statistical significance at the 0.05 level. Although a county's actual percent may be higher or lower than the state percent, the difference may not be statistically significant due to the small number of people in the county. Data for mothers who smoked cigarettes while pregnant are self-reported. See technical notes for more complete explanations. *The U.S. percent of cigarette smoking use by pregnant mothers is from 2018.

O Denotes that the county's percent is significantly higher than the state percent.

Map 4
Percent of Births Less Than 37 Weeks Gestation by County, 2014-2018
U.S. = 10.0% South Dakota = 8.9%




X Denotes that the county's rate is significantly lower than the state rate.

Note: "Significantly" refers to statistical significance at the 0.05 level. Although a county's actual percent may be higher or lower than the state percent, the difference may not be statistically significant due to the small number of people in the county. See technical notes for more complete explanations. The U.S. percent of births less than 37 weeks gestation is from 2018. Source: South Dakota Department of Health, Office of Health Statistics.

O Denotes that the county's rate is significantly higher than the state rate.

Map 5
Average Age of Mother by Resident County, 2014-2018
U.S. = 29.0
South Dakota = 28.1




X Denotes that the county's age is significantly lower than the state age.

Note: "Significantly" refers to statistical significance at the 0.05 level. Although a county's actual age may be higher or lower than the state age, the difference may not be statistically significant due to the small number of people in the county. The U.S. average age of mother is from 2018. See technical notes for more complete explanations.

Source: South Dakota Department of Health, Office of Health Statistics.

O Denotes that the county's age is significantly higher than the state age.

Map 6
Teenage Birth Rate by Resident County, 2014-2018
U.S. = 7.2
South Dakota = 10.8




X Denotes that the county's rate is significantly lower than the state rate.

O Denotes that the county's rate is significantly higher than the state rate.

Note: "Significantly" refers to statistical significance at the 0.05 level. Although a county's actual rate may be higher or lower than the state rate, the difference may not be statistically significant due to the small number of people in the county. The teenage birth rate is live births per 1,000 females age 15-17. The U.S. teenage birth rate is from 2018. See technical notes for more complete explanations.

Map 7
Death Rate Due to All Causes by County, 2014-2018
U.S. = 731.9
South Dakota = 719.1




X Denotes that the county's rate is significantly lower than the state rate.

O Denotes that the county's rate is significantly higher than the state rate.

Note: "Significantly" refers to statistical significance at the 0.05 level. Although a county's actual rate may be higher or lower than the state rate, the difference may not be statistically significant due to the small number of people in the county. The death rate is age-adjusted per 100,000 population. This eliminates age difference between populations, making them easier to compare. The U.S. age-adjusted death rate is from 2017. See technical notes for more complete explanations. Source: South Dakota Department of Health, Office of Health Statistics.

Map 8
Death Rate Due to Heart Disease by County, 2014-2018
U.S. = 165.0
South Dakota = 152.9




X Denotes that the county's rate is significantly lower than the state rate.

O Denotes that the county's rate is significantly higher than the state rate.

Note: "Significantly" refers to statistical significance at the 0.05 level. Although a county's actual rate may be higher or lower than the state rate, the difference may not be statistically significant due to the small number of people in the county. The death rate is age-adjusted per 100,000 population. This eliminates age difference between populations, making them easier to compare. Heart disease is defined as ICD-10 codes 100-109, 111, 113, and 120-151. The U.S. age-adjusted Heart Disease death rate is from 2017. See technical notes for more complete explanations. Source: South Dakota Department of Health. Office of Health Statistics.

Map 9
Death Rate Due to All Malignant Neoplasms (Cancer) by County, 2014-2018
U.S. = 152.5
South Dakota = 154.6




X Denotes that the county's rate is significantly lower than the state rate.

Note: "Significantly" refers to statistical significance at the 0.05 level. Although a county's actual rate may be higher or lower than the state rate, the difference may not be statistically significant due to the small number of people in the county. The death rate is age-adjusted per 100,000 population. This eliminates age difference between populations, making them easier to compare. Malignant Neoplasms (Cancer) are defined as ICD-10 codes C00-C97. The U.S. age-adjusted Malignant Neoplasms (Cancer) death rate is from 2017. See technical notes for more complete explanations. Source: South Dakota Department of Health, Office of Health Statistics.

O Denotes that the county's rate is significantly higher than the state rate.

Map 10
Death Rate Due to Trachea, Bronchus, and Lung Cancer by County, 2014-2018
U.S. = 36.6
South Dakota = 38.6




X Denotes that the county's rate is significantly lower than the state rate.

Note: "Significantly" refers to statistical significance at the 0.05 level. Although a county's actual rate may be higher or lower than the state rate, the difference may not be statistically significant due to the small number of people in the county. The death rate is age-adjusted per 100,000 population. This eliminates age difference between populations, making them easier to compare. Trachea, Bronchus, and Lung Cancer are defined as ICD-10 codes C33-C34. The U.S. age-adjusted Trachea, Bronchus, and Lung Cancer death rate is from 2017. See technical notes for more complete explanations.

O Denotes that the county's rate is significantly higher than the state rate.

Map 11
Death Rate Due to Colorectal Cancer by County, 2014-2018
U.S. = 13.7
South Dakota = 15.6




X Denotes that the county's rate is significantly lower than the state rate.

O Denotes that the county's rate is significantly higher than the state rate.

Note: "Significantly" refers to statistical significance at the 0.05 level. Although a county's actual rate may be higher or lower than the state rate, the difference may not be statistically significant due to the small number of people in the county. The death rate is age-adjusted per 100,000 population. This eliminates age difference between populations, making them easier to compare. Colorectal Cancer is defined as ICD-10 codes C18-C21. The U.S. age-adjusted Colorectal Cancer death rate is from 2017. See technical notes for more complete explanations.

Source: South Dakota Department of Health, Office of Health Statistics.

Map 12
Death Rate Due to Pancreatic Cancer by County, 2014-2018
U.S. = 11.1
South Dakota = 10.9




X Denotes that the county's rate is significantly lower than the state rate.

Note: "Significantly" refers to statistical significance at the 0.05 level. Although a county's actual rate may be higher or lower than the state rate, the difference may not be statistically significant due to the small number of people in the county. The death rate is age-adjusted per 100,000 population. This eliminates age difference between populations, making them easier to compare. Pancreatic Cancer is defined as ICD-10 code C25. The U.S. age-adjusted Pancreatic Cancer death rate is from 2017. See technical notes for more complete explanations.

Source: South Dakota Department of Health, Office of Health Statistics.

O Denotes that the county's rate is significantly higher than the state rate.

Map 13
Death Rate Due to Female Breast Cancer by County, 2014-2018
U.S. = 19.9
South Dakota = 18.6




X Denotes that the county's rate is significantly lower than the state rate.

Note: "Significantly" refers to statistical significance at the 0.05 level. Although a county's actual rate may be higher or lower than the state rate, the difference may not be statistically significant due to the small number of people in the county. The death rate is age-adjusted per 100,000 population. This eliminates age difference between populations, making them easier to compare. Female Breast Cancer is defined as ICD-10 code C50. The U.S. age-adjusted Female Breast Cancer death rate is from 2017. See technical notes for more complete explanations.

Source: South Dakota Department of Health, Office of Health Statistics.

O Denotes that the county's rate is significantly higher than the state rate.

Map 14
Death Rate Due to Chronic Lower Respiratory Diseases by County, 2014-2018
U.S. = 40.9
South Dakota = 42.7




X Denotes that the county's rate is significantly lower than the state rate.

Note: "Significantly" refers to statistical significance at the 0.05 level. Although a county's actual rate may be higher or lower than the state rate, the difference may not be statistically significant due to the small number of people in the county. The death rate is age-adjusted per 100,000 population. This eliminates age difference between populations, making them easier to compare. Chronic Lower Respiratory disease is defined as ICD-10 code J40-J47. The U.S. age-adjusted Chronic Lower Respiratory disease death rate is from 2017. See technical notes for more complete explanations. Source: South Dakota Department of Health, Office of Health Statistics.

O Denotes that the county's rate is significantly higher than the state rate.

Map 15
Death Rate Due to Accidents by County, 2014-2018
U.S. = 49.4
South Dakota = 50.9




X Denotes that the county's rate is significantly lower than the state rate.

Note: "Significantly" refers to statistical significance at the 0.05 level. Although a county's actual rate may be higher or lower than the state rate, the difference may not be statistically significant due to the small number of people in the county. The death rate is age-adjusted per 100,000 population. This eliminates age difference between populations, making them easier to compare. Accidents are defined as ICD-10 codes V01-X59, Y85-Y86. The U.S. age-adjusted Accident death rate is from 2017. See technical notes for more complete explanations.

Source: South Dakota Department of Health. Office of Health Statistics.

O Denotes that the county's rate is significantly higher than the state rate.

Map 16
Death Rate Due to Motor Vehicle Accidents by County, 2014-2018
U.S. = 12.0
South Dakota = 17.2




X Denotes that the county's rate is significantly lower than the state rate.

Note: "Significantly" refers to statistical significance at the 0.05 level. Although a county's actual rate may be higher or lower than the state rate, the difference may not be statistically significant due to the small number of people in the county. The death rate is age-adjusted per 100,000 population. This eliminates age difference between populations, making them easier to compare. Motor Vehicle Accidents are defined as ICD-10 codes (V02-V04, V09.0, V09.2, V12-V14, V19.0-V19.2, V19.4-V19.6, V20-V79, V80.3-V80.5, V81.0-V81.1, V82.0-V82.1, V83-V86, V87.0-V87.8, V88.0-V88.8, V89.0, and V89.2) The U.S. age-adjusted Motor Vehicle Accident death rate is from 2017. See technical notes for more complete explanations. Source: South Dakota Department of Health, Office of Health Statistics.

O Denotes that the county's rate is significantly higher than the state rate.

Map 17
Death Rate Due to Alzheimer's Disease by County, 2014-2018
U.S. = 31.0
South Dakota = 36.2




X Denotes that the county's rate is significantly lower than the state rate.

Note: "Significantly" refers to statistical significance at the 0.05 level. Although a county's actual rate may be higher or lower than the state rate, the difference may not be statistically significant due to the small number of people in the county. The death rate is age-adjusted per 100,000 population. This eliminates age difference between populations, making them easier to compare. Alzheimer's Disease is defined as ICD-10 code G30. The U.S. age-adjusted Alzheimer's Disease death rate is from 2017. See technical notes for more complete explanations. Source: South Dakota Department of Health, Office of Health Statistics.

O Denotes that the county's rate is significantly higher than the state rate.

Map 18
Death Rate Due to Cerebrovascular Disease by County, 2014-2018
U.S. = 37.6
South Dakota = 35.5




 $[\]ensuremath{\mathsf{X}}$ Denotes that the county's rate is significantly lower than the state rate.

Note: "Significantly" refers to statistical significance at the 0.05 level. Although a county's actual rate may be higher or lower than the state rate, the difference may not be statistically significant due to the small number of people in the county. The death rate is age-adjusted per 100,000 population. This eliminates age difference between populations, making them easier to compare. Cerebrovascular disease is defined as ICD-10 code I60-I69. The U.S. age-adjusted Cerebrovascular disease death rate is from 2017. See technical notes for more complete explanations. Source: South Dakota Department of Health, Office of Health Statistics.

O Denotes that the county's rate is significantly higher than the state rate.

Map 19
Death Rate Due to Diabetes Mellitus by County, 2014-2018
U.S. = 21.5
South Dakota = 23.9




X Denotes that the county's rate is significantly lower than the state rate.

Note: "Significantly" refers to statistical significance at the 0.05 level. Although a county's actual rate may be higher or lower than the state rate, the difference may not be statistically significant due to the small number of people in the county. The death rate is age-adjusted per 100,000 population. This eliminates age difference between populations, making them easier to compare. Diabetes Mellitus is defined as ICD-10 codes E10-E14. The U.S. age-adjusted Diabetes Mellitus death rate is from 2017. See technical notes for more complete explanations. Source: South Dakota Department of Health, Office of Health Statistics.

O Denotes that the county's rate is significantly higher than the state rate.

Map 20
Death Rate Due to Influenza and Pneumonia by County, 2014-2018
U.S. = 14.3
South Dakota = 18.2




X Denotes that the county's rate is significantly lower than the state rate.

Note: "Significantly" refers to statistical significance at the 0.05 level. Although a county's actual rate may be higher or lower than the state rate, the difference may not be statistically significant due to the small number of people in the county. The death rate is age-adjusted per 100,000 population. This eliminates age difference between populations, making them easier to compare. Influenza and Pneumonia are defined as ICD-10 codes J09-J18. The U.S. age-adjusted Influenza and Pneumonia death rate is from 2017. See technical notes for more complete explanations. Source: South Dakota Department of Health, Office of Health Statistics.

O Denotes that the county's rate is significantly higher than the state rate.

Map 21
Death Rate Due to Chronic Liver Disease and Cirrhosis by County, 2014-2018
U.S. = 10.9
South Dakota = 17.0




X Denotes that the county's rate is significantly lower than the state rate.

Note: "Significantly" refers to statistical significance at the 0.05 level. Although a county's actual rate may be higher or lower than the state rate, the difference may not be statistically significant due to the small number of people in the county. The death rate is age-adjusted per 100,000 population. This eliminates age difference between populations, making them easier to compare. Chronic Liver Disease and Cirrhosis are defined as ICD-10 codes K70, K73-K74. The U.S. age-adjusted Chronic Liver Disease and Cirrhosis death rate is from 2017. See technical notes for more complete explanations. Source: South Dakota Department of Health, Office of Health Statistics.

211

O Denotes that the county's rate is significantly higher than the state rate.

Map 22
Death Rate Due to Intentional Self-Harm (suicide) by County, 2014-2018
U.S. = 14.0
South Dakota = 19.9




X Denotes that the county's rate is significantly lower than the state rate.

Note: "Significantly" refers to statistical significance at the 0.05 level. Although a county's actual rate may be higher or lower than the state rate, the difference may not be statistically significant due to the small number of people in the county. The death rate is age-adjusted per 100,000 population. This eliminates age difference between populations, making them easier to compare. Intentional Self-Harm (suicide) is defined as ICD-10 codes *U03,X60-X84,Y87.0. The U.S. age-adjusted Intentional Self-Harm (suicide) death rate is from 2017. See technical notes for more complete explanations.

O Denotes that the county's rate is significantly higher than the state rate.

Map 23
Infant Mortality Rate by County, 2014-2018
U.S. = 5.79
South Dakota = 6.34



X Denotes that the county's rate is significantly lower than the state rate.

Note: "Significantly" refers to statistical significance at the 0.05 level. Although a county's actual rate may be higher or lower than the state rate, the difference may not be statistically significant due to the small number of people in the county. Infant mortality is calculated as the number of deaths to babies (less than 1 year old) per 1,000 live births. The U.S. infant mortality rate is from 2017. See technical notes for more complete explanations. Source: South Dakota Department of Health, Office of Health Statistics.

O Denotes that the county's rate is significantly higher than the state rate.