



## South Dakota Immunization Program Vaccines Available for VFC\* Eligible Children

Vaccine	Trade Name	Min Order	Presentation <sup>+</sup>	Ages/Indication
DT	Varies	1	Varies	6 weeks – 6 years Only to be administered when DTaP is medically contraindicated – call for approval to order.
DTaP	Daptacel <sup>®</sup>	10	SDV	6 weeks – 6 years
	Infanrix <sup>®</sup>	10	SDS	
DTaP-IPV	Kinrix <sup>®</sup>	10	SDS	4 years – 6 years
	Quadracel <sup>™</sup>	10	SDS	Approved for use as the fifth dose of DTaP and fourth dose of IPV.
DTaP-Hep B-IPV	Pediarix <sup>®</sup>	10	SDS	6 weeks – 6 years Approved only for the primary series (i.e., 2, 4, and 6 months).
DTaP-IPV-Hib	Pentacel <sup>®</sup>	5	SDV	6 weeks – 4 years
DTaP-IPV-Hib-HepB	Vaxelis <sup>™</sup>	10	SDS	6 weeks – 4 years
e-IPV	IPOL <sup>®</sup>	10	MDV	6 weeks – 18 years
Hepatitis A	Vaqta <sup>®</sup>	10	SDS	1 – 18 years
	Havrix <sup>®</sup>	10	SDS	Should be routinely administered to children 12 – 23 months of age.
Hepatitis A-B 18 only	Twinrix <sup>®</sup>	10	SDS	18 years of age only – call for approval to order.
Hepatitis B	Engerix B <sup>®</sup>	10	SDS	Birth – 18 years
	Recombivax HB <sup>®</sup>	10	SDS	
Hib	PedvaxHIB <sup>®</sup>	10	SDV	6 weeks – 59 months
	ActHIB <sup>®</sup>	5	SDV	PedvaxHIB <sup>®</sup> is a three-dose series at 2, 4, and 12 – 15 months. ActHIB <sup>®</sup> is a four-dose series at 2, 4, 6, and 12 – 15 months.
	Hiberix <sup>®</sup>	10	SDV	Hiberix <sup>®</sup> is a four-dose series at 2, 4, 6, and 12 – 15 months.
HPV	Gardasil <sup>®</sup> 9	10	SDS	Gardasil <sup>®</sup> 9 is available for both males and females ages 9 and older. Adolescents aged 9-14 years who begin vaccination before their 15 <sup>th</sup> birthday will only need two doses of HPV vaccine at least 6 months apart. Persons who start the series at age 15 or older or immunocompromised need 3 doses.
Meningococcal Conjugate	MenQuadfi <sup>™</sup>	5	SDV	2 months – 18 years Should be routinely administered to children 11-12 years of age, with a booster dose at 16 years of age.
	Menveo <sup>®</sup>	5	SDV	Minimum age for Menveo <sup>®</sup> is 2 months. Minimum age for MedQuadfi <sup>™</sup> is 2 years.
Meningococcal B	TRUMENBA <sup>®</sup>	10	SDS	16 – 18 years. High risk 10-18 years.
	BEXSERO <sup>®</sup>	1		TRUMENBA <sup>®</sup> is 2 doses at 0, and 6 months. During an outbreak or for high risk persons give 3 doses at 0, 1-2, and 6 months. BEXSERO <sup>®</sup> is 2 doses 1 month apart.
Measles, Mumps and Rubella	M-M-R <sup>®</sup> II Priorix	10	SDV	12 months – 18 years
MMR/Varicella	ProQuad <sup>®</sup>	10	SDV	12 months – 12 years For the first dose of MMR and varicella vaccines at ages 12 – 47 months, either MMR and varicella vaccines administered separately or MMRV vaccine may be used. For the second dose of MMR and varicella vaccines at any age (15 months--12 years) and for the first dose at age ≥48 months.
Pneumococcal Conjugate	Prevnar 13 <sup>™</sup> (PCV13)	10	SDS	6 weeks – 59 months The ACIP recommends PCV13 for all children ages two through 59 months and for children ages 60 through 71 months who have underlying medical conditions that increase their risk of pneumococcal disease or complications.
	Vaxneuvance <sup>™</sup> (PCV15)			
Pneumococcal Polysaccharide	Pneumovax <sup>®</sup> 23	1	SDS	2 – 18 years • Available for high-risk children with one or more of the following conditions: <b>1.</b> Chronic illnesses such as cardiovascular disease, chronic pulmonary disease, diabetes mellitus, alcoholism, chronic liver disease or CSF leaks. <b>2.</b> Functional or anatomic asplenia (splenectomy) <b>3.</b> Conditions associated with immunosuppression, including HIV infection, leukemia, lymphoma, Hodgkin's disease, multiple myeloma, generalized malignancy, chronic renal failure, nephritic syndrome, or other conditions associated with immunosuppression (organ or bone marrow transplantation); and people receiving immunosuppressive chemotherapy, including long term high-dose corticosteroids. <b>4.</b> Cochlear implants
Rotavirus, Live, Oral	RotaTeq <sup>®</sup>	10	SDT	6 weeks – 8 months, 0 days
	Rotarix <sup>®</sup>	10	SDT	Rotateq <sup>®</sup> (three dose rotavirus vaccine to be administered at 2, 4, and 6 months of age) Rotarix <sup>®</sup> (two dose rotavirus vaccine to be administered at 2 and 4 months of age)
Tetanus & Diphtheria Toxoids	TDVAX <sup>™</sup>	1	SDV	7 – 18 years • Tdap is recommended instead of Td. • Td is only necessary for children and adolescents who have not completed the primary series of DTaP, or who are contraindicated for Tdap.
Tdap	Boostrix <sup>®</sup>	10	SDV	7 – 18 years
	Adacel <sup>®</sup>	10	SDV	
Varicella	Varivax <sup>®</sup>	10	SDV	12 months – 18 years
Influenza	Varies	10	Varies	For VFC eligible children 6 months – 18 years of age.

\*VFC-eligible children are those who are 18 and younger and meet one of the following criteria (child does not need to be a resident of, or live in SD):

- No health insurance
- Medicaid eligible
- American Indian or Alaskan Native
- Underinsured – have health insurance, but it does not cover a particular vaccine (VFC vaccine can only be administered to underinsured children at Rural Health Clinics, Federally Qualified Health Centers, SD Department of Health Community Health Offices)

\*Presentations (this can change without notice):

SDS – Single Dose Syringe      SDV – Single Dose Vial      SDT – Single Dose Tube      MDV – Multi-dose vial