
United States

Demographic Information		Health Sta	atus I	ndicators		
2017 Population I	2017 Population Information		Natality – 2016		Mortality⁴ - 2016	
Subject	Number	Percent	Percent of Low Birth Weight Infants	8.2	All Causes	728.8
•			Percent of Mothers Receiving		Heart Disease	165.5
Total population White	325,719,178 249,619,493	100 76.6	Care in 1st Trimester	77.1	Malignant Neoplasms (Cancer)	155.8
Black or African American	43.499.874	13.4	Percent of Mothers Who	7.0	Trachea, Bronchus, & Lung	38.4
American Indian & Alaska Native	4,104,295	1.3	Used Tobacco While Pregnant ² Percent of Births Less Than 37 Wks. of Gestation	7.2 9.8	Colon, Rectum, & Anus Female Breast	13.9 20.1
Asian	18,989,059	5.8	Average Age of Mother	28.7	Pancreas	11.1
Native Hawaiian & Other Pacific			Teenage Birth Rate ³	8.8	Prostate	19.3
Islander	785,471	0.2	Percent White Births (2015)	75.7	Leukemia	6.2
Two or More Races	8,720,986	2.7	Percent American Indian Births (2015)	1.1	Chronic Lower Respiratory Diseases	40.6
			Percent Unmarried	39.8	Alzheimer's Disease	30.3
Under 5 years	19.938.860	6.1	Percent WIC births	39.6	Cerebrovascular Disease	37.3
Under 18 years	73,655,378	22.6	Percent Breastfeeding at discharge	83.1	Accidents	47.4
65 years and over	50,858,679	15.6	Percent Payment-Private Insurance	49.4	Motor Vehicle Accidents	12.1
,	,,-		Percent Payment-Medicaid	42.6	Diabetes	21.0
			Percent C-Section	31.9	Influenza and Pneumonia	13.5
					Intentional Self-Harm (Suicide)	13.5
					Chronic Liver Disease and Cirrhosis	10.7
					Infant Mortality	5.87
						Total
					Leading Causes of Death	Deaths
					1. Heart Disease	635,260
					Malignant Neoplasms (Cancer)	598,038
					3. Accidents	161,374
					4. Chronic Lower Respiratory Disease	154,596
					5. Cerebrovascular Disease	142,142
					6. Alzheimer's Disease	116,103
					7. Diabetes	80,058
					Influenza and Pneumonia	51,537
					Nephritis, Nephrotic Syndrome, and	50,046
					Nephrosis	•
					10. Intentional Self-Harm (Suicide)	44,965
					4-1	114
					⁴ The mortality rates, except infant mortal	ality are age
					adjusted death rates per 100,000 pop	
					adjusting to the standard million populati	
			Only one year of U.S. data are given to compare with five			
			of state and county data because the numbers on the r		ontrio.	
Course Haitad Ctate - Course Du	0047 Damidatia		level are much greater and do not fluctuate as much annua	ally.	Source: National Center for Health Stati	stics. Centers
Source: United States Census Bureau, 2 Estimates	2017 Population		² Data for mothers who used tobacco are self-reported. ³ Teenage birth rate is live births per 1,000 females age 15-	17	for Disease Control and Prevention, U.S. I	,
Louinaleo			reenage birtii rate is live birtiis per 1,000 remaies age 15-	17.	Health and Human Services, Hyattsville, Ma	

South Dakota

Demographic Information

Health Status Indicators 2013-2017



South Dakota is located in the north central portion of the United States and averages 10.7 persons per square mile.

2017 Population Information

Subject	Number	Percent
Total population	869,666	100.0
White	738,554	84.9
Black or African American	18,479	2.1
American Indian & Alaska Native	78,456	9.0
Asian	12,901	1.5
Native Hawaiian & Other Pacific Islander	719	0.1
Two or More Races	20,557	2.4
Under 5 years	61,759	7.1
Under 18 years	214,856	24.7
65 years and over	141,624	16.3

Source: United States Census Bureau, 2017 Population

Estimates

Natality	
 Percent of Low Birth Weight Infants 	6.5
Percent of Mothers Receiving	
Care in 1st Trimester	72.4
 Percent of Mothers Who 	
Used Tobacco While Pregnant ¹	14.0
 Percent of Births Less Than 37 Wks. of Gestation 	8.7
Average Age of Mother	27.9
○ Teenage Birth Rate ²	11.5
Percent White Births	74.5
Percent American Indian Births	15.6
 Percent Unmarried 	37.4
 Percent WIC births 	33.1
 Percent Breastfeeding at discharge 	79.4
 Percent Payment-Private Insurance 	58.5
Percent Payment-Medicaid	32.0
Percent C-Section	25.2

Mortality ³	
All Causes	711.5
Heart Disease	151.4
Malignant Neoplasms (Cancer)	156.6
Trachea, Bronchus, & Lung	39.8
 Colon, Rectum, & Anus 	15.9
Female Breast	18.5
Pancreas	11.0
Prostate	18.4
Leukemia	7.0
Chronic Lower Respiratory Diseases	41.8
 Alzheimer's Disease 	36.0
Cerebrovascular Disease	36.3
 Accidents 	50.9
 Motor Vehicle Accidents 	17.2
 Diabetes 	23.8
 Influenza and Pneumonia 	17.3
 Intentional Self-Harm (Suicide) 	19.7
 Chronic Liver Disease and Cirrhosis 	15.7
Infant Mortality	6.47

Leading Causes of Death	iotai	
Leading Causes of Death	Deaths	
1. Heart Disease	8,464	
Malignant Neoplasms (Cancer)	8,293	
3. Accidents	2,392	
4. Chronic Lower Respiratory Diseases	2,285	
5. Alzheimer's Disease	2,167	
Cerebrovascular Disease	2,064	
7. Diabetes	1,259	
8. Influenza and Pneumonia	991	
Intentional Self-Harm (Suicide)	814	
10. Chronic Liver Disease and Cirrhosis	696	
Percent of Deaths due to tobacco use	17.1	
Median age at death	79	

Total

See technical notes for more information. Source: South Dakota Department of Health, Office of **Health Statistics**

[•]Denotes a health status indicator which is significantly lower than the national average.

oDenotes a health status indicator which is significantly higher than the national average.

¹Data for mothers who used tobacco are self-reported.

²Teenage birth rate is live births per 1,000 females age 15-17.

[•]Denotes a health status indicator which is significantly lower than the national average.

oDenotes a health status indicator which is significantly higher than the national average.

³All mortality rates except infant mortality are ageadjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

Aurora County

Demographic Information

Aurora County is located in the south central portion of the state and averages 3.8 persons per square mile.

2017 Population Information

Subject	Number	Percent
Total population	2,738	100.0
White	2,595	94.8
Black or African American	22	0.8
American Indian & Alaska Native	72	2.6
Asian	20	0.7
Native Hawaiian & Other Pacific Islander	0	0.0
Two or More Races	29	1.1
Under 5 years	168	6.1
Under 18 years	690	25.2
65 years and over	581	21.2

Health Status Indicators 2013-2017

Natality	
Percent of Low Birth Weight Infants	8.0
Percent of Mothers Receiving	
Care in 1st Trimester	72.0
Percent of Mothers Who	
Used Tobacco While Pregnant ¹	10.6
Percent of Births Less Than 37 Wks. of Gestation	10.0
Average Age of Mother	28.1
Teenage Birth Rate ²	9.4
Percent White Births	99.0
Percent American Indian Births	LNE
 Percent Unmarried 	28.0
Percent WIC births	38.5
Percent Breastfeeding at discharge	74.5
 Percent Payment-Private Insurance 	74.4
Percent Payment-Medicaid	16.6
Percent C-Section	30.0

Notality

Mortality ³	
All Causes	600.7
Heart Disease	150.6
Malignant Neoplasms (Cancer)	126.8
Trachea, Bronchus, & Lung	35.0
Colon, Rectum, & Anus	LNE
Female Breast	43.2
Pancreas	LNE
Prostate	38.3
Leukemia	LNE
Chronic Lower Respiratory Diseases	21.5
Alzheimer's Disease	42.5
Cerebrovascular Disease	62.1
Accidents	51.4
Motor Vehicle Accidents	33.1
Diabetes	20.1
Influenza and Pneumonia	21.2

INF

LNE LNE

Intentional Self-Harm (Suicide)

Infant Mortality

Chronic Liver Disease and Cirrhosis

Leading Causes of Death	Total Deaths
1. Heart Disease	35
Malignant Neoplasms (Cancer)	28
Cerebrovascular Disease	14
4. Alzheimer's Disease	11
5. Accidents	8
Influenza and Pneumonia	6
T7. Chronic Lower Respiratory Diseases	5
T7. Diabetes	5
Percent of Deaths due to tobacco use	14.1
Median age at death	83

See technical notes for more information.

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

³All mortality rates except infant mortality are ageadjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

Beadle County

Demographic Information

Beadle County is located in the center of eastern South Dakota and averages 13.8 persons per square mile.

2017 Population Information

Subject	Number	Percent
Total population	18,157	100.0
White	15,561	85.7
Black or African American	232	1.3
American Indian & Alaska Native	259	1.4
Asian	1,702	9.4
Native Hawaiian & Other Pacific Islander	48	0.3
Two or More Races	355	2.0
Under 5 years	1,666	9.2
Under 18 years	4,990	27.5
65 years and over	3,155	17.4

Denotes a health status indicator which is significantly lower than the state average.
Denotes a health status indicator which is significantly higher

 Denotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Health Status Indicators 2013-2017

Natality	
Percent of Low Birth Weight Infants	6.8
Percent of Mothers Receiving	
Care in 1st Trimester	56.8
Percent of Mothers Who	
Used Tobacco While Pregnant ¹	12.8
Percent of Births Less Than 37 Wks. of Gestation	8.8
 Average Age of Mother 	26.8
 Teenage Birth Rate² 	22.4
Percent White Births	75.1
Percent American Indian Births	1.4
Percent Unmarried	39.0
 Percent WIC births 	48.5
Percent Breastfeeding at discharge	77.2
Percent Payment-Private Insurance	50.3
Percent Payment-Medicaid	39.2
 Percent C-Section 	31.2

Mortality ³	
All Causes	727.0
Heart Disease	134.0
Malignant Neoplasms (Cancer)	173.3
Trachea, Bronchus, & Lung	41.3
Colon, Rectum, & Anus	10.7
Female Breast	18.0
Pancreas	13.2
Prostate	19.8
Leukemia	5.8
Chronic Lower Respiratory Diseases	32.1
Alzheimer's Disease	44.6
Cerebrovascular Disease	46.5
Accidents	47.7
Motor Vehicle Accidents	13.9
Diabetes	34.6
 Influenza and Pneumonia 	35.5
Intentional Self-Harm (Suicide)	18.5
Chronic Liver Disease and Cirrhosis	15.4

Leading Causes of Death	Total Deaths
Malignant Neoplasms (Cancer)	214
2. Heart Disease	187
3. Alzheimer's Disease	73
Cerebrovascular Disease	69
Influenza and Pneumonia	52
6. Accidents	51
7. Diabetes	46
8. Chronic Lower Respiratory Diseases	42
Intentional Self-Harm (Suicide)	15
10. Chronic Liver Disease and Cirrhosis	14
Percent of Deaths due to tobacco use	19.8
Median age at death	82

Infant Mortality

7.41

See technical notes for more information.

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

³All mortality rates except infant mortality are ageadjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

Bennett County

Demographic Information

Bennett County is located on the southern border in western South Dakota and averages 2.9 persons per square mile.

2017 Population Information

Subject	Number	Percent
Total population	3,454	100.0
White	1,204	34.9
Black or African American	19	0.6
American Indian & Alaska Native	2,038	59.0
Asian	15	0.4
Native Hawaiian & Other Pacific Islander	1	0.0
Two or More Races	177	5.1
Under 5 years	328	9.5
Under 18 years	1,151	33.3
65 years and over	431	12.5

Health Status Indicators 2013-2017

Natality	
Percent of Low Birth Weight Infants	5.6
 Percent of Mothers Receiving 	
Care in 1st Trimester	56.1
 Percent of Mothers Who 	
Used Tobacco While Pregnant ¹	25.5
Percent of Births Less Than 37 Wks. of Gestation	9.5
 Average Age of Mother 	25.5
 Teenage Birth Rate² 	36.7
Percent White Births	15.1
Percent American Indian Births	77.8
 Percent Unmarried 	74.3
Percent WIC births	65.6
Percent Breastfeeding at discharge	51.2
Percent Payment-Private Insurance	15.2
Percent Payment-Medicaid Percent C. Castian	56.3
Percent C-Section	21.3

Mortality ³	
o All Causes	1,072.6
Heart Disease	199.0
Malignant Neoplasms (Cancer)	204.8
Trachea, Bronchus, & Lung	28.3
Colon, Rectum, & Anus	24.1
Female Breast	LNE
Pancreas	LNE
Prostate	34.9
Leukemia	LNE
Chronic Lower Respiratory Diseases	63.9
Alzheimer's Disease	17.7
Cerebrovascular Disease	22.8
 Accidents 	115.1
Motor Vehicle Accidents	41.7
o Diabetes	93.1
Influenza and Pneumonia	26.5
Intentional Self-Harm (Suicide)	30.7
Chronic Liver Disease and Cirrhosis	43.2

Leading Causes of Death	Total Deaths
Malignant Neoplasms (Cancer)	35
2. Heart Disease 3. Accidents	34 19
4. Diabetes	15
Chronic Lower Respiratory Diseases	11
Chronic Liver Disease and Cirrhosis	7
7. Septicemia	6
T8. Influenza and Pneumonia	5
T8. Intentional Self-Harm (Suicide)	5
Percent of Deaths due to tobacco use	22.0
Median age at death	72

23.67

Infant Mortality

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Source: United States Census Bureau, 2017 Population

Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

³All mortality rates except infant mortality are ageadjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

Bon Homme County

Demographic Information

Bon Homme County is located on the Nebraska border in eastern South Dakota and averages 12.5 persons per square mile.

2017 Population Information

Subject	Number	Percent
Total population White	6,984 6.218	100.0 89.0
Black or African American	84	1.2
American Indian & Alaska Native	566	8.1
Asian	12	0.2
Native Hawaiian & Other Pacific Islander	1	0.0
Two or More Races	103	1.5
Under 5 years	374	5.4
Under 18 years	1,367	19.6
65 years and over	1,432	20.5

Health Status Indicators 2013-2017

Natality	
Percent of Low Birth Weight Infants	5.0
Percent of Mothers Receiving	
Care in 1st Trimester	78.3
Percent of Mothers Who	
Used Tobacco While Pregnant ¹	10.9
Percent of Births Less Than 37 Wks. of Gestation	7.5
Average Age of Mother	28.4
Teenage Birth Rate ²	LNE
Percent White Births	96.0
Percent American Indian Births	1.9
 Percent Unmarried 	21.2
Percent WIC births	30.9
Percent Breastfeeding at discharge	82.2
 Percent Payment-Private Insurance 	74.7
 Percent Payment-Medicaid 	19.7
Percent C-Section	26.8

Notality

Mortality ³		
All Causes	613.1	
 Heart Disease 	111.1	
Malignant Neoplasms (Cancer)	143.1	
 Trachea, Bronchus, & Lung 	22.9	
Colon, Rectum, & Anus	14.6	
Female Breast	32.4	
Pancreas	13.7	
Prostate	12.6	
Leukemia	14.1	
Chronic Lower Respiratory Diseases	44.0	
 Alzheimer's Disease 	22.1	
Cerebrovascular Disease	41.1	
Accidents	43.2	
Motor Vehicle Accidents	15.6	
Diabetes	22.9	
Influenza and Pneumonia	22.1	
Intentional Self-Harm (Suicide)	11.6	
Chronic Liver Disease and Cirrhosis	LNE	
Infant Mortality	9.35	

Leading Causes of Death	iotai
Leading Causes of Death	Deaths
1. Malignant Neoplasms (Cancer	r) 84
Heart Disease	76
Cerebrovascular Disease	29
Chronic Lower Respiratory Dis	seases 27
Accidents	19
Influenza and Pneumonia	18
Alzheimer's Disease	16
 Essential (Primary) Hypertensi Hypertensive Renal Disease 	ion and 15
9. Diabetes	14
10. Unspecified Dementia	12
Percent of Deaths due to tobacco Median age at death	o use 11.1 85

See technical notes for more information.

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

³All mortality rates except infant mortality are ageadjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

Brookings County

Demographic Information

Brookings County is located on the eastern border of the state and averages 40.3 persons per square mile.

2017 Population Information

Subject	Number	Percent
Total population	34,255	100.0
White	31,883	93.1
Black or African American	535	1.6
American Indian & Alaska Native	397	1.2
Asian	884	2.6
Native Hawaiian & Other Pacific Islander	28	0.1
Two or More Races	528	1.5
Under 5 years	2,226	6.5
Under 18 years	7,013	20.5
65 years and over	4,000	11.7

Health Status Indicators 2013-2017

Natality	
Percent of Low Birth Weight Infants	5.5
 Percent of Mothers Receiving 	
Care in 1st Trimester	80.6
 Percent of Mothers Who 	
Used Tobacco While Pregnant ¹	9.1
 Percent of Births Less Than 37 Wks. of Gestation 	6.9
 Average Age of Mother 	28.7
 Teenage Birth Rate² 	4.8
Percent White Births	86.2
Percent American Indian Births	2.1
 Percent Unmarried 	21.6
 Percent WIC births 	21.2
 Percent Breastfeeding at discharge 	86.9
 Percent Payment-Private Insurance 	76.3
 Percent Payment-Medicaid 	16.9
Percent C-Section	20.4

Mortality ³	
All Causes	622.6
Heart Disease	145.5
Malignant Neoplasms (Cancer)	147.7
Trachea, Bronchus, & Lung	31.2
Colon, Rectum, & Anus	17.2
Female Breast	14.4
Pancreas	10.8
Prostate	17.1
Leukemia	7.9
 Chronic Lower Respiratory Diseases 	27.8
 Alzheimer's Disease 	25.8
Cerebrovascular Disease	45.7
 Accidents 	36.6
 Motor Vehicle Accidents 	8.7
Diabetes	22.7
 Influenza and Pneumonia 	10.7
Intentional Self-Harm (Suicide)	14.0
 Chronic Liver Disease and Cirrhosis 	8.6
Infant Mortality	6.45
	Total

Leading Causes of Death	ıotai
Leading Causes of Death	Deaths
1. Heart Disease	223
Malignant Neoplasms (Cancer)	217
Cerebrovascular Disease	71
4. Accidents	58
T5. Chronic Lower Respiratory Diseases	42
T5. Alzheimer's Disease	42
7. Diabetes	34
Intentional Self-Harm (Suicide)	21
Influenza and Pneumonia	17
Essential (Primary) Hypertension and	14
Hypertensive Renal Disease	
Percent of Deaths due to tobacco use	14.6
Median age at death	82

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Source: United States Census Bureau, 2017 Population

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

³All mortality rates except infant mortality are ageadjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

Brown County

Demographic Information

Brown County is located on the North Dakota border in the eastern part of the state and averages 21.3 persons per square mile.

2017 Population Information

Subject	Number	Percent
Total population	39,178	100.0
White	34,890	89.1
Black or African American	862	2.2
American Indian & Alaska Native	1,381	3.5
Asian	1,148	2.9
Native Hawaiian & Other Pacific Islander	80	0.2
Two or More Races	817	2.1
Under 5 years	2,635	6.7
Under 18 years	9,469	24.2
65 years and over	6,522	16.6

Source: United States Census Bureau, 2017 Population

Estimates

Health Status Indicators 2013-2017

Natality			
Percent of Low Birth Weight Infants	6.5		
Percent of Mothers Receiving			
Care in 1st Trimester	69.8		
 Percent of Mothers Who 			
Used Tobacco While Pregnant ¹	16.0		
Percent of Births Less Than 37 Wks. of Gestation	7.7		
Average Age of Mother	28.0		
 Teenage Birth Rate² 	6.3		
Percent White Births	84.4		
Percent American Indian Births	5.3		
 Percent Unmarried 	33.0		
 Percent WIC births 	29.3		
Percent Breastfeeding at discharge	82.3		
 Percent Payment-Private Insurance 	67.1		
Percent Payment-Medicaid	28.2		
Percent C-Section	27.6		

Mortality ³			
All Causes	695.2		
Heart Disease	139.8		
Malignant Neoplasms (Cancer)	165.3		
Trachea, Bronchus, & Lung	46.8		
Colon, Rectum, & Anus	18.0		
Female Breast	25.2		
Pancreas	10.8		
Prostate	17.3		
Leukemia	5.6		
Chronic Lower Respiratory Diseases	36.3		
Alzheimer's Disease	37.7		
Cerebrovascular Disease	39.6		
 Accidents 	40.2		
 Motor Vehicle Accidents 	9.3		
Diabetes	26.6		
Influenza and Pneumonia	18.9		
Intentional Self-Harm (Suicide)	15.2		
 Chronic Liver Disease and Cirrhosis 	8.8		
Infant Mortality	3.92		
	T-1-1		

Leading Causes of Death	Total Deaths
1. Malignant Neoplasms (Cancer)	418
2. Heart Disease	411
3. Alzheimer's Disease	120
Cerebrovascular Disease	112
Chronic Lower Respiratory Diseases	100
6. Accidents	95
7. Diabetes	70
Influenza and Pneumonia	55
Unspecified Dementia	30
10. Intentional Self-Harm (Suicide)	29
Percent of Deaths due to tobacco use	14.7
Median age at death	82

See technical notes for more information.

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

³All mortality rates except infant mortality are ageadjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

Brule County

Demographic Information

Brule County is located in the south central part of the state and averages 6.4 persons per square mile.

2017 Population Information

Subject	Number	Percent
Total population	5,312	100.0
White	4,528	85.2
Black or African American	23	0.4
American Indian & Alaska Native	548	10.3
Asian	21	0.4
Native Hawaiian & Other Pacific Islander	2	0.0
Two or More Races	190	3.6
Under 5 years	335	6.3
Under 18 years	1,410	26.5
65 years and over	956	18.0

Health Status Indicators 2013-2017

Natality	
Percent of Low Birth Weight Infants	5.8
Percent of Mothers Receiving	
Care in 1st Trimester	64.6
Percent of Mothers Who	
Used Tobacco While Pregnant ¹	15.2
Percent of Births Less Than 37 Wks. of Gestation	9.1
Average Age of Mother	27.7
Teenage Birth Rate ²	4.9
Percent White Births	76.2
Percent American Indian Births	19.5
Percent Unmarried	35.3
Percent WIC births	35.9
 Percent Breastfeeding at discharge 	69.9
Percent Payment-Private Insurance	63.1
Percent Payment-Medicaid	32.0
Percent C-Section	34.2

Notality

Mortality ³	
All Causes	675.9
Heart Disease	163.2
Malignant Neoplasms (Cancer)	141.5
Trachea, Bronchus, & Lung	24.2
Colon, Rectum, & Anus	19.2
Female Breast	20.8
Pancreas	9.9
Prostate	41.1
Leukemia	10.6
Chronic Lower Respiratory Diseases	34.6
Alzheimer's Disease	31.3
 Cerebrovascular Disease 	15.8
Accidents	41.2
Motor Vehicle Accidents	11.4
Diabetes	27.7
Influenza and Pneumonia	18.7
Intentional Self-Harm (Suicide)	18.1
Chronic Liver Disease and Cirrhosis	8.2
Infant Mortality	8.22
Landing Organisation	Total

Leading Causes of Death	Deaths
1. Heart Disease	68
Malignant Neoplasms (Cancer)	55
Alzheimer's Disease	15
T4. Accidents	14
T4. Chronic Lower Respiratory Diseases	14
6. Diabetes	10
Influenza and Pneumonia	8
Cerebrovascular Disease	7
T9. Intentional Self-Harm (Suicide)	5
T9. Pneumonitis Due to Solids and Liquids	5
Percent of Deaths due to tobacco use	13.2

•Denotes a health status indicator which is significantly lower than the state average.

See technical notes for more information.

Median age at death

 $[\]circ \mbox{Denotes}$ a health status indicator which is significantly higher than the state average.

³All mortality rates except infant mortality are ageadjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.


¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Source: United States Census Bureau, 2017 Population Estimates

Buffalo County

Demographic Information



Buffalo County is located in the central portion of the state and averages 4.1 persons per square mile.

2017 Population Information

Subject	Number	Percent
Total population	1,999	100.0
White	317	15.9
Black or African American	15	8.0
American Indian & Alaska Native	1,626	81.3
Asian	1	0.1
Native Hawaiian & Other Pacific Islander	0	0.0
Two or More Races	40	2.0
Under 5 years	226	11.3
Under 18 years	767	38.4
65 years and over	145	7.3

Health Status Indicators 2013-2017

Natality	
Percent of Low Birth Weight Infants	8.4
 Percent of Mothers Receiving 	
Care in 1st Trimester	29.5
 Percent of Mothers Who 	
Used Tobacco While Pregnant ¹	29.8
Percent of Births Less Than 37 Wks. of Gestation	10.8
 Average Age of Mother 	25.7
○ Teenage Birth Rate ²	64.4
Percent White Births	13.5
Percent American Indian Births	82.9
Percent Unmarried	74.6
Percent WIC births	65.3
Percent Breastfeeding at discharge	38.8
Percent Payment-Private Insurance Percent Payment-Madical description	12.7
Percent Payment-Medicaid Percent C Section	75.0
Percent C-Section	22.2

Mortality ³			
o All Causes	1,552.9		
Heart Disease	208.8		
Malignant Neoplasms (Cancer)	281.2		
 Trachea, Bronchus, & Lung 	140.7		
Colon, Rectum, & Anus	69.0		
Female Breast	LNE		
Pancreas	LNE		
Prostate	LNE		
Leukemia	LNE		
Chronic Lower Respiratory Diseases	49.4		
Alzheimer's Disease	LNE		
Cerebrovascular Disease	44.2		
 Accidents 	141.5		
Motor Vehicle Accidents	38.2		
 Diabetes 	124.6		
Influenza and Pneumonia	56.4		
 Intentional Self-Harm (Suicide) 	110.9		
 Chronic Liver Disease and Cirrhosis 	113.6		
Infant Mortality	11.90		

Leading Causes of Death	Total Deaths
Malignant Neoplasms (Cancer)	19
2. Heart Disease	15
3. Accidents	10
T4. Intentional Self-Harm (Suicide)	9
T4. Chronic Liver Disease and Cirrhosis	9
6. Diabetes	8
Percent of Deaths due to tobacco use Median age at death	19.6 61

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Source: United States Census Bureau, 2017 Population

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

³All mortality rates except infant mortality are ageadjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

Butte County

Demographic Information

Butte County is located on the western border of the state and averages 4.5 people per square mile.

2017 Population Information

Subject	Number	Percent
Total population	10,107	100.0
White	9,521	94.2
Black or African American	57	0.6
American Indian & Alaska Native	231	2.3
Asian	44	0.4
Native Hawaiian & Other Pacific Islander	9	0.1
Two or More Races	245	2.4
Under 5 years	612	6.1
Under 18 years	2,441	24.2
65 years and over	1,974	19.5

Source: United States Census Bureau, 2017 Population

Estimates

Natality

Health Status Indicators 2013-2017

ivalanty	
Percent of Low Birth Weight Infants	7.2
Percent of Mothers Receiving	
Care in 1st Trimester	77.9
 Percent of Mothers Who 	
Used Tobacco While Pregnant ¹	20.7
Percent of Births Less Than 37 Wks. of Gestation	9.2
 Average Age of Mother 	27.1
Teenage Birth Rate ²	11.8
Percent White Births	93.6
Percent American Indian Births	2.7
Percent Unmarried	35.9
 Percent WIC births 	45.2
Percent Breastfeeding at discharge	85.4
Percent Payment-Private Insurance	54.9
 Percent Payment-Medicaid 	39.7
Percent C-Section	22.9

Mortality ³	
All Causes	735.5
 Heart Disease 	185.6
Malignant Neoplasms (Cancer)	170.6
Trachea, Bronchus, & Lung	48.8
Colon, Rectum, & Anus	23.5
Female Breast	13.7
Pancreas	7.4
Prostate	16.7
Leukemia	3.4
 Chronic Lower Respiratory Diseases 	62.9
Alzheimer's Disease	29.2
Cerebrovascular Disease	34.1
Accidents	50.2
Motor Vehicle Accidents	22.0
Diabetes	15.0
 Influenza and Pneumonia 	7.2
 Intentional Self-Harm (Suicide) 	6.6
Chronic Liver Disease and Cirrhosis	17.3

Leading Causes of Death	Total Deaths
1. Heart Disease	140
2. Malignant Neoplasms (Cancer)	124
3. Chronic Lower Respiratory Diseases	46
4. Accidents	28
5. Cerebrovascular Disease	24
6. Alzheimer's Disease	23
7. Diabetes	12
8. Chronic Liver Disease and Cirrhosis	10
9. Septicemia	8
T10. Unspecified Dementia	6
T10. Parkinson's Disease	6
Percent of Deaths due to tobacco use Median age at death	20.0 79

7.80

Infant Mortality

See technical notes for more information.

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

³All mortality rates except infant mortality are ageadjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

Campbell County

Demographic Information

Campbell County is located in the north central portion of the state and averages 2.0 persons per square mile.

2017 Population Information

Subject	Number	Percent
Total population White	1,379 1.348	100.0 97.8
Black or African American	3	0.2
American Indian & Alaska Native	13	0.9
Asian	6	0.4
Native Hawaiian & Other Pacific Islander	0	0.0
Two or More Races	9	0.7
Under 5 years	56	4.1
Under 18 years	211	15.3
65 years and over	379	27.5

Health Status Indicators 2013-2017

Natality	
Percent of Low Birth Weight Infants	LNE
Percent of Mothers Receiving	
Care in 1st Trimester	79.0
 Percent of Mothers Who 	
Used Tobacco While Pregnant ¹	4.8
Percent of Births Less Than 37 Wks. of Gestation	9.7
Average Age of Mother	29.1
Teenage Birth Rate ²	LNE
Percent White Births	90.3
Percent American Indian Births	LNE
 Percent Unmarried 	16.1
 Percent WIC births 	17.7
Percent Breastfeeding at discharge	86.9
Percent Payment-Private Insurance	72.9
 Percent Payment-Medicaid 	16.9
Percent C-Section	29.0

Mortality ³	
All Causes	601.1
Heart Disease	119.2
Malignant Neoplasms (Cancer)	110.7
Trachea, Bronchus, & Lung	40.8
Colon, Rectum, & Anus	LNE
Female Breast	LNE
Pancreas	LNE
Prostate	LNE
Leukemia	LNE
Chronic Lower Respiratory Diseases	25.0
Alzheimer's Disease	18.5
Cerebrovascular Disease	31.0
Accidents	123.0
Motor Vehicle Accidents	LNE
Diabetes	37.1
Influenza and Pneumonia	LNE
Intentional Self-Harm (Suicide)	LNE
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	LNE

Total Deaths
18
15
8
5
5
20.8 77

See technical notes for more information.

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

[•]Denotes a health status indicator which is significantly lower than the state average.

 $[\]circ \mbox{Denotes}$ a health status indicator which is significantly higher than the state average.

³All mortality rates except infant mortality are ageadjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

Charles Mix County

Demographic Information

Charles Mix County is located in the south central area of the state and averages 8.3 persons per square mile.

2017 Population Information

Subject	Number	Percent
Total population	9,428	100.0
White	6,025	63.9
Black or African American	54	0.6
American Indian & Alaska Native	3,012	31.9
Asian	25	0.3
Native Hawaiian & Other Pacific Islander	1	0.0
Two or More Races	311	3.3
Under 5 years	829	8.8
Under 18 years	2,782	29.5
65 years and over	1,713	18.2

Health Status Indicators 2013-2017

Natality	
Percent of Low Birth Weight Infants	5.9
 Percent of Mothers Receiving 	
Care in 1st Trimester	64.6
 Percent of Mothers Who 	
Used Tobacco While Pregnant ¹	21.6
Percent of Births Less Than 37 Wks. of Gestation	8.4
 Average Age of Mother 	26.6
 Teenage Birth Rate² 	24.8
Percent White Births	43.2
Percent American Indian Births	50.6
 Percent Unmarried 	54.4
 Percent WIC births 	58.1
 Percent Breastfeeding at discharge 	68.5
Percent Payment-Private Insurance	37.3
Percent Payment-Medicaid	52.1
Percent C-Section	30.8

Mortality ³			
All Causes	857.9		
Heart Disease	183.1		
Malignant Neoplasms (Cancer)	172.9		
Trachea, Bronchus, & Lung	49.3		
Colon, Rectum, & Anus	17.8		
Female Breast	26.3		
Pancreas	7.3		
Prostate	28.5		
Leukemia	4.3		
Chronic Lower Respiratory Diseases	39.7		
Alzheimer's Disease	49.9		
Cerebrovascular Disease	41.3		
 Accidents 	79.7		
Motor Vehicle Accidents	22.2		
 Diabetes 	45.2		
Influenza and Pneumonia	21.6		
 Intentional Self-Harm (Suicide) 	44.4		
Chronic Liver Disease and Cirrhosis	31.1		
Infant Mortality	3.69		

Leading Causes of Death	Total Deaths
1. Heart Disease	120
Malignant Neoplasms (Cancer)	116
3. Alzheimer's Disease	41
4. Accidents	40
Cerebrovascular Disease	29
6. Chronic Lower Respiratory Diseases	27
7. Diabetes	24
8. Influenza and Pneumonia	18
Intentional Self-Harm (Suicide)	16
10. Chronic Liver Disease and Cirrhosis	12
Percent of Deaths due to tobacco use	15.6
Median age at death	77

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Source: United States Census Bureau, 2017 Population

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

³All mortality rates except infant mortality are ageadjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

Clark County

Demographic Information

Clark County is located in east central South Dakota and averages 3.9 persons per square mile.

2017 Population Information

Subject	Number	Percent
Total population	3,668	100.0
White	3,539	96.5
Black or African American	63	1.7
American Indian & Alaska Native	12	0.3
Asian	18	0.5
Native Hawaiian & Other Pacific Islander	0	0.0
Two or More Races	36	1.0
Under 5 years	372	10.1
Under 18 years	938	25.6
65 years and over	778	21.2

Health Status Indicators 2013-2017

Natality	
Percent of Low Birth Weight Infants	6.5
Percent of Mothers Receiving	
Care in 1st Trimester	65.9
 Percent of Mothers Who 	
Used Tobacco While Pregnant ¹	9.4
Percent of Births Less Than 37 Wks. of Gestation	6.1
Average Age of Mother	28.4
Teenage Birth Rate ²	9.6
Percent White Births	94.2
Percent American Indian Births	1.6
 Percent Unmarried 	15.2
 Percent WIC births 	21.0
Percent Breastfeeding at discharge	88.0
 Percent Payment-Private Insurance 	82.3
 Percent Payment-Medicaid 	14.5
Percent C-Section	21.9

Mortality ³			
All Causes	697.7		
Heart Disease	128.0		
Malignant Neoplasms (Cancer)	153.7		
Trachea, Bronchus, & Lung	34.7		
Colon, Rectum, & Anus	14.0		
Female Breast	13.8		
Pancreas	19.2		
Prostate	20.8		
Leukemia	16.5		
Chronic Lower Respiratory Diseases	33.6		
Alzheimer's Disease	32.8		
Cerebrovascular Disease	51.1		
Accidents	64.2		
Motor Vehicle Accidents	18.8		
Diabetes	35.7		
Influenza and Pneumonia	16.6		
Intentional Self-Harm (Suicide)	LNE		
Chronic Liver Disease and Cirrhosis	LNE		
Infant Mortality	9.68		

Leading Causes of Death	
Ecading Gadaca of Death	Deaths
Malignant Neoplasms (Cancer)	51
2. Heart Disease	39
3. Accidents	19
Cerebrovascular Disease	17
5. Alzheimer's Disease	13
6. Chronic Lower Respiratory Diseases	12
7. Diabetes	10
8. Influenza and Pneumonia	6
9. Unspecified Dementia	5
Percent of Deaths due to tobacco use	14.3
Median age at death	83

Total

See technical notes for more information.

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

³All mortality rates except infant mortality are ageadjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

Clay County

Demographic Information

Clay County is located in the southeastern part of the state and averages 33.6 persons per square mile.

2017 Population Information

Subject	Number	Percent
Total population	13,990	100.0
White	12,540	89.6
Black or African American	244	1.7
American Indian & Alaska Native	495	3.5
Asian	361	2.6
Native Hawaiian & Other Pacific Islander	10	0.1
Two or More Races	340	2.4
Under 5 years	756	5.4
Under 18 years	2,479	17.7
65 years and over	1,645	11.8

Health Status Indicators 2013-2017

Natality	
Percent of Low Birth Weight Infants	5.4
 Percent of Mothers Receiving 	
Care in 1st Trimester	80.7
 Percent of Mothers Who 	
Used Tobacco While Pregnant ¹	9.9
Percent of Births Less Than 37 Wks. of Gestation	7.0
 Average Age of Mother 	28.7
 Teenage Birth Rate² 	5.9
Percent White Births	84.3
Percent American Indian Births	6.9
 Percent Unmarried 	29.1
Percent WIC births	29.2
Percent Breastfeeding at discharge	81.0
 Percent Payment-Private Insurance 	68.2
Percent Payment-Medicaid	28.2
 Percent C-Section 	31.7

Notality

Mortality ³	
All Causes	785.7
 Heart Disease 	240.6
Malignant Neoplasms (Cancer)	178.0
Trachea, Bronchus, & Lung	43.0
Colon, Rectum, & Anus	12.0
Female Breast	17.6
Pancreas	13.0
Prostate	26.4
Leukemia	9.8
Chronic Lower Respiratory Diseases	45.9
Alzheimer's Disease	24.2
Cerebrovascular Disease	42.5
 Accidents 	29.2
Motor Vehicle Accidents	16.5
Diabetes	20.2
Influenza and Pneumonia	28.8
 Intentional Self-Harm (Suicide) 	7.7
Chronic Liver Disease and Cirrhosis	9.9
Infant Mortality	LNE
Landing Course of Double	Total

Leading Causes of Death	Deaths
1. Heart Disease	167
Malignant Neoplasms (Cancer)	113
T3. Chronic Lower Respiratory Diseases	29
T3. Cerebrovascular Disease	29
5. Influenza and Pneumonia	20
6. Accidents	18
7. Alzheimer's Disease	17
8. Diabetes	14
T9. Unspecified Dementia	7
T9. Nephritis, Nephrotic Syndrome, and Nephrosis	7
Percent of Deaths due to tobacco use Median age at death	15.9 80

•Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

See technical notes for more information.

²Teenage Birth rate is live births per 1,000 females age 15-17.

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

³All mortality rates except infant mortality are ageadjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

Codington County

Demographic Information

Codington County is located in eastern South Dakota and averages 39.5 persons per square mile.

2017 Population Information

Number	Percent
28,099	100.0
26,556	94.5
199	0.7
712	2.5
193	0.7
3	0.0
436	1.6
4 000	0.5
,	6.5
,	24.2
4,787	17.0
	28,099 26,556 199 712 193 3

Health Status Indicators 2013-2017

Natality	
Percent of Low Birth Weight Infants	7.2
 Percent of Mothers Receiving 	
Care in 1st Trimester	79.9
 Percent of Mothers Who 	
Used Tobacco While Pregnant ¹	20.5
Percent of Births Less Than 37 Wks. of Gestation	7.9
 Average Age of Mother 	27.5
Teenage Birth Rate ²	7.4
Percent White Births	89.1
Percent American Indian Births	5.1
Percent Unmarried	37.5
Percent WIC births	35.1
Percent Breastfeeding at discharge	77.8
 Percent Payment-Private Insurance 	67.1
Percent Payment-Medicaid	29.8
Percent C-Section	23.5

Notality

Mortality ³	
All Causes	649.8
Heart Disease	149.9
Malignant Neoplasms (Cancer)	176.0
Trachea, Bronchus, & Lung	45.7
Colon, Rectum, & Anus	20.4
Female Breast	28.7
Pancreas	15.0
Prostate	17.7
Leukemia	7.0
Chronic Lower Respiratory Diseases	39.7
Alzheimer's Disease	30.5
Cerebrovascular Disease	32.5
Accidents	40.2
Motor Vehicle Accidents	16.2
Diabetes	23.7
 Influenza and Pneumonia 	11.4
 Intentional Self-Harm (Suicide) 	9.7
 Chronic Liver Disease and Cirrhosis 	5.5

2.68

Leading Causes of Death 1. Malignant Neoplasms (Cancer) 2. Heart Disease 3. Chronic Lower Respiratory Diseases 4. Cerebrovascular Disease 5. Accidents 6. Alzheimer's Disease	Total Deaths 319 292 73 64 63 62 44
8. Influenza and Pneumonia9. Unspecified Dementia10. Essential (Primary) Hypertension and Hypertensive Renal Disease	23 19 15
Percent of Deaths due to tobacco use Median age at death	21.1 81

Infant Mortality

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Source: United States Census Bureau, 2017 Population Estimates

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

³All mortality rates except infant mortality are ageadjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

Corson County

Demographic Information

Corson County is located in the north central portion of the state and averages 1.6 persons per square mile.

2017 Population Information

S	Subject	Number	Percent
Т	otal population	4,203	100.0
	White	1,279	30.4
	Black or African American	20	0.5
	American Indian & Alaska Native	2,738	65.1
	Asian	21	0.5
	Native Hawaiian & Other Pacific Islander	2	0.0
	Two or More Races	143	3.4
	Inder 5 years	439	10.4
	Inder 18 years	1,532	36.5
6	5 years and over	455	10.8

Source: United States Census Bureau, 2017 Population

Estimates

Health Status Indicators 2013-2017

Natality	
Percent of Low Birth Weight Infants	9.4
 Percent of Mothers Receiving 	
Care in 1st Trimester	42.9
 Percent of Mothers Who 	
Used Tobacco While Pregnant ¹	25.2
 Percent of Births Less Than 37 Wks. of Gestation 	14.0
 Average Age of Mother 	26.7
Teenage Birth Rate ²	17.6
Percent White Births	19.1
Percent American Indian Births	73.6
 Percent Unmarried 	71.2
 Percent WIC births 	62.7
 Percent Breastfeeding at discharge 	52.7
Percent Payment-Private Insurance	17.5
Percent Payment-Medicaid	74.2
Percent C-Section	28.5

1,292.0
315.3
209.3
31.2
31.4
LNE
LNE
LNE
LNE
38.8
LNE
26.3
132.3
23.2
64.6
33.6
58.5
64.9
8.51

Leading Causes of Death	Total Deaths
1. Heart Disease	57
Malignant Neoplasms (Cancer)	40
3. Accidents	25
4. Chronic Liver Disease and Cirrhosis	13
T5. Diabetes	12
T5. Intentional Self-Harm (Suicide)	12
7. Chronic Lower Respiratory Diseases	7
T8. Influenza and Pneumonia	6
T8. Nephritis, Nephrotic Syndrome, and Nephrosis	6
T8. III-Defined and Unknown Causes of Mortality	6
Percent of Deaths due to tobacco use Median age at death	16.3 63

See technical notes for more information.

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

³All mortality rates except infant mortality are ageadjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

Custer County

Demographic Information

Custer County is located in the southern Black Hills and averages 5.3 persons per square mile.

2017 Population Information

Subject	Number	Percent
Total population	8,691	100.0
White	8,067	92.8
Black or African American	52	0.6
American Indian & Alaska Native	338	3.9
Asian	50	0.6
Native Hawaiian & Other Pacific Islander	2	0.0
Two or More Races	182	2.1
Under 5 years	350	4.0
Under 18 years	1,369	15.8
65 years and over	2,487	28.6

Source: United States Census Bureau, 2017 Population

Estimates

Health Status Indicators 2013-2017

inatality	
Percent of Low Birth Weight Infants	8.1
Percent of Mothers Receiving	
Care in 1st Trimester	72.1
Percent of Mothers Who	
Used Tobacco While Pregnant ¹	18.8
Percent of Births Less Than 37 Wks. of Gestation	8.4
 Average Age of Mother 	28.7
Teenage Birth Rate ²	LNE
Percent White Births	85.4
Percent American Indian Births	7.3
Percent Unmarried	35.1
Percent WIC births	31.9
Percent Breastfeeding at discharge	87.2
Percent Payment-Private Insurance	52.4
Percent Payment-Medicaid	36.4
Percent C-Section	20.6

Natality

Mortality ³		
All Causes	644.9	
Heart Disease	134.2	
 Malignant Neoplasms (Cancer) 	120.6	
 Trachea, Bronchus, & Lung 	26.2	
Colon, Rectum, & Anus	13.5	
Female Breast	18.7	
Pancreas	8.9	
Prostate	10.1	
Leukemia	9.4	
 Chronic Lower Respiratory Diseases 	27.4	
Alzheimer's Disease	23.5	
Cerebrovascular Disease	36.8	
Accidents	66.6	
Motor Vehicle Accidents	18.2	
Diabetes	16.1	
Influenza and Pneumonia	28.8	
Intentional Self-Harm (Suicide)	22.3	
Chronic Liver Disease and Cirrhosis	18.6	
Infant Mortality	LNE	

Leading Causes of Death	Deaths
1. Heart Disease	106
Malignant Neoplasms (Cancer)	98
3. Accidents	35
Cerebrovascular Disease	27
Chronic Lower Respiratory Diseases	21
6. Influenza and Pneumonia	19
7. Alzheimer's Disease	17
T8. Diabetes	12
T8. Intentional Self-Harm (Suicide)	12
T8. Chronic Liver Disease and Cirrhosis	12
Percent of Deaths due to tobacco use	15.9
Median age at death	77

See technical notes for more information.

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

³All mortality rates except infant mortality are ageadjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

Davison County

Demographic Information

Davison County is located in eastern South Dakota and averages 44.8 persons per square mile.

2017 Population Information

Subject	Number	Percent
Total population	19,704	100.0
White	18,354	93.1
Black or African American	186	0.9
American Indian & Alaska Native	648	3.3
Asian	127	0.6
Native Hawaiian & Other Pacific Islander	17	0.1
Two or More Races	372	1.9
Under 5 years	1,238	6.3
Under 18 years	4,586	23.3
65 years and over	3,645	18.5

Health Status Indicators 2013-2017

Natality	
Percent of Low Birth Weight Infants	6.6
 Percent of Mothers Receiving 	
Care in 1st Trimester	80.3
 Percent of Mothers Who 	
Used Tobacco While Pregnant ¹	19.3
Percent of Births Less Than 37 Wks. of Gestation	9.6
 Average Age of Mother 	27.4
Teenage Birth Rate ²	9.5
Percent White Births	86.0
Percent American Indian Births	8.9
 Percent Unmarried 	41.6
Percent WIC births	36.0
 Percent Breastfeeding at discharge 	73.7
 Percent Payment-Private Insurance 	64.2
Percent Payment-Medicaid	30.8
 Percent C-Section 	29.2

Mortality ³		
All Causes	708.1	
Heart Disease	174.0	
Malignant Neoplasms (Cancer)	163.0	
Trachea, Bronchus, & Lung	47.1	
Colon, Rectum, & Anus	17.6	
Female Breast	12.0	
Pancreas	10.2	
Prostate	11.5	
Leukemia	9.3	
Chronic Lower Respiratory Diseases	35.6	
 Alzheimer's Disease 	25.1	
Cerebrovascular Disease	38.8	
Accidents	43.6	
 Motor Vehicle Accidents 	7.0	
Diabetes	26.2	
Influenza and Pneumonia	21.5	
Intentional Self-Harm (Suicide)	13.2	
Chronic Liver Disease and Cirrhosis	10.4	
Infant Mortality	6.35	

Leading Causes of Death	Deaths
1. Heart Disease	269
Malignant Neoplasms (Cancer)	237
Cerebrovascular Disease	70
4. Accidents	54
5. Chronic Lower Respiratory Diseases	51
6. Alzheimer's Disease	49
7. Influenza and Pneumonia	38
8. Unspecified Dementia	37
9. Diabetes	35
10. Essential (Primary) Hypertension and	22
Hypertensive Renal Disease	
••	
Percent of Deaths due to tobacco use	17.4
Median age at death	82

Total

See technical notes for more information.

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

³All mortality rates except infant mortality are ageadjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

Day County

Demographic Information

Day County is located in the northeastern part of the state and averages 5.6 persons per square mile.

2017 Population Information

Subject	Number	Percent
Total population	5,521	100.0
White	4,831	87.5
Black or African American	21	0.4
American Indian & Alaska Native	531	9.6
Asian	39	0.7
Native Hawaiian & Other Pacific Islander	0	0.0
Two or More Races	99	1.8
Under 5 years	306	5.5
Under 18 years	1,237	22.4
65 years and over	1,386	25.1

Source: United States Census Bureau, 2017 Population

Estimates

Health Status Indicators 2013-2017

ivatanty	
Percent of Low Birth Weight Infants	5.1
Percent of Mothers Receiving	
Care in 1st Trimester	72.6
 Percent of Mothers Who 	
Used Tobacco While Pregnant ¹	24.0
Percent of Births Less Than 37 Wks. of Gestation	5.8
Average Age of Mother	28.0
Teenage Birth Rate ²	13.8
Percent White Births	75.4
Percent American Indian Births	19.1
Percent Unmarried	36.0
Percent WIC births	40.4
Percent Breastfeeding at discharge	76.5
Percent Payment-Private Insurance	62.1
Percent Payment-Medicaid	33.8
Percent C-Section	28.0

Natality

Mortality ³			
All Causes	702.1		
Heart Disease	155.2		
Malignant Neoplasms (Cancer)	131.1		
Trachea, Bronchus, & Lung	33.3		
Colon, Rectum, & Anus	14.5		
Female Breast	16.5		
Pancreas	12.1		
Prostate	LNE		
Leukemia	LNE		
Chronic Lower Respiratory Diseases	30.8		
Alzheimer's Disease	37.8		
Cerebrovascular Disease	49.6		
Accidents	76.9		
Motor Vehicle Accidents	33.5		
Diabetes	26.5		
Influenza and Pneumonia	13.6		
Intentional Self-Harm (Suicide)	13.0		
Chronic Liver Disease and Cirrhosis	18.8		
Infant Mortality	LNE		

Leading Causes of Death	lotai	
•	Deaths	
1. Heart Disease	91	
Malignant Neoplasms (Cancer)	62	
Cerebrovascular Disease	32	
4. Accidents	26	
5. Alzheimer's Disease	25	
6. Chronic Lower Respiratory Diseases	18	
7. Diabetes	14	
T8. Influenza and Pneumonia	9	
T8. Parkinson's Disease	9	
10. Essential (Primary) Hypertension and Hypertensive Renal Disease	7	
Percent of Deaths due to tobacco use	15.1	
Median age at death	83	

See technical notes for more information.

[•]Denotes a health status indicator which is significantly lower than the state average.

 $[\]circ \mbox{Denotes}$ a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

³All mortality rates except infant mortality are ageadjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

Deuel County

Demographic Information

Deuel County borders Minnesota and averages 7.0 persons per square mile.

2017 Population Information

Subject	Number	Percent
Total population	4,281	100.0
White	4,169	97.4
Black or African American	37	0.9
American Indian & Alaska Native	21	0.5
Asian	8	0.2
Native Hawaiian & Other Pacific Islander	0	0.0
Two or More Races	46	1.1
Under 5 years	266	6.2
Under 18 years	996	23.3
65 years and over	948	22.1

Health Status Indicators 2013-2017

Natality	
Percent of Low Birth Weight Infants	4.3
Percent of Mothers Receiving	
Care in 1st Trimester	80.7
 Percent of Mothers Who 	
Used Tobacco While Pregnant ¹	9.5
Percent of Births Less Than 37 Wks. of Gestation	6.3
Average Age of Mother	28.3
Teenage Birth Rate ²	LNE
Percent White Births	97.2
Percent American Indian Births	LNE
 Percent Unmarried 	20.1
 Percent WIC births 	25.6
Percent Breastfeeding at discharge	85.8
 Percent Payment-Private Insurance 	73.7
Percent Payment-Medicaid	17.1
Percent C-Section	21.3

Mortality ³	
All Causes	620.1
Heart Disease	123.8
Malignant Neoplasms (Cancer)	156.1
Trachea, Bronchus, & Lung	36.5
Colon, Rectum, & Anus	8.0
Female Breast	LNE
Pancreas	11.4
Prostate	LNE
Leukemia	LNE
Chronic Lower Respiratory Diseases	41.2
Alzheimer's Disease	27.8
Cerebrovascular Disease	48.2
 Accidents 	25.8
Motor Vehicle Accidents	13.0
Diabetes	13.8
Influenza and Pneumonia	17.1
Intentional Self-Harm (Suicide)	LNE
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	LNE

Leading Causes of Death	lotal
· ·	Deaths
Malignant Neoplasms (Cancer)	55
2. Heart Disease	44
Cerebrovascular Disease	18
Chronic Lower Respiratory Diseases	17
5. Alzheimer's Disease	11
T6. Accidents	8
T6. Influenza and Pneumonia	8
T8. Essential (Primary) Hypertension and	6
Hypertensive Renal Disease	
T8. Septicemia	6
T8. Pneumonitis due to Solids and Liquids	6
Percent of Deaths due to tobacco use	17.4
Median age at death	82
modian ago at acam	

See technical notes for more information.

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

³All mortality rates except infant mortality are ageadjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

Dewey County

Demographic Information

Dewey County is located in the north central region of the state and averages 2.3 persons per square mile.

2017 Population Information

Subject	Number	Percent
Total population	5,835	100.0
White	1,222	20.9
Black or African American	23	0.4
American Indian & Alaska Native	4,320	74.0
Asian	14	0.2
Native Hawaiian & Other Pacific Islander	1	0.0
Two or More Races	255	4.4
Under 5 years	726	12.4
Under 18 years	2,154	36.9
65 years and over	559	9.6

•Denotes a health status indicator which is significantly lower than the state average. oDenotes a health status indicator which is significantly higher

than the state average. ¹Data for mothers who used tobacco are self-reported.

Health Status Indicators 2013-2017

o All Causes

Heart Disease

Pancreas

Prostate

Accidents

Diabetes

Infant Mortality

Leukemia

o Motor Vehicle Accidents

o Chronic Liver Disease and Cirrhosis

o Influenza and Pneumonia

Intentional Self-Harm (Suicide)

Natality	
 Percent of Low Birth Weight Infants 	9.4
 Percent of Mothers Receiving 	
Care in 1st Trimester	50.6
 Percent of Mothers Who 	
Used Tobacco While Pregnant ¹	20.5
 Percent of Births Less Than 37 Wks. of Gestation 	15.7
 Average Age of Mother 	26.1
○ Teenage Birth Rate ²	49.4
Percent White Births	10.2
Percent American Indian Births	83.7
 Percent Unmarried 	68.4
 Percent WIC births 	78.0
 Percent Breastfeeding at discharge 	54.0
 Percent Payment-Private Insurance 	16.5
Percent Payment-Medicaid	71.9
Percent C-Section	33.2

Mortality³ 1,353.3 206.6 Malignant Neoplasms (Cancer) 191.2 Trachea, Bronchus, & Lung 47.4 Colon, Rectum, & Anus 21.8 Female Breast 26.6 11.6 LNE LNE Chronic Lower Respiratory Diseases 44.8 • Alzheimer's Disease 14.6 Cerebrovascular Disease 32.3 175.8

79.1

101.5

65.7

46.6

95.0

10.44

Leading Causes of Death	Total Deaths
T1. Heart Disease	46
T1. Accidents	46
3. Malignant Neoplasms (Cancer)	45
T4. Diabetes	22
T4. Chronic Liver Disease and Cirrhosis	22
Influenza and Pneumonia	15
7. Intentional Self-Harm (Suicide)	12
T8. Chronic Lower Respiratory Diseases	10
T8. Nephritis, Nephrotic Syndrome, and Nephrosis	10
10. Cerebrovascular Disease	8
Percent of Deaths due to tobacco use	13.5
Median age at death	65

See technical notes for more information.

²Teenage Birth rate is live births per 1,000 females age 15-17.

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

³All mortality rates except infant mortality are ageadjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

Source: United States Census Bureau, 2017 Population **Estimates**

Douglas County

Demographic Information

Douglas County is located in south central South Dakota and averages 7.0 persons per square mile.

2017 Population Information

Subject	Number	Percent
Total population	2,931	100.0
White	2,805	95.7
Black or African American	16	0.5
American Indian & Alaska Native	67	2.3
Asian	3	0.1
Native Hawaiian & Other Pacific Islander	0	0.0
Two or More Races	40	1.4
Under 5 years	214	7.3
Under 18 years	729	24.9
65 years and over	707	24.1

Health Status Indicators 2013-2017

Natality		
	Percent of Low Birth Weight Infants	5.8
	Percent of Mothers Receiving	
	Care in 1st Trimester	76.1
	Percent of Mothers Who	
	Used Tobacco While Pregnant ¹	9.8
	Percent of Births Less Than 37 Wks. of Gestation	7.3
	Average Age of Mother	27.8
	Teenage Birth Rate ²	LNE
	Percent White Births	97.1
	Percent American Indian Births	LNE
	 Percent Unmarried 	12.1
	Percent WIC births	21.0
	Percent Breastfeeding at discharge	89.7
	 Percent Payment-Private Insurance 	71.4
	Percent Payment-Medicaid	15.3
	Percent C-Section	28.6

Motality

Mortality ³			
All Causes	627.8		
Heart Disease	102.0		
Malignant Neoplasms (Cancer)	150.5		
Trachea, Bronchus, & Lung	34.3		
Colon, Rectum, & Anus	21.3		
Female Breast	51.3		
Pancreas	11.4		
Prostate	LNE		
Leukemia	LNE		
Chronic Lower Respiratory Diseases	54.2		
Alzheimer's Disease	47.5		
Cerebrovascular Disease	20.8		
Accidents	55.6		
Motor Vehicle Accidents	LNE		
Diabetes	25.3		
Influenza and Pneumonia	12.8		
Intentional Self-Harm (Suicide)	LNE		
Chronic Liver Disease and Cirrhosis	29.1		
Infant Mortality	LNE		

Leading Causes of Death	Total Deaths
1. Heart Disease	43
Malignant Neoplasms (Cancer)	41
3. Alzheimer's Disease	22
4. Accidents	18
Chronic Lower Respiratory Diseases	16
T6. Cerebrovascular Disease	8
T6. Diabetes	8
8. Influenza and Pneumonia	6
Essential (Primary) Hypertension and Hypertensive Renal Disease	5
Percent of Deaths due to tobacco use	16.3
Median age at death	86

See technical notes for more information.

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

³All mortality rates except infant mortality are ageadjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

Edmunds County

Demographic Information

Edmunds County is located in the north central region of the state and averages 3.6 persons per square mile.

2017 Population Information

Subject	Number	Percent
Total population	3,919	100.0
White	3,796	96.9
Black or African American	16	0.4
American Indian & Alaska Native	39	1.0
Asian	28	0.7
Native Hawaiian & Other Pacific Islander	0	0.0
Two or More Races	40	1.0
Under 5 years	279	7.1
Under 18 years	909	23.2
65 years and over	840	21.4

Health Status Indicators 2013-2017

Natality	
Percent of Low Birth Weight Infants	3.8
Percent of Mothers Receiving	
Care in 1st Trimester	64.3
 Percent of Mothers Who 	
Used Tobacco While Pregnant ¹	7.7
Percent of Births Less Than 37 Wks. of Gestation	5.5
 Average Age of Mother 	28.6
Teenage Birth Rate ²	LNE
Percent White Births	95.8
Percent American Indian Births	1.7
Percent Unmarried	14.8
Percent WIC births	16.1
Percent Breastfeeding at discharge	89.4
Percent Payment-Private Insurance	84.3
Percent Payment-Medicaid	13.6
Percent C-Section	20.3

Mortality ³		
All Causes	598.8	
Heart Disease	126.7	
Malignant Neoplasms (Cancer)	128.0	
Trachea, Bronchus, & Lung	41.6	
Colon, Rectum, & Anus	8.1	
Female Breast	LNE	
Pancreas	17.3	
Prostate	18.3	
Leukemia	LNE	
Chronic Lower Respiratory Diseases	31.5	
Alzheimer's Disease	27.3	
Cerebrovascular Disease	26.5	
Accidents	55.7	
Motor Vehicle Accidents	18.0	
Diabetes	25.7	
Influenza and Pneumonia	19.8	
Intentional Self-Harm (Suicide)	LNE	
Chronic Liver Disease and Cirrhosis	LNE	
Infant Mortality	LNE	

Leading Causes of Death	Total Deaths
1. Heart Disease	47
Malignant Neoplasms (Cancer)	40
3. Accidents	16
T4. Chronic Lower Respiratory Diseases	12
T4. Alzheimer's Disease	12
Cerebrovascular Disease	10
7. Influenza and Pneumonia	9
T8. Diabetes	7
T8. Unspecified Dementia	7
T8. Essential (Primary) Hypertension and Hypertensive Renal Disease	7
Percent of Deaths due to tobacco use	17.5
Median age at death	83

•Denotes a health status indicator which is significantly lower than the state average.

See technical notes for more information. Source: South Dakota Department of Health, Office of Health Statistics

Estimates

Source: United States Census Bureau, 2017 Population

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

oDenotes a health status indicator which is significantly higher than the state average.

³All mortality rates except infant mortality are ageadjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

Fall River County

Demographic Information

Fall River County is located in the southwestern corner of the state and averages 4.1 persons per square mile.

2017 Population Information

Subject	Number	Percent
Total population	6,687	100.0
White	5,866	87.7
Black or African American	90	1.3
American Indian & Alaska Native	471	7.0
Asian	42	0.6
Native Hawaiian & Other Pacific Islander	2	0.0
Two or More Races	216	3.2
Under 5 years	304	4.5
Under 18 years	1,171	17.5
65 years and over	1,896	28.4

Source: United States Census Bureau, 2017 Population

Estimates

Health Status Indicators 2013-2017

inatality	
Percent of Low Birth Weight Infants	7.5
Percent of Mothers Receiving	
Care in 1st Trimester	68.2
Percent of Mothers Who	
Used Tobacco While Pregnant ¹	17.5
Percent of Births Less Than 37 Wks. of Gestation	8.2
Average Age of Mother	27.3
Teenage Birth Rate ²	10.5
Percent White Births	80.7
Percent American Indian Births	8.9
Percent Unmarried	42.3
 Percent WIC births 	41.1
Percent Breastfeeding at discharge	80.7
 Percent Payment-Private Insurance 	46.8
Percent Payment-Medicaid	38.5
Percent C-Section	17.4

Natality

Mortality ³		
o All Causes	974.4	
 Heart Disease 	216.9	
Malignant Neoplasms (Cancer)	185.9	
Trachea, Bronchus, & Lung	35.5	
Colon, Rectum, & Anus	24.5	
Female Breast	24.9	
Pancreas	8.1	
Prostate	22.7	
Leukemia	7.0	
 Chronic Lower Respiratory Diseases 	84.5	
Alzheimer's Disease	36.8	
Cerebrovascular Disease	42.5	
Accidents	72.3	
Motor Vehicle Accidents	26.9	
Diabetes	37.1	
Influenza and Pneumonia	21.9	
Intentional Self-Harm (Suicide)	23.3	
Chronic Liver Disease and Cirrhosis	19.5	
Infant Mortality	LNE	
	T-1-1	

Leading Causes of Death	Total Deaths
1. Heart Disease	138
Malignant Neoplasms (Cancer)	120
Chronic Lower Respiratory Diseases	59
4. Accidents	32
Cerebrovascular Disease	28
6. Alzheimer's Disease	26
7. Diabetes	24
8. Influenza and Pneumonia	13
Unspecified Dementia	12
10. Intentional Self-Harm (Suicide)	10
Percent of Deaths due to tobacco use	23.7
Median age at death	79

See technical notes for more information.

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

³All mortality rates except infant mortality are ageadjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

Faulk County

Demographic Information

Faulk County is located in north central South Dakota and averages 2.4 persons per square mile.

2017 Population Information

C...l. ! - -4

Subject	Number	Percent
Total population	2,329	100.0
White	2,282	98.0
Black or African American	9	0.4
American Indian & Alaska Native	11	0.5
Asian	10	0.4
Native Hawaiian & Other Pacific Islander	0	0.0
Two or More Races	17	0.7
Under 5 years	182	7.8
Under 18 years	566	24.3
65 years and over	540	23.2

Health Status Indicators 2013-2017

Natality	
Percent of Low Birth Weight Infants	8.0
 Percent of Mothers Receiving 	
Care in 1st Trimester	56.8
 Percent of Mothers Who 	
Used Tobacco While Pregnant ¹	2.5
Percent of Births Less Than 37 Wks. of Gestation	5.6
 Average Age of Mother 	29.2
Teenage Birth Rate ²	LNE
Percent White Births	99.4
Percent American Indian Births	LNE
 Percent Unmarried 	6.8
 Percent WIC births 	13.7
Percent Breastfeeding at discharge	90.7
 Percent Payment-Private Insurance 	87.7
Percent Payment-Medicaid	7.4
Percent C-Section	22.2

Mortality ³	
All Causes	697.8
Heart Disease	128.3
Malignant Neoplasms (Cancer)	159.4
Trachea, Bronchus, & Lung	21.3
Colon, Rectum, & Anus	LNE
Female Breast	36.8
Pancreas	16.5
Prostate	26.4
Leukemia	LNE
Chronic Lower Respiratory Diseases	49.5
Alzheimer's Disease	40.1
Cerebrovascular Disease	33.2
Accidents	42.4
Motor Vehicle Accidents	LNE
Diabetes	26.5
Influenza and Pneumonia	10.2
Intentional Self-Harm (Suicide)	26.3
Chronic Liver Disease and Cirrhosis	28.6
Infant Mortality	LNE
	Total

Leading Causes of Death	Deaths
T1. Heart Disease	34
T1. Malignant Neoplasms (Cancer)	34
Alzheimer's Disease	13
Chronic Lower Respiratory Diseases	12
5. Accidents	10
Cerebrovascular Disease	9
7. Diabetes	5
Percent of Deaths due to tobacco use Median age at death	11.3 83

See technical notes for more information.

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

³All mortality rates except infant mortality are ageadjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

Grant County

Demographic Information

Grant County borders Minnesota in northeastern South Dakota and averages 10.8 persons per square mile.

2017 Population Information

Subject	Number	Percent
Total population	7,061	100.0
White	6,799	96.3
Black or African American	50	0.7
American Indian & Alaska Native	85	1.2
Asian	28	0.4
Native Hawaiian & Other Pacific Islander	5	0.1
Two or More Races	94	1.3
Under 5 years	440	6.2
Under 18 years	1,598	22.6
65 years and over	1,480	21.0

Source: United States Census Bureau, 2017 Population Estimates

Health Status Indicators 2013-2017

Natality	
Percent of Low Birth Weight Infants	5.0
Percent of Mothers Receiving	
Care in 1st Trimester	78.7
Percent of Mothers Who	
Used Tobacco While Pregnant ¹	12.8
Percent of Births Less Than 37 Wks. of Gestation	6.7
Average Age of Mother	28.0
Teenage Birth Rate ²	LNE
Percent White Births	91.2
Percent American Indian Births	1.4
 Percent Unmarried 	25.1
Percent WIC births	28.7
Percent Breastfeeding at discharge	83.2
 Percent Payment-Private Insurance 	69.9
 Percent Payment-Medicaid 	20.4
Percent C-Section	26.1

Motality

Mortality ³	
All Causes	767.5
Heart Disease	176.3
Malignant Neoplasms (Cancer)	141.4
Trachea, Bronchus, & Lung	42.4
Colon, Rectum, & Anus	8.1
Female Breast	17.9
Pancreas	8.1
Prostate	15.6
Leukemia	6.9
Chronic Lower Respiratory Diseases	56.0
Alzheimer's Disease	50.3
 Cerebrovascular Disease 	59.5
Accidents	45.7
Motor Vehicle Accidents	24.1
Diabetes	28.9
Influenza and Pneumonia	24.5
Intentional Self-Harm (Suicide)	24.7
 Chronic Liver Disease and Cirrhosis 	5.0
Infant Mortality	LNE

Leading Causes of Death	Deaths
Heart Disease Malignant Neoplasms (Cancer) Cerebrovascular Disease Chronic Lower Respiratory Diseases Alzheimer's Disease Accidents Diabetes Influenza and Pneumonia Intentional Self-Harm (Suicide) T10. Essential (Primary) Hypertension and Hypertensive Renal Disease Nephritis, Nephrotic Syndrome, and Nephrosis	110 81 39 35 34 22 17 15 9 7
Percent of Deaths due to tobacco use Median age at death	15.7 83

Total

See technical notes for more information.

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

³All mortality rates except infant mortality are ageadjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

Gregory County

Demographic Information

Gregory County borders the west bank of the Missouri River and the state of Nebraska and averages 4.2 persons per square mile.

2017 Population Information

Subject	Number	Percent
Total population	4,226	100.0
White	3,757	88.9
Black or African American	15	0.4
American Indian & Alaska Native	306	7.2
Asian	19	0.4
Native Hawaiian & Other Pacific Islander	4	0.1
Two or More Races	125	3.0
Under 5 years	269	6.4
Under 18 years	981	23.2
65 years and over	1,046	24.8

Health Status Indicators 2013-2017

Natality	
 Percent of Low Birth Weight Infants 	3.4
Percent of Mothers Receiving	
Care in 1st Trimester	65.3
Percent of Mothers Who	
Used Tobacco While Pregnant ¹	11.3
Percent of Births Less Than 37 Wks. of Gestation	6.4
Average Age of Mother	28.0
Teenage Birth Rate ²	16.0
Percent White Births	77.2
Percent American Indian Births	15.7
Percent Unmarried	34.5
 Percent WIC births 	41.7
Percent Breastfeeding at discharge	78.9
Percent Payment-Private Insurance	57.7
Percent Payment-Medicaid	38.6
Percent C-Section	27.3

Motality

Mortality ³	
All Causes	800.1
Heart Disease	165.9
Malignant Neoplasms (Cancer)	176.6
Trachea, Bronchus, & Lung	42.6
Colon, Rectum, & Anus	13.9
Female Breast	20.4
Pancreas	21.0
Prostate	LNE
Leukemia	LNE
Chronic Lower Respiratory Diseases	54.4
Alzheimer's Disease	27.1
Cerebrovascular Disease	39.0
Accidents	85.5
 Motor Vehicle Accidents 	59.6
Diabetes	27.3
 Influenza and Pneumonia 	5.8
Intentional Self-Harm (Suicide)	13.0
Chronic Liver Disease and Cirrhosis	31.3
Infant Mortality	11.24

Leading Causes of Death	Total Deaths
1. Heart Disease	69
2. Malignant Neoplasms (Cancer)	68
3. Chronic Lower Respiratory Diseases	26
T4. Accidents	19
T4. Cerebrovascular Disease	19
6. Alzheimer's Disease	15
7. Diabetes	11
Disorders of Lipoprotein Metabolism and Other Lipidemias	8
9. Chronic Liver Disease and Cirrhosis	7
Percent of Deaths due to tobacco use Median age at death	19.9 82

See technical notes for more information.

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

[•]Denotes a health status indicator which is significantly lower than the state average.

 $[\]circ \mbox{Denotes}$ a health status indicator which is significantly higher than the state average.

³All mortality rates except infant mortality are ageadjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

Haakon County

Demographic Information

Haakon County is located in the west central region of the state and averages 1.1 persons per square mile.

2017 Population Information

Subject	Number	Percent
Total population	1,943	100.0
White	1,822	93.8
Black or African American	5	0.3
American Indian & Alaska Native	45	2.3
Asian	5	0.3
Native Hawaiian & Other Pacific Islander	2	0.1
Two or More Races	64	3.3
Under 5 years	118	6.1
Under 18 years	462	23.8
65 years and over	460	23.7

Health Status Indicators 2013-2017

Natality	
Percent of Low Birth Weight Infants	2.8
Percent of Mothers Receiving	
Care in 1st Trimester	73.8
Percent of Mothers Who	
Used Tobacco While Pregnant ¹	15.7
Percent of Births Less Than 37 Wks. of Gestation	5.6
Average Age of Mother	28.4
Teenage Birth Rate ²	LNE
Percent White Births	86.1
Percent American Indian Births	2.8
 Percent Unmarried 	24.1
Percent WIC births	30.8
Percent Breastfeeding at discharge	84.3
Percent Payment-Private Insurance	72.2
Percent Payment-Medicaid	25.0
Percent C-Section	21.3

Mortality ³	
All Causes	727.0
Heart Disease	140.6
Malignant Neoplasms (Cancer)	166.4
Trachea, Bronchus, & Lung	47.4
Colon, Rectum, & Anus	26.2
Female Breast	LNE
Pancreas	LNE
Prostate	LNE
Leukemia	33.0
Chronic Lower Respiratory Diseases	76.0
Alzheimer's Disease	LNE
Cerebrovascular Disease	36.2
 Accidents 	23.7
Motor Vehicle Accidents	LNE
Diabetes	LNE
Influenza and Pneumonia	31.2
Intentional Self-Harm (Suicide)	33.4
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	LNE

Leading Causes of Death	Deaths
Malignant Neoplasms (Cancer) Heart Disease	27 25
Chronic Lower Respiratory Diseases	16
4. Cerebrovascular Disease	9
5. Influenza and Pneumonia	7
T6. Pneumonitis Due to Solids and Liquids	5
T6. Urinary Tract Infection, Site Not Specified	5
Percent of Deaths due to tobacco use Median age at death	12.2 84

See technical notes for more information.

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

³All mortality rates except infant mortality are ageadjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

Hamlin County

Demographic Information

Hamlin County is located in the northeastern region of the state and averages 11.6 persons per square mile.

2017 Population Information

Subject Total population White Black or African American American Indian & Alaska Native Asian Native Hawaiian & Other Pacific Islander Two or More Races	Number 5,948 5,803 25 47 23 1	Percent 100.0 97.6 0.4 0.8 0.4 0.0 0.8
Under 5 years	545	9.2
Under 18 years	1,892	31.8
65 years and over	935	15.7

Health Status Indicators 2013-2017

Natality	
Percent of Low Birth Weight Infants	4.1
Percent of Mothers Receiving	
Care in 1st Trimester	73.5
Percent of Mothers Who	
Used Tobacco While Pregnant1	13.2
Percent of Births Less Than 37 Wks. of Gestation	6.4
Average Age of Mother	27.6
Teenage Birth Rate ²	4.2
Percent White Births	96.2
Percent American Indian Births	0.5
Percent Unmarried	13.7
Percent WIC births	30.4
 Percent Breastfeeding at discharge 	88.4
 Percent Payment-Private Insurance 	76.6
Percent Payment-Medicaid	16.7
Percent C-Section	16.3

Mortality ³	
All Causes	732.4
Heart Disease	118.5
Malignant Neoplasms (Cancer)	149.9
Trachea, Bronchus, & Lung	52.1
Colon, Rectum, & Anus	13.4
Female Breast	15.4
Pancreas	10.5
Prostate	16.9
Leukemia	7.5
Chronic Lower Respiratory Diseases	31.2
 Alzheimer's Disease 	72.3
Cerebrovascular Disease	46.7
Accidents	45.6
Motor Vehicle Accidents	15.6
Diabetes	22.2
Influenza and Pneumonia	19.7
Intentional Self-Harm (Suicide)	14.1
Chronic Liver Disease and Cirrhosis	13.5
Infant Mortality	5.14

Leading Causes of Death	Total Deaths
1. Malignant Neoplasms (Cancer) 2. Heart Disease 3. Alzheimer's Disease 4. Cerebrovascular Disease 5. Accidents 6. Chronic Lower Respiratory Diseases 7. Vascular Dementia T8. Diabetes T8. Influenza and Pneumonia 10. Unspecified Dementia	61 51 37 20 15 13 12 10 10 6
Percent of Deaths due to tobacco use Median age at death	13.6 84

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Source: United States Census Bureau, 2017 Population

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

³All mortality rates except infant mortality are ageadjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

Hand County

Demographic Information

Hand County is located in central South Dakota and averages 2.4 persons per square mile.

2017 Population Information

Subject	Number	Percent
Total population	3,277	100.0
White	3,209	97.9
Black or African American	5	0.2
American Indian & Alaska Native	17	0.5
Asian	12	0.4
Native Hawaiian & Other Pacific Islander	1	0.0
Two or More Races	33	1.0
Lindar E vaara	100	F.C
Under 5 years	182	5.6
Under 18 years	698	21.3
65 years and over	824	25.1

Health Status Indicators 2013-2017

Natality	
Percent of Low Birth Weight Infants	7.0
 Percent of Mothers Receiving 	
Care in 1st Trimester	54.3
 Percent of Mothers Who 	
Used Tobacco While Pregnant ¹	8.6
Percent of Births Less Than 37 Wks. of Gestation	9.6
Average Age of Mother	28.4
Teenage Birth Rate ²	LNE
Percent White Births	97.9
Percent American Indian Births	LNE
Percent Unmarried	16.6
Percent WIC births	14.4
Percent Breastfeeding at discharge	86.6
Percent Payment-Private Insurance	84.5
Percent Payment-Medicaid	11.8
Percent C-Section	33.7

Mortality ³	
All Causes	645.3
Heart Disease	115.6
Malignant Neoplasms (Cancer)	147.3
 Trachea, Bronchus, & Lung 	21.8
Colon, Rectum, & Anus	22.1
Female Breast	LNE
Pancreas	18.5
Prostate	40.8
Leukemia	LNE
Chronic Lower Respiratory Diseases	55.5
 Alzheimer's Disease 	18.8
Cerebrovascular Disease	53.3
Accidents	48.3
 Motor Vehicle Accidents 	6.6
Diabetes	27.2
Influenza and Pneumonia	18.9
Intentional Self-Harm (Suicide)	23.8
Chronic Liver Disease and Cirrhosis	12.3
Infant Mortality	LNE

Leading Causes of Death	lotai
Leading Causes of Death	Deaths
Malignant Neoplasms (Cancer)	47
2. Heart Disease	46
Cerebrovascular Disease	23
4. Chronic Lower Respiratory Diseases	20
5. Accidents	13
6. Diabetes	10
T7. Alzheimer's Disease	9
T7. Essential (Primary) Hypertension and Hypertensive Renal Disease	9
T9. Influenza and Pneumonia	6
T9. Atherosclerosis	6
19. Atheroscierosis	O
Percent of Deaths due to tobacco use	27.4
Median age at death	84

See technical notes for more information. Source: South Dakota Department of Health, Office of Health Statistics

[•]Denotes a health status indicator which is significantly lower than the state average.

 $[\]circ \mbox{Denotes}$ a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.


[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

³All mortality rates except infant mortality are ageadjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

Hanson County

Demographic Information



Hanson County is located in southeastern South Dakota and averages 7.7 people per square mile.

2017 Population Information

Subject	Number	Percent
Total population	3,423	100.0
White	3,343	97.7
Black or African American	16	0.5
American Indian & Alaska Native	14	0.4
Asian	12	0.4
Native Hawaiian & Other Pacific Islander	1	0.0
Two or More Races	37	1.1
Under 5 years	273	8.0
Under 18 years	1,057	30.9
65 years and over	523	15.3

Health Status Indicators 2013-2017

Natality	
Percent of Low Birth Weight Infants	7.2
Percent of Mothers Receiving	
Care in 1st Trimester	69.2
 Percent of Mothers Who 	
Used Tobacco While Pregnant ¹	6.8
Percent of Births Less Than 37 Wks. of Gestation	7.2
 Average Age of Mother 	29.7
Teenage Birth Rate ²	LNE
Percent White Births	99.1
Percent American Indian Births	LNE
Percent Unmarried	12.6
Percent WIC births	11.4
Percent Breastfeeding at discharge	85.6
Percent Payment-Private Insurance	81.1
Percent Payment-Medicaid Percent C Section	6.8
Percent C-Section	24.3

Mortality ³	
o All Causes	920.3
Heart Disease	207.0
 Malignant Neoplasms (Cancer) 	279.2
Trachea, Bronchus, & Lung	60.4
 Colon, Rectum, & Anus 	75.9
Female Breast	LNE
Pancreas	24.6
Prostate	LNE
Leukemia	LNE
Chronic Lower Respiratory Diseases	LNE
Alzheimer's Disease	41.7
Cerebrovascular Disease	32.7
Accidents	53.6
Motor Vehicle Accidents	34.7
Diabetes	29.8
Influenza and Pneumonia	23.7
Intentional Self-Harm (Suicide)	18.9
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	22.52

Leading Causes of Death	Total Deaths
Malignant Neoplasms (Cancer)	40
2. Heart Disease	29
3. Accidents	8
T4. Alzheimer's Disease	5
T4. Certain Conditions Originating in the Perinatal Period	5
Percent of Deaths due to tobacco use Median age at death	13.1 76

See technical notes for more information.

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

³All mortality rates except infant mortality are ageadjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

Harding County

Demographic Information

Harding County is located in the northwestern corner of the state and averages 0.5 persons per square mile.

2017 Population Information

Subject	Number	Percent
Total population	1,242	100.0
White	1,180	95.0
Black or African American	7	0.6
American Indian & Alaska Native	26	2.1
Asian	3	0.2
Native Hawaiian & Other Pacific Islander	0	0.0
Two or More Races	26	2.1
Lindor E voore	05	6.0
Under 5 years	85	6.8
Under 18 years	284	22.9
65 years and over	209	16.8

Health Status Indicators 2013-2017

Natality	
Percent of Low Birth Weight Infants	3.4
Percent of Mothers Receiving	
Care in 1st Trimester	79.5
 Percent of Mothers Who 	
Used Tobacco While Pregnant ¹	6.7
 Percent of Births Less Than 37 Wks. of Gestation 	3.4
Average Age of Mother	28.8
Teenage Birth Rate ²	LNE
Percent White Births	92.1
Percent American Indian Births	LNE
 Percent Unmarried 	15.9
 Percent WIC births 	10.1
Percent Breastfeeding at discharge	94.2
 Percent Payment-Private Insurance 	81.6
Percent Payment-Medicaid	11.5
Percent C-Section	19.1

Motality

Mortality ³	
All Causes	448.2
Heart Disease	99.0
Malignant Neoplasms (Cancer)	91.1
Trachea, Bronchus, & Lung	LNE
Colon, Rectum, & Anus	LNE
Female Breast	LNE
Pancreas	LNE
Prostate	LNE
Leukemia	LNE
Chronic Lower Respiratory Diseases	LNE
Alzheimer's Disease	LNE
Cerebrovascular Disease	51.3
Accidents	43.3
Motor Vehicle Accidents	LNE
Diabetes	65.1
Influenza and Pneumonia	LNE
Intentional Self-Harm (Suicide)	LNE
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	LNE

Leading Causes of Death	Total Deaths
T1. Heart Disease	8
T1. Malignant Neoplasms (Cancer)	8
3. Diabetes	5
Percent of Deaths due to tobacco use	13.5
Median age at death	75

See technical notes for more information.

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

³All mortality rates except infant mortality are ageadjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

Hughes County

Demographic Information

Hughes County is located in the center of the state and averages 23.0 persons per square mile.

2017 Population Information

Number	Percent
17,666	100.0
14,847	84.0
138	0.8
2,057	11.6
136	0.8
6	0.0
482	2.7
4.000	0.0
, -	6.9
, -	24.1
2,954	16.7
	17,666 14,847 138 2,057 136 6

Health Status Indicators 2013-2017

Natality	
Percent of Low Birth Weight Infants	6.0
 Percent of Mothers Receiving 	
Care in 1st Trimester	50.1
 Percent of Mothers Who 	
Used Tobacco While Pregnant ¹	18.2
Percent of Births Less Than 37 Wks. of Gestation	9.4
Average Age of Mother	27.8
Teenage Birth Rate ²	9.7
Percent White Births	72.7
Percent American Indian Births	20.1
Percent Unmarried	41.7
Percent WIC births	34.4
Percent Breastfeeding at discharge	77.2
Percent Payment-Private Insurance	58.1
Percent Payment-Medicaid Percent C. Castian	37.6
Percent C-Section	29.2

Mortality ³	
All Causes	660.7
Heart Disease	139.2
Malignant Neoplasms (Cancer)	144.7
Trachea, Bronchus, & Lung	38.8
Colon, Rectum, & Anus	26.6
Female Breast	21.5
Pancreas	9.9
Prostate	7.2
Leukemia	2.7
Chronic Lower Respiratory Diseases	49.5
 Alzheimer's Disease 	24.6
Cerebrovascular Disease	44.9
Accidents	43.0
Motor Vehicle Accidents	14.6
Diabetes	34.0
 Influenza and Pneumonia 	29.7
Intentional Self-Harm (Suicide)	15.6
 Chronic Liver Disease and Cirrhosis 	8.0
Infant Mortality	9.27

Leading Causes of Death	I otal
Leading Causes of Death	Deaths
Malignant Neoplasms (Cancer)	160
2. Heart Disease	158
3. Chronic Lower Respiratory Diseases	56
Cerebrovascular Disease	53
5. Accidents	43
6. Diabetes	38
7. Influenza and Pneumonia	35
8. Alzheimer's Disease	30
Intentional Self-Harm (Suicide)	14
10. Unspecified Dementia	9
Percent of Deaths due to tobacco use	18.3
Median age at death	80

See technical notes for more information.

Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

³All mortality rates except infant mortality are ageadjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

Hutchinson County

Demographic Information

Hutchinson County is located in the southeastern region of the state and averages 9.0 persons per square mile.

2017 Population Information

Subject	Number	Percent
Total population	7,358	100.0
White	7,079	96.2
Black or African American	76	1.0
American Indian & Alaska Native	99	1.3
Asian	19	0.3
Native Hawaiian & Other Pacific Islander	3	0.0
Two or More Races	82	1.1
Under 5 years	611	8.3
Under 18 years	1,821	24.7
65 years and over	1,672	22.7

Health Status Indicators 2013-2017

Natality	
Percent of Low Birth Weight Infants	5.3
Percent of Mothers Receiving	
Care in 1st Trimester	68.6
 Percent of Mothers Who 	
Used Tobacco While Pregnant ¹	7.7
Percent of Births Less Than 37 Wks. of Gestation	7.9
 Average Age of Mother 	28.5
 Teenage Birth Rate² 	4.9
Percent White Births	96.6
Percent American Indian Births	0.9
Percent Unmarried	18.6
Percent WIC births	19.5
Percent Breastfeeding at discharge	83.5
Percent Payment-Private Insurance	78.3
Percent Payment-Medicaid	15.3
Percent C-Section	23.9

Mortality ³	
All Causes	698.7
Heart Disease	158.6
 Malignant Neoplasms (Cancer) 	126.2
 Trachea, Bronchus, & Lung 	24.8
Colon, Rectum, & Anus	14.4
Female Breast	13.0
Pancreas	5.3
Prostate	24.3
Leukemia	LNE
Chronic Lower Respiratory Diseases	32.2
Alzheimer's Disease	48.2
 Cerebrovascular Disease 	57.1
Accidents	69.8
Motor Vehicle Accidents	31.1
Diabetes	15.2
Influenza and Pneumonia	17.8
Intentional Self-Harm (Suicide)	LNE
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	LNE

Leading Causes of Death	Total Deaths
1. Heart Disease	141
Malignant Neoplasms (Cancer)	89
3. Alzheimer's Disease	55
Cerebrovascular Disease	52
5. Accidents	37
Chronic Lower Respiratory Diseases	23
Influenza and Pneumonia	16
T8. Diabetes	13
T8. Unspecified Dementia	13
 Essential (Primary) Hypertension and Hypertensive Renal Disease 	12
Percent of Deaths due to tobacco use Median age at death	8.5 86

See technical notes for more information.

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

³All mortality rates except infant mortality are ageadjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

Hyde County

Demographic Information

Hyde County is located in the central region of the state and averages 1.7 people per square mile.

2017 Population Information

Subject	Number	Percent
Total population	1,318	100.0
White	1,155	87.6
Black or African American	5	0.4
American Indian & Alaska Native	123	9.3
Asian	3	0.2
Native Hawaiian & Other Pacific Islander	2	0.2
Two or More Races	30	2.3
Under 5 years	75	5.7
Under 18 years	279	21.2
65 years and over	332	25.2

Health Status Indicators 2013-2017

Natality		
Percent of Low Birth Weight Infants	7.0	
Percent of Mothers Receiving		
Care in 1st Trimester	61.2	
Percent of Mothers Who		
Used Tobacco While Pregnant ¹	10.5	
Percent of Births Less Than 37 Wks. of Gestation	9.3	
Average Age of Mother	27.4	
Teenage Birth Rate ²	LNE	
Percent White Births	86.0	
Percent American Indian Births	11.6	
Percent Unmarried	29.1	
Percent WIC births	23.5	
Percent Breastfeeding at discharge	80.2	
Percent Payment-Private Insurance	74.4	
Percent Payment-Medicaid	23.3	
Percent C-Section	20.9	

Motality

Mortality ³	
All Causes	747.3
Heart Disease	220.0
Malignant Neoplasms (Cancer)	117.6
Trachea, Bronchus, & Lung	24.2
Colon, Rectum, & Anus	LNE
Female Breast	LNE
Pancreas	LNE
Prostate	LNE
Leukemia	LNE
Chronic Lower Respiratory Diseases	28.3
Alzheimer's Disease	26.0
Cerebrovascular Disease	49.8
Accidents	71.3
Motor Vehicle Accidents	42.8
Diabetes	22.3
Influenza and Pneumonia	28.5
Intentional Self-Harm (Suicide)	LNE
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	LNE

Leading Causes of Death	Total Deaths
1. Heart Disease	31
Malignant Neoplasms (Cancer)	17
Cerebrovascular Disease	9
4. Accidents	8
T5. Alzheimer's Disease	5
T5. Influenza and Pneumonia	5
Percent of Deaths due to tobacco use Median age at death	11.2 86

See technical notes for more information.

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

³All mortality rates except infant mortality are ageadjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

Jackson County

Demographic Information

Jackson County is located in western South Dakota and averages 1.6 persons per square mile.

2017 Population Information

Subject	Number	Percent
Total population	3,289	100.0
White	1,387	42.2
Black or African American	35	1.1
American Indian & Alaska Native	1,694	51.5
Asian	6	0.2
Native Hawaiian & Other Pacific Isla	ınder 1	0.0
Two or More Races	166	5.0
Under 5 years	349	10.6
Under 18 years	1,064	32.4
65 years and over	446	13.6

Health Status Indicators 2013-2017

	Natality	
Perc	ent of Low Birth Weight Infants	7.2
• Pe	ercent of Mothers Receiving	
	Care in 1st Trimester	54.3
∘ Pe	ercent of Mothers Who	
	Used Tobacco While Pregnant ¹	21.0
∘ Pe	ercent of Births Less Than 37 Wks. of Gestation	12.5
	verage Age of Mother	26.2
o Te	enage Birth Rate ²	46.8
	ent White Births	17.5
Perc	ent American Indian Births	78.4
	ercent Unmarried	74.8
	ercent WIC births	67.6
	ercent Breastfeeding at discharge	61.9
	ercent Payment-Private Insurance	14.8
	ercent Payment-Medicaid	61.9
Perc	ent C-Section	23.9

Mortality ³	
o All Causes	1,072.3
 Heart Disease 	229.5
Malignant Neoplasms (Cancer)	200.3
Trachea, Bronchus, & Lung	60.5
Colon, Rectum, & Anus	LNE
Female Breast	LNE
Pancreas	LNE
Prostate	30.6
Leukemia	LNE
Chronic Lower Respiratory Diseases	54.3
Alzheimer's Disease	LNE
Cerebrovascular Disease	26.3
 Accidents 	151.3
 Motor Vehicle Accidents 	108.7
Diabetes	49.0
Influenza and Pneumonia	14.9
Intentional Self-Harm (Suicide)	42.2
 Chronic Liver Disease and Cirrhosis 	58.2
Infant Mortality	10.28

Leading Causes of Death	Total Deaths
1. Heart Disease	41
Malignant Neoplasms (Cancer)	38
3. Accidents	23
Chronic Lower Respiratory Diseases	10
T5. Diabetes	8
T5. Chronic Liver Disease and Cirrhosis	8
7. Intentional Self-Harm (Suicide)	7
T8. Cerebrovascular Disease	5
T8. Aortic Aneurysm and Dissection	5
Percent of Deaths due to tobacco use	10.2
Median age at death	69

See technical notes for more information.

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

³All mortality rates except infant mortality are ageadjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

Jerauld County

Demographic Information

Jerauld County is located in the central region of the state and averages 3.9 persons per square mile.

2017 Population Information

Subject	Number	Percent
Total population	2,028	100.0
White	1,984	97.8
Black or African American	3	0.1
American Indian & Alaska Native	12	0.6
Asian	4	0.2
Native Hawaiian & Other Pacific Islander	2	0.1
Two or More Races	23	1.1
Under 5 years	125	6.2
Under 18 years	474	23.4
65 years and over	542	26.7

Health Status Indicators 2013-2017

Accidents

Diabetes

Infant Mortality

Median age at death

Cerebrovascular Disease

Influenza and Pneumonia

Motor Vehicle Accidents

Intentional Self-Harm (Suicide)

Chronic Liver Disease and Cirrhosis

Natality	
Percent of Low Birth Weight Infants	5.5
Percent of Mothers Receiving	
Care in 1st Trimester	77.1
Percent of Mothers Who	
Used Tobacco While Pregnant ¹	14.5
Percent of Births Less Than 37 Wks. of Gestation	8.2
 Average Age of Mother 	29.1
Teenage Birth Rate ²	LNE
Percent White Births	97.3
Percent American Indian Births	LNE
 Percent Unmarried 	25.5
 Percent WIC births 	21.1
Percent Breastfeeding at discharge	79.1
Percent Payment-Private Insurance	81.8
Percent Payment-Medicaid	12.7
Percent C-Section	22.7

Wortality	
All Causes	565.3
Heart Disease	97.8
Malignant Neoplasms (Cancer)	165.9
Trachea, Bronchus, & Lung	30.7
Colon, Rectum, & Anus	19.3
Female Breast	LNE
Pancreas	32.3
Prostate	LNE
Leukemia	13.7
 Chronic Lower Respiratory Diseases 	14.5
 Alzheimer's Disease 	85.6

34.6

59.5

47.9

LNE

19.6

INF

LNE LNE

86

Martality3

Leading Causes of Death 1. Malignant Neoplasms (Cancer) 2. Alzheimer's Disease 3. Heart Disease 4. Cerebrovascular Disease	Total Deaths 34 27 21 9
Cerebrovascular Disease Accidents	9 7
Percent of Deaths due to tobacco use	14.2

See technical notes for more information.

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

³All mortality rates except infant mortality are ageadjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

Jones County

Demographic Information

Jones County is located in western South Dakota and averages 1.0 persons per square mile.

2017 Population Information

Subject	Number	Percent
Total population	936	100.0
White	857	91.6
Black or African American	3	0.3
American Indian & Alaska Native	35	3.7
Asian	1	0.1
Native Hawaiian & Other Pacific Islander	7	0.7
Two or More Races	33	3.5
Under 5 years	53	5.7
Under 18 years	205	21.9
65 years and over	220	23.5

Health Status Indicators 2013-2017

Natality	
Percent of Low Birth Weight Infants	5.8
Percent of Mothers Receiving	
Care in 1st Trimester	55.8
Percent of Mothers Who	
Used Tobacco While Pregnant ¹	15.4
Percent of Births Less Than 37 Wks. of Gestation	11.5
Average Age of Mother	27.7
Teenage Birth Rate ²	LNE
Percent White Births	90.4
Percent American Indian Births	7.7
Percent Unmarried	28.8
Percent WIC births	26.9
Percent Breastfeeding at discharge	88.5
Percent Payment-Private Insurance	69.2
Percent Payment-Medicaid	26.9
Percent C-Section	13.5

Motality

Mortality ³	
All Causes	635.6
Heart Disease	94.4
Malignant Neoplasms (Cancer)	149.7
Trachea, Bronchus, & Lung	47.4
Colon, Rectum, & Anus	31.0
Female Breast	LNE
Pancreas	LNE
Prostate	LNE
Leukemia	LNE
Chronic Lower Respiratory Diseases	57.5
Alzheimer's Disease	LNE
Cerebrovascular Disease	LNE
Accidents	LNE
Motor Vehicle Accidents	LNE
Diabetes	LNE
Influenza and Pneumonia	LNE
Intentional Self-Harm (Suicide)	LNE
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	LNE

Leading Causes of Death	Total Deaths
Malignant Neoplasms (Cancer) Heart Disease Chronic Lower Respiratory Diseases	13 9 5
Percent of Deaths due to tobacco use Median age at death	25.0 78

See technical notes for more information.

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

³All mortality rates except infant mortality are ageadjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

Kingsbury County

Demographic Information

Kingsbury County is located in east central South Dakota and averages 6.2 persons per square mile.

2017 Population Information

Subject	Number	Percent
Total population	4,952	100.0
White	4,762	96.2
Black or African American	24	0.5
American Indian & Alaska Native	57	1.2
Asian	33	0.7
Native Hawaiian & Other Pacific Islander	1	0.0
Two or More Races	75	1.5
Under 5 years	324	6.5
Under 18 years	1.120	22.6
65 years and over	1,126	22.7

Health Status Indicators 2013-2017

Natality	
Percent of Low Birth Weight Infants	5.8
Percent of Mothers Receiving	
Care in 1st Trimester	80.1
Percent of Mothers Who	
Used Tobacco While Pregnant ¹	10.5
Percent of Births Less Than 37 Wks. of Gestation	6.2
 Average Age of Mother 	28.6
Teenage Birth Rate ²	6.5
Percent White Births	96.3
Percent American Indian Births	LNE
Percent Unmarried	21.8
Percent WIC births	22.2
Percent Breastfeeding at discharge	88.0
Percent Payment-Private Insurance	80.2
Percent Payment-Medicaid Percent C-Section	16.7
Percent C-Section	18.5

Mortality ³	
All Causes	796.5
Heart Disease	172.1
Malignant Neoplasms (Cancer)	202.0
Trachea, Bronchus, & Lung	46.3
Colon, Rectum, & Anus	19.1
Female Breast	29.7
 Pancreas 	29.4
Prostate	LNE
Leukemia	18.7
Chronic Lower Respiratory Diseases	36.3
Alzheimer's Disease	28.9
Cerebrovascular Disease	41.0
Accidents	73.3
 Motor Vehicle Accidents 	50.9
Diabetes	19.7
Influenza and Pneumonia	25.2
Intentional Self-Harm (Suicide)	34.8
Chronic Liver Disease and Cirrhosis	20.1
Infant Mortality	LNE

Leading Causes of Death	Total Deaths
1. Heart Disease	89
2. Malignant Neoplasms (Cancer)	85
T3. Accidents	21
T3. Cerebrovascular Disease	21
Chronic Lower Respiratory Diseases	18
6. Alzheimer's Disease	16
7. Influenza and Pneumonia	11
8. Diabetes	9
T9. Intentional Self-Harm (Suicide)	8
T9. Parkinson's Disease	8
Percent of Deaths due to tobacco use	11.2
Median age at death	83

See technical notes for more information.

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

³All mortality rates except infant mortality are ageadjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

Lake County

Demographic Information

Lake County is located in the east central region of the state and averages 19.9 persons per square mile.

2017 Population Information

Subject	Number	Percent
Total population	12,809	100.0
White	12,196	95.2
Black or African American	132	1.0
American Indian & Alaska Native	159	1.2
Asian	137	1.1
Native Hawaiian & Other Pacific Islander	7	0.1
Two or More Races	178	1.4
Under 5 years	688	5.4
Under 18 years	2,557	20.0
65 years and over	2.705	21.1
	_,	

Health Status Indicators 2013-2017

Natality	
Percent of Low Birth Weight Infants	5.8
Percent of Mothers Receiving	
Care in 1st Trimester	73.8
Percent of Mothers Who	
Used Tobacco While Pregnant ¹	11.2
Percent of Births Less Than 37 Wks. of Gestation	8.5
 Average Age of Mother 	28.4
 Teenage Birth Rate² 	4.8
Percent White Births	92.7
Percent American Indian Births	2.7
 Percent Unmarried 	25.9
 Percent WIC births 	25.2
Percent Breastfeeding at discharge	82.5
Percent Payment-Private Insurance	72.0
Percent Payment-Medicaid	20.3
Percent C-Section	23.8

Mortality ³	
All Causes	644.6
Heart Disease	152.0
 Malignant Neoplasms (Cancer) 	121.6
Trachea, Bronchus, & Lung	36.3
Colon, Rectum, & Anus	18.4
 Female Breast 	8.5
Pancreas	9.2
Prostate	14.7
Leukemia	6.0
Chronic Lower Respiratory Diseases	45.3
Alzheimer's Disease	36.9
 Cerebrovascular Disease 	54.2
 Accidents 	33.0
Motor Vehicle Accidents	11.7
Diabetes	30.4
Influenza and Pneumonia	10.6
Intentional Self-Harm (Suicide)	24.0
Chronic Liver Disease and Cirrhosis	10.7
Infant Mortality	9.09

Leading Causes of Death	Total Deaths
Heart Disease Malignant Neoplasms (Cancer)	149 121
3. Cerebrovascular Disease4. Chronic Lower Respiratory Diseases5. Alzheimer's Disease	50 41 34
6. Diabetes 7. Accidents	28 26 22
8. Unspecified Dementia 9. Intentional Self-Harm (Suicide) 10. Influenza and Pneumonia	12 10
Percent of Deaths due to tobacco use Median age at death	22.3 80

See technical notes for more information.

Denotes a health status indicator which is significantly lower than the state average.

 $[\]circ \mbox{Denotes}$ a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

³All mortality rates except infant mortality are ageadjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

Lawrence County

Demographic Information

Lawrence County is located along the Wyoming border and averages 30.1 persons per square mile.

2017 Population Information

Number	Percent
25,429	100.0
23,862	93.8
209	8.0
600	2.4
238	0.9
16	0.1
504	2.0
1.159	4.6
4.614	18.1
5,293	20.8
	25,429 23,862 209 600 238 16 504 1,159 4,614

Source: United States Census Bureau, 2017 Population

Estimates

Health Status Indicators 2013-2017

Natality	
 Percent of Low Birth Weight Infants 	8.9
Percent of Mothers Receiving	
Care in 1st Trimester	79.5
Percent of Mothers Who	
Used Tobacco While Pregnant ¹	16.2
 Percent of Births Less Than 37 Wks. of Gestation 	11.9
Average Age of Mother	28.1
 Teenage Birth Rate² 	4.8
Percent White Births	91.6
Percent American Indian Births	2.7
Percent Unmarried	37.6
Percent WIC births	30.1
Percent Breastfeeding at discharge	85.0
Percent Payment-Private Insurance	61.0
Percent Payment-Medicaid	31.8
Percent C-Section	20.5

Mortality ³	
All Causes	664.8
Heart Disease	126.5
Malignant Neoplasms (Cancer)	140.6
 Trachea, Bronchus, & Lung 	28.7
Colon, Rectum, & Anus	19.0
Female Breast	21.0
Pancreas	8.7
Prostate	16.6
 Leukemia 	3.1
 Chronic Lower Respiratory Diseases 	54.8
Alzheimer's Disease	29.0
Cerebrovascular Disease	27.8
Accidents	53.2
Motor Vehicle Accidents	19.0
Diabetes	22.2
Influenza and Pneumonia	17.3
Intentional Self-Harm (Suicide)	21.2

11.5

9.35

Chronic Liver Disease and Cirrhosis

Infant Mortality

Leading Causes of Death	Total Deaths
1. Malignant Neoplasms (Cancer)	247
2. Heart Disease	233
Chronic Lower Respiratory Diseases	104
4. Accidents	73
5. Alzheimer's Disease	57
Cerebrovascular Disease	52
7. Diabetes	42
8. Influenza and Pneumonia	31
Intentional Self-Harm (Suicide)	28
10. Vascular Dementia	25
Percent of Deaths due to tobacco use	21.7
Median age at death	81

See technical notes for more information.

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

[•]Denotes a health status indicator which is significantly lower than the state average.

 $[\]circ \mbox{Denotes}$ a health status indicator which is significantly higher than the state average.

³All mortality rates except infant mortality are ageadjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

Lincoln County

Demographic Information

Lincoln County is located in southeastern South Dakota and averages 77.7 persons per square mile.

2017 Population Information

Subject	Number	Percent
Total population	56,664	100.0
White	53,660	94.7
Black or African American	837	1.5
American Indian & Alaska Native	406	0.7
Asian	765	1.4
Native Hawaiian & Other Pacific Islander	31	0.1
Two or More Races	965	1.7
Under 5 years	4.206	7.4
Under 18 years	16.009	28.3
	-,	
65 years and over	7,025	12.4

Health Status Indicators 2013-2017

Natality	
Percent of Low Birth Weight Infants	5.8
 Percent of Mothers Receiving 	
Care in 1st Trimester	85.5
 Percent of Mothers Who 	
Used Tobacco While Pregnant ¹	5.4
Percent of Births Less Than 37 Wks. of Gestation	7.6
 Average Age of Mother 	29.6
 Teenage Birth Rate² 	3.4
Percent White Births	95.4
Percent American Indian Births	0.5
 Percent Unmarried 	17.0
 Percent WIC births 	8.7
Percent Breastfeeding at discharge	85.1
Percent Payment-Private Insurance	84.5
Percent Payment-Medicaid	10.5
Percent C-Section	25.2

Mortality ³	
All Causes	476.8
Heart Disease	101.0
 Malignant Neoplasms (Cancer) 	114.9
 Trachea, Bronchus, & Lung 	29.7
 Colon, Rectum, & Anus 	9.4
Female Breast	13.0
Pancreas	14.0
Prostate	20.3
 Leukemia 	3.7
 Chronic Lower Respiratory Diseases 	25.8
Alzheimer's Disease	40.7
Cerebrovascular Disease	26.7
Accidents	23.2
 Motor Vehicle Accidents 	6.8
Diabetes	10.1
 Influenza and Pneumonia 	10.0
 Intentional Self-Harm (Suicide) 	8.4
 Chronic Liver Disease and Cirrhosis 	2.9

Leading Causes of Death	Total Deaths
1. Malignant Neoplasms (Cancer)	277
2. Heart Disease	247
3. Alzheimer's Disease	94
4. Cerebrovascular Disease	65
Chronic Lower Respiratory Diseases	63
6. Accidents	61
7. Unspecified Dementia	25
T8. Diabetes	24
T8. Influenza and Pneumonia	24
10. Intentional Self-Harm (Suicide)	20
Percent of Deaths due to tobacco use	16.9
Median age at death	80

6.37

Infant Mortality

See technical notes for more information.

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

[•]Denotes a health status indicator which is significantly lower than the state average.

 $[\]circ \mbox{Denotes}$ a health status indicator which is significantly higher than the state average.

³All mortality rates except infant mortality are ageadjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

Lyman County

Demographic Information

Lyman County is located in central South Dakota and averages 2.3 persons per square mile.

2017 Population Information

Subject	Number	Percent
Total population	3,904	100.0
White	2,201	56.4
Black or African American	22	0.6
American Indian & Alaska Native	1,532	39.2
Asian	12	0.3
Native Hawaiian & Other Pacific Islander	2	0.1
Two or More Races	135	3.5
Under 5 years	348	8.9
Under 18 years	1,148	29.4
65 years and over	610	15.6

Health Status Indicators 2013-2017

Natality	
Percent of Low Birth Weight Infants	8.4
Percent of Mothers Receiving	
Care in 1st Trimester	45.9
 Percent of Mothers Who 	
Used Tobacco While Pregnant ¹	25.2
Percent of Births Less Than 37 Wks. of Gestation	11.8
 Average Age of Mother 	26.5
 Teenage Birth Rate² 	39.0
Percent White Births	38.7
Percent American Indian Births	56.0
 Percent Unmarried 	62.7
 Percent WIC births 	59.1
 Percent Breastfeeding at discharge 	62.5
 Percent Payment-Private Insurance 	34.9
 Percent Payment-Medicaid 	57.8
Percent C-Section	29.8

Mortality ³	
All Causes	822.5
Heart Disease	159.1
Malignant Neoplasms (Cancer)	160.9
Trachea, Bronchus, & Lung	42.9
Colon, Rectum, & Anus	16.5
Female Breast	40.9
Pancreas	LNE
Prostate	55.4
Leukemia	LNE
Chronic Lower Respiratory Diseases	40.2
Alzheimer's Disease	42.3
Cerebrovascular Disease	42.5
Accidents	83.6
Motor Vehicle Accidents	33.7
Diabetes	39.3
Influenza and Pneumonia	17.8
 Intentional Self-Harm (Suicide) 	68.0
Chronic Liver Disease and Cirrhosis	30.5
Infant Mortality	LNE

Leading Causes of Death	l otal Deaths
1. Malignant Neoplasms (Cancer)	36
2. Heart Disease	33
3. Accidents	14
4. Intentional Self-Harm (Suicide)	11
T5. Chronic Lower Respiratory Diseases	9
T5. Cerebrovascular Disease	9
T5. Diabetes	9
8. Alzheimer's Disease	8
Percent of Deaths due to tobacco use	24.6
Median age at death	71

See technical notes for more information.

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

³All mortality rates except infant mortality are ageadjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

McCook County

Demographic Information

McCook County is located in eastern South Dakota and averages 9.8 persons per square mile.

2017 Population Information

Number	Percent
5,499	100.0
5,318	96.7
32	0.6
65	1.2
13	0.2
4	0.1
67	1.2
425	7.7
1,516	27.6
1,019	18.5
	5,499 5,318 32 65 13 4 67 425 1,516

Source: United States Census Bureau, 2017 Population

Estimates

Health Status Indicators 2013-2017

Natality	
Percent of Low Birth Weight Infants	4.8
Percent of Mothers Receiving	
Care in 1st Trimester	75.0
 Percent of Mothers Who 	
Used Tobacco While Pregnant ¹	8.9
Percent of Births Less Than 37 Wks. of Gestation	7.4
 Average Age of Mother 	28.7
Teenage Birth Rate ²	6.8
Percent White Births	95.2
Percent American Indian Births	0.8
 Percent Unmarried 	23.1
Percent WIC births	26.7
Percent Breastfeeding at discharge	81.0
Percent Payment-Private Insurance	74.8
Percent Payment-Medicaid	17.6
Percent C-Section	23.9

Mortality ³	
All Causes	860.9
 Heart Disease 	197.1
 Malignant Neoplasms (Cancer) 	230.9
Trachea, Bronchus, & Lung	50.3
Colon, Rectum, & Anus	31.9
Female Breast	27.3
Pancreas	16.2
Prostate	47.0
Leukemia	LNE
 Chronic Lower Respiratory Diseases 	13.9
 Alzheimer's Disease 	63.9
Cerebrovascular Disease	51.7
Accidents	47.0
Motor Vehicle Accidents	20.3
Diabetes	35.0
Influenza and Pneumonia	12.2
Intentional Self-Harm (Suicide)	15.5
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	LNE

Leading Causes of Death	Total Deaths
1. Malignant Neoplasms (Cancer)	94
2. Heart Disease	91
3. Alzheimer's Disease	34
Cerebrovascular Disease	25
5. Accidents	15
6. Diabetes	14
7. Septicemia	9
8. Influenza and Pneumonia	7
9. Chronic Lower Respiratory Diseases	6
10. Parkinson's Disease	5
Percent of Deaths due to tobacco use Median age at death	11.4 82

177

See technical notes for more information.

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.


[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

³All mortality rates except infant mortality are ageadjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

McPherson County

Demographic Information



McPherson County is located in the north central region of the state and averages 2.2 persons per square mile.

2017 Population Information

Subject	Number	Percent
Total population	2,426	100.0
White	2,368	97.6
Black or African American	15	0.6
American Indian & Alaska Native	6	0.2
Asian	12	0.5
Native Hawaiian & Other Pacific Islander	2	0.1
Two or More Races	23	0.9
Under 5 years	172	7.1
Under 18 years	587	24.2
65 years and over	679	28.0

Health Status Indicators 2013-2017

Natality	
Percent of Low Birth Weight Infants	7.4
Percent of Mothers Receiving	
Care in 1st Trimester	53.3
Percent of Mothers Who	
Used Tobacco While Pregnant ¹	9.1
Percent of Births Less Than 37 Wks. of Gestation	9.1
 Average Age of Mother 	28.9
Teenage Birth Rate ²	LNE
Percent White Births	98.3
Percent American Indian Births	LNE
 Percent Unmarried 	18.2
Percent WIC births	24.2
Percent Breastfeeding at discharge	83.5
Percent Payment-Private Insurance	73.3
Percent Payment-Medicaid	20.0
Percent C-Section	25.6

Mortality ³	
All Causes	717.6
 Heart Disease 	232.8
Malignant Neoplasms (Cancer)	146.2
Trachea, Bronchus, & Lung	30.3
Colon, Rectum, & Anus	12.4
Female Breast	LNE
Pancreas	LNE
Prostate	LNE
Leukemia	LNE
 Chronic Lower Respiratory Diseases 	22.7
Alzheimer's Disease	29.2
Cerebrovascular Disease	38.4
Accidents	46.7
Motor Vehicle Accidents	18.7
Diabetes	32.7
Influenza and Pneumonia	LNE
Intentional Self-Harm (Suicide)	52.1
Chronic Liver Disease and Cirrhosis	19.1
Infant Mortality	LNE

Leading Causes of Death	Deaths
1. Heart Disease	66
Malignant Neoplasms (Cancer)	36
Cerebrovascular Disease	15
4. Accidents	13
Alzheimer's Disease	12
Chronic Lower Respiratory Diseases	8
7. Diabetes	7
Percent of Deaths due to tobacco use	9.9
Median age at death	85

See technical notes for more information.

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

 $[\]bullet \mbox{Denotes}$ a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

³All mortality rates except infant mortality are ageadjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

Marshall County

Demographic Information

Marshall County is located in the northeastern part of the state and averages 5.6 persons per square mile.

2017 Population Information

Subject	Number	Percent
Total population	4,804	100.0
White	4,232	88.1
Black or African American	35	0.7
American Indian & Alaska Native	437	9.1
Asian	11	0.2
Native Hawaiian & Other Pacific Islander	0	0.0
Two or More Races	89	1.9
Under 5 years	361	7.5
Under 18 years	1,111	23.1
65 years and over	997	20.8

Health Status Indicators 2013-2017

Natality	
Percent of Low Birth Weight Infants	5.4
 Percent of Mothers Receiving 	
Care in 1st Trimester	56.4
Percent of Mothers Who	
Used Tobacco While Pregnant ¹	14.0
Percent of Births Less Than 37 Wks. of Gestation	6.8
 Average Age of Mother 	28.7
Teenage Birth Rate ²	LNE
Percent White Births	79.2
Percent American Indian Births	10.8
 Percent Unmarried 	28.2
 Percent WIC births 	24.8
Percent Breastfeeding at discharge	84.3
Percent Payment-Private Insurance	70.5
Percent Payment-Medicaid	20.6
Percent C-Section	24.5

Mortality ³		
All Causes	655.4	
Heart Disease	134.6	
Malignant Neoplasms (Cancer)	135.4	
Trachea, Bronchus, & Lung	35.8	
Colon, Rectum, & Anus	7.9	
Female Breast	23.5	
Pancreas	20.0	
Prostate	25.2	
Leukemia	7.1	
Chronic Lower Respiratory Diseases	50.5	
Alzheimer's Disease	26.7	
 Cerebrovascular Disease 	68.9	
Accidents	55.3	
Motor Vehicle Accidents	18.6	
Diabetes	14.1	
Influenza and Pneumonia	10.0	
Intentional Self-Harm (suicide)	21.7	
Chronic Liver Disease and Cirrhosis	7.3	
Infant Mortality	8.55	

Leading Causes of Death	Total Deaths
1. Heart Disease	53
Malignant Neoplasms (Cancer)	48
Cerebrovascular Disease	25
4. Chronic Lower Respiratory Diseases	20
5. Accidents	15
6. Alzheimer's Disease	11
7. Intentional Self-Harm (Suicide)	6
Percent of Deaths due to tobacco use	18.6
Median age at death	80

See technical notes for more information.

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

³All mortality rates except infant mortality are ageadjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

Meade County

Demographic Information

Meade County is located in west central South Dakota and averages 7.3 persons per square mile.

2017 Population Information

Subject	Number	Percent
Total population	28,018	100.0
White	25,432	90.8
Black or African American	583	2.1
American Indian & Alaska Native	781	2.8
Asian	348	1.2
Native Hawaiian & Other Pacific Islander	34	0.1
Two or More Races	840	3.0
Under 5 years	1,599	5.7
Under 18 years	6,509	23.2
65 years and over	4,107	14.7

Source: United States Census Bureau, 2017 Population Estimates

Health Status Indicators 2013-2017

Natality	
Percent of Low Birth Weight Infants	7.7
Percent of Mothers Receiving	
Care in 1st Trimester	78.4
Percent of Mothers Who	
Used Tobacco While Pregnant ¹	15.0
Percent of Births Less Than 37 Wks. of Gestation	9.7
Average Age of Mother	28.1
 Teenage Birth Rate² 	5.5
Percent White Births	90.3
Percent American Indian Births	3.8
 Percent Unmarried 	28.5
Percent WIC births	28.9
 Percent Breastfeeding at discharge 	87.8
Percent Payment-Private Insurance	57.5
Percent Payment-Medicaid	26.1
Percent C-Section	21.0

Mortality ³		
All Causes	695.3	
Heart Disease	142.7	
Malignant Neoplasms (Cancer)	161.1	
Trachea, Bronchus, & Lung	47.5	
Colon, Rectum, & Anus	13.6	
Female Breast	14.4	
Pancreas	11.9	
Prostate	25.8	
Leukemia	5.8	
 Chronic Lower Respiratory Diseases 	64.5	
 Alzheimer's Disease 	20.7	
Cerebrovascular Disease	32.1	
Accidents	39.9	
Motor Vehicle Accidents	14.2	
 Diabetes 	15.3	
Influenza and Pneumonia	17.8	
Intentional Self-Harm (suicide)	24.5	
Chronic Liver Disease and Cirrhosis	11.1	
Infant Mortality	3.40	

Leading Causes of Death	Total
Leading Causes of Death	Deaths
Malignant Neoplasms (Cancer)	240
2. Heart Disease	203
3. Chronic Lower Respiratory Diseases	92
4. Accidents	54
Cerebrovascular Disease	44
Intentional Self-Harm (Suicide)	33
7. Alzheimer's Disease	28
Influenza and Pneumonia	25
9. Diabetes	24
10. Unspecified Dementia	23
Percent of Deaths due to tobacco use	19.6
Median age at death	77

See technical notes for more information.

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

³All mortality rates except infant mortality are ageadjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

Mellette County

Demographic Information

Mellette County is located in the south central region of the state and averages 1.6 persons per square mile.

2017 Population Information

Subject	Number	Percent
Total population	2,088	100.0
White	840	40.2
Black or African American	6	0.3
American Indian & Alaska Native	1,123	53.8
Asian	5	0.2
Native Hawaiian & Other Pacific Islander	0	0.0
Two or More Races	114	5.5
Under 5 years	197	9.4
Under 18 years	631	30.2
65 years and over	328	15.7

Source: United States Census Bureau, 2017 Population Estimates

Natality	
Percent of Low Birth Weight Infants	5.0
 Percent of Mothers Receiving 	
Care in 1st Trimester	48.2
Percent of Mothers Who	
Used Tobacco While Pregnant ¹	19.3
Percent of Births Less Than 37 Wks. of Gestation	7.5
 Average Age of Mother 	25.6
 ○ Teenage Birth Rate² 	37.8
Percent White Births	19.9
Percent American Indian Births	66.7
 Percent Unmarried 	72.1
 Percent WIC births 	61.1
Percent Breastfeeding at discharge	67.8
 Percent Payment-Private Insurance 	15.5
 Percent Payment-Medicaid 	64.5
o Percent C-Section	40.8

Health Status Indicators 2013-2017

Mortality ³	
o All Causes	1,161.9
Heart Disease	203.8
Malignant Neoplasms (Cancer)	148.8
Trachea, Bronchus, & Lung	30.2
Colon, Rectum, & Anus	LNE
Female Breast	LNE
Pancreas	LNE
Prostate	LNE
Leukemia	LNE
Chronic Lower Respiratory Diseases	98.9
Alzheimer's Disease	47.4
Cerebrovascular Disease	31.6
Accidents	101.0
Motor Vehicle Accidents	56.6
Diabetes	66.5
Influenza and Pneumonia	22.5
Intentional Self-Harm (suicide)	34.4
 Chronic Liver Disease and Cirrhosis 	146.3
Infant Mortality	LNE

Leading Causes of Death	l otal Deaths
1. Heart Disease	26
Malignant Neoplasms (Cancer)	19
Chronic Lower Respiratory Diseases	12
Chronic Liver Disease and Cirrhosis	11
5. Accidents	10
6. Diabetes	7
7. Alzheimer's Disease	6
Percent of Deaths due to tobacco use Median age at death	17.7 75

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

¹Data for mothers who used tobacco are self-reported.

than the state average.

than the state average.

•Denotes a health status indicator which is significantly lower

oDenotes a health status indicator which is significantly higher

²Teenage Birth rate is live births per 1,000 females age 15-17.

[•]Denotes a health status indicator which is significantly lower than the state average.


oDenotes a health status indicator which is significantly higher than the state average.

³All mortality rates except infant mortality are ageadjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

Miner County

Health Status Indicators 2013-2017

Demographic Information



Miner County is a located in the east central region of the state and averages 4.2 persons per square mile.

2017 Population Information

Subject	Number	Percent
Total population	2,228	100.0
White	2,147	96.4
Black or African American	22	1.0
American Indian & Alaska Native	15	0.7
Asian	10	0.4
Native Hawaiian & Other Pacific Islander	0	0.0
Two or More Races	34	1.5
Under 5 years	143	6.4
Under 18 years	547	24.6
65 years and over	452	20.3

Source: United States Census Bureau, 2017 Population Estimates

Natality	
Percent of Low Birth Weight Infants	7.7
Percent of Mothers Receiving	
Care in 1st Trimester	65.6
Percent of Mothers Who	
Used Tobacco While Pregnant1	7.7
Percent of Births Less Than 37 Wks. of Gestation	6.9
Average Age of Mother	29.4
Teenage Birth Rate ²	LNE
Percent White Births	95.4
Percent American Indian Births	LNE
Percent Unmarried	16.2
Percent WIC births	14.6
Percent Breastfeeding at discharge	83.1
Percent Payment-Private Insurance	77.5
Percent Payment-Medicaid	18.6
Percent C-Section	30.8

Mortality ³			
All Causes	820.7		
Heart Disease	183.8		
Malignant Neoplasms (Cancer)	197.9		
Trachea, Bronchus, & Lung	62.4		
Colon, Rectum, & Anus	LNE		
Female Breast	LNE		
Pancreas	17.9		
Prostate	LNE		
Leukemia	13.5		
Chronic Lower Respiratory Diseases	34.0		
Alzheimer's Disease	38.9		
Cerebrovascular Disease	44.0		
Accidents	73.5		
Motor Vehicle Accidents	LNE		
Diabetes	46.9		
Influenza and Pneumonia	LNE		
Intentional Self-Harm (Suicide)	LNE		
Chronic Liver Disease and Cirrhosis	LNE		
Infant Mortality	LNE		

Leading Causes of Death	Deaths
1. Heart Disease	39
Malignant Neoplasms (Cancer)	38
3. Accidents	12
Cerebrovascular Disease	11
T5. Alzheimer's Disease	10
T5. Diabetes	10
Chronic Lower Respiratory Diseases	7
8. Unspecified Dementia	5
Percent of Deaths due to tobacco use	18.9
Median age at death	82

[•]Denotes a health status indicator which is significantly lower than the state average.

See technical notes for more information.

oDenotes a health status indicator which is significantly higher than the state average.

³All mortality rates except infant mortality are ageadjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

[•]Denotes a health status indicator which is significantly lower than the state average.


 $[\]circ \mbox{Denotes}$ a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Minnehaha County

Demographic Information



Minnehaha County is located in southeastern South Dakota and averages 210.0 persons per square mile.

2017 Population Information

Subject	Number	Percent
Total population	188,616	100.0
White	163,780	86.8
Black or African American	10,644	5.6
American Indian & Alaska Native	5,256	2.8
Asian	4,142	2.2
Native Hawaiian & Other Pacific Islander	230	0.1
Two or More Races	4,564	2.4
Under 5 years	14,635	7.8
Under 18 years	47,882	25.4
65 years and over	24,161	12.8

Matality	

Health Status Indicators 2013-2017

ı	ivatanty	
I	Percent of Low Birth Weight Infants	6.6
l	 Percent of Mothers Receiving 	
l	Care in 1st Trimester	77.0
l	 Percent of Mothers Who 	
l	Used Tobacco While Pregnant ¹	10.8
l	Percent of Births Less Than 37 Wks. of Gestation	8.0
l	Average Age of Mother	28.4
l	Teenage Birth Rate ²	9.8
l	Percent White Births	77.7
l	Percent American Indian Births	4.5
l	Percent Unmarried Percent WIChirtha	33.6
l	Percent WIC births	25.8
l	Percent Breastfeeding at discharge O Percent Payment-Private Insurance	80.9 65.8
l	Percent Payment-Private insurance Percent Payment-Medicaid	29.0
I	Percent C-Section	25.1
ı	1 Cicciii O Occilori	20.1

Mortality ³				
All Causes	713.0			
Heart Disease	150.5			
Malignant Neoplasms (Cancer)	165.9			
Trachea, Bronchus, & Lung	42.9			
Colon, Rectum, & Anus	15.7			
Female Breast	21.4			
Pancreas	11.5			
Prostate	20.0			
Leukemia	7.0			
Chronic Lower Respiratory Diseases	43.8			
 Alzheimer's Disease 	44.7			
Cerebrovascular Disease	37.2			
Accidents	44.9			
 Motor Vehicle Accidents 	9.2			
 Diabetes 	15.5			

13.2

18.9

10.7

5.03

Influenza and Pneumonia

Infant Mortality

Intentional Self-Harm (Suicide)

• Chronic Liver Disease and Cirrhosis

Leading Causes of Death	Total Deaths
1. Malignant Neoplasms (Cancer)	1,566
2. Heart Disease	1,456
3. Alzheimer's Disease	434
4. Accidents	416
5. Chronic Lower Respiratory Diseases	409
Cerebrovascular Disease	353
7. Intentional Self-Harm (Suicide)	171
8. Diabetes	144
Influenza and Pneumonia	132
10. Chronic Liver Disease and Cirrhosis	101
Percent of Deaths due to tobacco use Median age at death	17.5 78
iviculari aye at ucatri	70

See technical notes for more information.

[•]Denotes a health status indicator which is significantly lower than the state average.

 $[\]circ \mbox{Denotes}$ a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.


[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

³All mortality rates except infant mortality are ageadjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

Moody County

Demographic Information



Moody County is located on the Minnesota border and averages 12.5 persons per square mile.

2017 Population Information

Subject	Number	Percent
Total population	6,579	100.0
White	5,189	78.9
Black or African American	131	2.0
American Indian & Alaska Native	890	13.5
Asian	151	2.3
Native Hawaiian & Other Pacific Islander	0	0.0
Two or More Races	218	3.3
Under 5 years	499	7.6
Under 18 years	1,722	26.2
65 years and over	1,208	18.4

Source: United States Census Bureau, 2017 Population Estimates

Natality	
Percent of Low Birth Weight Infants	5.7
Percent of Mothers Receiving	
Care in 1st Trimester	73.8
Percent of Mothers Who	
Used Tobacco While Pregnant ¹	15.1
Percent of Births Less Than 37 Wks. of Gestation	9.2
Average Age of Mother	28.2
Teenage Birth Rate ²	12.2
Percent White Births	68.9
Percent American Indian Births	21.9
Percent Unmarried	41.1
Percent WIC births	35.2
Percent Breastfeeding at discharge	78.9
Percent Payment-Private Insurance	60.0
Percent Payment-Medicaid	31.1
Percent C-Section	24.5

	Mortality ³	
5.7	All Causes	633.1
	Heart Disease	154.1
73.8	Malignant Neoplasms (Cancer)	136.8
	 Trachea, Bronchus, & Lung 	22.7
15.1	Colon, Rectum, & Anus	20.9
9.2	Female Breast	LNE
28.2	Pancreas	13.5
12.2	Prostate	28.1
68.9	Leukemia	LNE
21.9	 Chronic Lower Respiratory Diseases 	21.4
41.1	Alzheimer's Disease	46.2
35.2	Cerebrovascular Disease	31.8
78.9	Accidents	36.9
60.0	Motor Vehicle Accidents	11.1
31.1	Diabetes	27.1
24.5	Influenza and Pneumonia	LNE
	Intentional Self-Harm (Suicide)	22.5
	Chronic Liver Disease and Cirrhosis	8.1
	Infant Mortality	6.56

Health Status Indicators 2013-2017

Leading Causes of Death	Total
Ecading Gauses of Death	Deaths
Heart Disease	69
Malignant Neoplasms (Cancer)	56
3. Alzheimer's Disease	23
Cerebrovascular Disease	15
5. Accidents	13
6. Diabetes	12
7. Vascular Dementia	11
8. Chronic Lower Respiratory Diseases	10
Peripheral Vascular Disease, Unspecified	8
10. Intentional Self-Harm (Suicide)	7
Percent of Deaths due to tobacco use Median age at death	17.8 79

Total

- •Denotes a health status indicator which is significantly lower than the state average.
- $\circ \mbox{Denotes}$ a health status indicator which is significantly higher than the state average.
- ³All mortality rates except infant mortality are ageadjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.


^{Denotes a health status indicator which is significantly lower} than the state average.
Denotes a health status indicator which is significantly higher

than the state average.

¹Data for mothers who used tobacco are self-reported. ²Teenage Birth rate is live births per 1,000 females age 15-17.

Oglala Lakota County

Demographic Information



Oglala Lakota County (formerly known as Shannon County) is located in the southwestern part of the state, along the Nebraska border and averages 6.5 persons per square mile.

2017 Population Information

Subject	Number	Percent
Total population	14,354	100.0
White	746	5.2
Black or African American	52	0.4
American Indian & Alaska Native	13,291	92.6
Asian	19	0.1
Native Hawaiian & Other Pacific Islander	8	0.1
Two or More Races	238	1.7
Under 5 years	1,519	10.6
Under 18 years	5,391	37.6
65 years and over	1,007	7.0

Source: United States Census Bureau, 2017 Population

Estimates

Natality	
Percent of Low Birth Weight Infants	7.5
Percent of Mothers Receiving	
Care in 1st Trimester	59.3
 Percent of Mothers Who 	
Used Tobacco While Pregnant ¹	17.3
 Percent of Births Less Than 37 Wks. of Gestation 	11.5
Average Age of Mother	25.3
 ○ Teenage Birth Rate² 	38.6
Percent White Births	1.4
Percent American Indian Births	96.6
 Percent Unmarried 	88.1
Percent WIC births	71.6
 Percent Breastfeeding at discharge 	56.8

Percent Payment-Private Insurance

Percent Payment-Medicaid

Percent C-Section

Health Status Indicators 2013-2017

3.8

62.5

23.1

Mortality ³	
o All Causes	1,447.1
Heart Disease	187.1
 Malignant Neoplasms (Cancer) 	213.2
Trachea, Bronchus, & Lung	59.3
Colon, Rectum, & Anus	24.2
Female Breast	12.8
Pancreas	13.9
Prostate	28.8
Leukemia	LNE
Chronic Lower Respiratory Diseases	49.1
 Alzheimer's Disease 	14.7
Cerebrovascular Disease	44.0
 Accidents 	169.6
 Motor Vehicle Accidents 	79.7
 Diabetes 	139.3
Influenza and Pneumonia	36.0
 Intentional Self-Harm (Suicide) 	49.1
 Chronic Liver Disease and Cirrhosis 	164.2
 Infant Mortality 	15.57

Leading Causes of Death	rotai
Louding Gudoco of Doutin	Deaths
1. Accidents	105
Malignant Neoplasms (Cancer)	92
3. Chronic Liver Disease and Cirrhosis	86
4. Heart Disease	79
5. Diabetes	64
Intentional Self-Harm (Suicide)	39
7. Cerebrovascular Disease	20
T8. Chronic Lower Respiratory Diseases	19
T8. Septicemia	19
10. Assault (Homicide)	16
Percent of Deaths due to tobacco use	11.1
Median age at death	55

See technical notes for more information.

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

[•]Denotes a health status indicator which is significantly lower than the state average.


oDenotes a health status indicator which is significantly higher than the state average.

³All mortality rates except infant mortality are ageadjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

Pennington County

Health Status Indicators 2013-2017

Demographic Information



Pennington County is located on the Wyoming border and averages 36.4 persons per square mile.

2017 Population Information

Subject	Number	Percent
Total population	110,141	100.0
White	91,949	83.5
Black or African American	1,477	1.3
American Indian & Alaska Native	11,105	10.1
Asian	1,299	1.2
Native Hawaiian & Other Pacific Islander	108	0.1
Two or More Races	4,203	3.8
Under 5 years	7,355	6.7
Under 18 years	25,591	23.2
65 years and over	19,101	17.3

Source: United States Census Bureau, 2017 Population Estimates

Natality	
Percent of Low Birth Weight Infants	6.9
Percent of Mothers Receiving	
Care in 1st Trimester	73.6
 Percent of Mothers Who 	
Used Tobacco While Pregnant ¹	16.1
Percent of Births Less Than 37 Wks. of Gestation	9.5
 Average Age of Mother 	27.4
Teenage Birth Rate ²	13.5
Percent White Births	71.0
Percent American Indian Births	18.1
 Percent Unmarried 	42.9
 Percent WIC births 	37.1
 Percent Breastfeeding at discharge 	84.2
Percent Payment-Private Insurance	42.3
Percent Payment-Medicaid	37.6
 Percent C-Section 	20.0

Mortality ³		
All Causes	653.3	
Heart Disease	149.9	
Malignant Neoplasms (Cancer)	152.3	
Trachea, Bronchus, & Lung	40.8	
 Colon, Rectum, & Anus 	11.2	
Female Breast	13.6	
Pancreas	11.2	
Prostate	17.4	
Leukemia	9.6	
 Chronic Lower Respiratory Diseases 	31.4	
 Alzheimer's Disease 	27.8	
 Cerebrovascular Disease 	26.4	
Accidents	46.1	
Motor Vehicle Accidents	13.8	
 Diabetes 	13.4	
 Influenza and Pneumonia 	12.1	
Intentional Self-Harm (Suicide)	20.4	
Chronic Liver Disease and Cirrhosis	15.8	
Infant Mortality	6.18	

Leading Causes of Death	Total
	Deaths
Heart Disease	1,035
Malignant Neoplasms (Cancer)	1,018
3. Accidents	268
4. Chronic Lower Respiratory Diseases	213
5. Alzheimer's Disease	198
Cerebrovascular Disease	182
7. Intentional Self-Harm (Suicide)	106
Unspecified Dementia	102
Chronic Liver Disease and Cirrhosis	91
10. Diabetes	88
Percent of Deaths due to tobacco use	17.4
Median age at death	77

See technical notes for more information.

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

[•]Denotes a health status indicator which is significantly lower than the state average.


oDenotes a health status indicator which is significantly higher than the state average.

³All mortality rates except infant mortality are ageadjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

Perkins County

Health Status Indicators 2013-2017

Demographic Information



Perkins County is located in northwestern South Dakota and averages 1.0 persons per square mile.

2017 Population Information

Subject	Number	Percent
Total population	2,974	100.0
White	2,843	95.6
Black or African American	15	0.5
American Indian & Alaska Native	65	2.2
Asian	8	0.3
Native Hawaiian & Other Pacific Islander	2	0.1
Two or More Races	41	1.4
Under 5 years	182	6.1
Under 18 years	637	21.4
65 years and over	716	24.1

Source: United States Census Bureau, 2017 Population

Estimates

Natality	
Percent of Low Birth Weight Infants	5.9
Percent of Mothers Receiving	
Care in 1st Trimester	72.4
Percent of Mothers Who	
Used Tobacco While Pregnant ¹	13.0
Percent of Births Less Than 37 Wks. of Gestation	6.5
 Average Age of Mother 	28.8
Teenage Birth Rate ²	LNE
Percent White Births	94.1
Percent American Indian Births	2.7
Percent Unmarried	23.2
Percent WIC births	36.7
Percent Breastfeeding at discharge	86.2
Percent Payment-Private Insurance	60.2
Percent Payment-Medicaid	25.0
Percent C-Section	25.9

Mortality ³	
All Causes	750.3
Heart Disease	144.1
Malignant Neoplasms (Cancer)	154.3
 Trachea, Bronchus, & Lung 	18.1
Colon, Rectum, & Anus	18.4
Female Breast	48.4
Pancreas	LNE
Prostate	LNE
Leukemia	12.8
Chronic Lower Respiratory Diseases	59.4
Alzheimer's Disease	40.4
Cerebrovascular Disease	32.9
Accidents	36.8
Motor Vehicle Accidents	26.9
Diabetes	22.9
Influenza and Pneumonia	18.5
Intentional Self-Harm (Suicide)	LNE
Chronic Liver Disease and Cirrhosis	15.2
Infant Mortality	LNE

Leading Causes of Death	Deaths
1. Heart Disease	44
Malignant Neoplasms (Cancer)	43
3. Chronic Lower Respiratory Diseases	20
4. Alzheimer's Disease	16
Cerebrovascular Disease	11
6. Diabetes	7
7. Accidents	6
8. Aortic Aneurysm and Dissection	5
Percent of Deaths due to tobacco use	20.6
Median age at death	82

See technical notes for more information.

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

[•]Denotes a health status indicator which is significantly lower than the state average.

 $[\]circ \mbox{Denotes}$ a health status indicator which is significantly higher than the state average.

³All mortality rates except infant mortality are ageadjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

Potter County

Health Status Indicators 2013-2017

Demographic Information

Potter County is located in north central South Dakota and averages 2.7 persons per square mile.

2017 Population Information

Subject	Number	Percent
Total population	2,231	100.0
White	2,109	94.5
Black or African American	10	0.4
American Indian & Alaska Native	49	2.2
Asian	23	1.0
Native Hawaiian & Other Pacific Islander	1	0.0
Two or More Races	39	1.7
Under 5 years	113	5.1
Under 18 years	475	21.3
65 years and over	639	28.6

· · · ·	
Natality	
Percent of Low Birth Weight Infants	6.3
 Percent of Mothers Receiving 	
Care in 1st Trimester	56.8
Percent of Mothers Who	
Used Tobacco While Pregnant ¹	9.9
Percent of Births Less Than 37 Wks. of Gestation	8.1
Average Age of Mother	28.6
Teenage Birth Rate ²	LNE
Percent White Births	90.1
Percent American Indian Births	9.0
 Percent Unmarried 	21.6
 Percent WIC births 	19.8
Percent Breastfeeding at discharge	87.4
 Percent Payment-Private Insurance 	75.7
 Percent Payment-Medicaid 	20.7
Percent C-Section	22.5

Mortality ³	
All Causes	701.4
Heart Disease	116.5
Malignant Neoplasms (Cancer)	118.1
Trachea, Bronchus, & Lung	40.4
Colon, Rectum, & Anus	LNE
Female Breast	LNE
Pancreas	11.2
Prostate	33.4
Leukemia	LNE
Chronic Lower Respiratory Diseases	41.8
Alzheimer's Disease	42.3
 Cerebrovascular Disease 	14.0
Accidents	82.8
Motor Vehicle Accidents	35.1
Diabetes	39.6
Influenza and Pneumonia	20.7
Intentional Self-Harm (Suicide)	31.6
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	LNE
	Total

Leading Causes of Death	Deaths
1. Heart Disease	35
Malignant Neoplasms (Cancer)	33
Alzheimer's Disease	16
T4. Accidents	12
T4. Chronic Lower Respiratory Diseases	12
6. Diabetes	10
7. Septicemia	7
T8. Cerebrovascular Disease	5
T8. Influenza and Pneumonia	5
T8. Essential (Primary) Hypertension and Hypertensive Renal Disease	5
Percent of Deaths due to tobacco use Median age at death	15.9 85

- •Denotes a health status indicator which is significantly lower than the state average.
- $\circ\mbox{Denotes}$ a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

•Denotes a health status indicator which is significantly lower than the state average.

 $\circ \mbox{Denotes}$ a health status indicator which is significantly higher than the state average.

³All mortality rates except infant mortality are ageadjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Roberts County

Demographic Information

Roberts County is located in the extreme northeastern corner of the state and averages 9.2 persons per square mile.

2017 Population Information

Subject	Number	Percent
Total population	10,278	100.0
White	5,950	57.9
Black or African American	76	0.7
American Indian & Alaska Native	3,888	37.8
Asian	24	0.2
Native Hawaiian & Other Pacific Islander	1	0.0
Two or More Races	339	3.3
Under 5 years	906	8.8
Under 18 years	2,896	28.2
65 years and over	1,976	19.2

Source: United States Census Bureau, 2017 Population

Estimates

Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Natality	
Percent of Low Birth Weight Infants	3.9
Percent of Mothers Receiving	
Care in 1st Trimester	59.1
 Percent of Mothers Who 	
Used Tobacco While Pregnant ¹	29.2
Percent of Births Less Than 37 Wks. of Gestation	10.3
 Average Age of Mother 	26.3
 Teenage Birth Rate² 	28.9
Percent White Births	30.1
Percent American Indian Births	58.6
 Percent Unmarried 	63.7
Percent WIC births	57.8
Percent Breastfeeding at discharge	68.9
Percent Payment-Private Insurance	33.3
Percent Payment-Medicaid	57.5
Percent C-Section	25.0

Health Status Indicators 2013-2017

Mortality ³	
o All Causes	815.3
Heart Disease	144.3
Malignant Neoplasms (Cancer)	172.5
Trachea, Bronchus, & Lung	37.6
Colon, Rectum, & Anus	19.4
Female Breast	11.6
Pancreas	LNE
Prostate	22.7
Leukemia	10.1
Chronic Lower Respiratory Diseases	47.5
Alzheimer's Disease	27.6
Cerebrovascular Disease	43.5
 ○ Accidents 	76.5
 Motor Vehicle Accidents 	38.6
Diabetes	26.4
Influenza and Pneumonia	19.5
 Intentional Self-Harm (Suicide) 	42.6
Chronic Liver Disease and Cirrhosis	24.1
Infant Mortality	5.61

Leading Causes of Death	Total Deaths
1. Malignant Neoplasms (Cancer)	123
2. Heart Disease	109
3. Accidents	44
4. Chronic Lower Respiratory Diseases	37
Cerebrovascular Disease	32
6. Alzheimer's Disease	23
T7. Diabetes	18
T7. Intentional Self-Harm (Suicide)	18
T9. Influenza and Pneumonia	13
T9. Chronic Liver Disease and Cirrhosis	13
Percent of Deaths due to tobacco use Median age at death	17.3 77

[•]Denotes a health status indicator which is significantly lower than the state average.

See technical notes for more information.


oDenotes a health status indicator which is significantly higher than the state average.

³All mortality rates except infant mortality are ageadjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

Sanborn County

Health Status Indicators 2013-2017

Demographic Information



Sanborn County is located in east central South Dakota and averages 4.1 persons per square mile.

2017 Population Information

Subject	Number	Percent
Total population	2,450	100.0
White	2,395	97.8
Black or African American	5	0.2
American Indian & Alaska Native	10	0.4
Asian	7	0.3
Native Hawaiian & Other Pacific Islander	1	0.0
Two or More Races	32	1.3
Under 5 years	219	8.9
Under 18 years	638	26.0
65 years and over	452	18.4

Source: United States Census Bureau, 2017 Population

Estimates

Natality	
Percent of Low Birth Weight Infants	4.9
Percent of Mothers Receiving	
Care in 1st Trimester	73.6
Percent of Mothers Who	
Used Tobacco While Pregnant ¹	16.9
Percent of Births Less Than 37 Wks. of Gestation	7.1
Average Age of Mother	28.0
Teenage Birth Rate ²	LNE
Percent White Births	99.5
Percent American Indian Births	LNE
 Percent Unmarried 	23.5
Percent WIC births	24.9
Percent Breastfeeding at discharge	83.0
 Percent Payment-Private Insurance 	79.8
Percent Payment-Medicaid	19.1
 Percent C-Section 	34.4

Mortality ³	
All Causes	753.9
Heart Disease	125.2
Malignant Neoplasms (Cancer)	167.5
Trachea, Bronchus, & Lung	55.7
Colon, Rectum, & Anus	15.9
Female Breast	LNE
Pancreas	14.5
Prostate	LNE
Leukemia	LNE
Chronic Lower Respiratory Diseases	34.1
 Alzheimer's Disease 	114.0
Cerebrovascular Disease	22.2
Accidents	38.1
Motor Vehicle Accidents	LNE
Diabetes	18.1
Influenza and Pneumonia	17.0
Intentional Self-Harm (Suicide)	LNE
Chronic Liver Disease and Cirrhosis	22.0
Infant Mortality	LNE

Leading Causes of Death	Total Deaths
Malignant Neoplasms (Cancer)	29
T2. Heart Disease	26
T2. Alzheimer's Disease	26
Chronic Lower Respiratory Diseases	7
5. Accidents	6
Cerebrovascular Disease	5
Percent of Deaths due to tobacco use Median age at death	12.5 84

See technical notes for more information.

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

³All mortality rates except infant mortality are ageadjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

Spink County

Health Status Indicators 2013-2017

Demographic Information

Spink County is located in the center of eastern South Dakota and averages 4.3 persons per square mile.

2017 Population Information

Subject	Number	Percent
Total population	6,410	100.0
White	6,177	96.4
Black or African American	43	0.7
American Indian & Alaska Native	112	1.7
Asian	7	0.1
Native Hawaiian & Other Pacific Islander	1	0.0
Two or More Races	70	1.1
Under 5 years	382	6.0
Under 18 years	1,503	23.4
65 years and over	1,299	20.3

Source: United States Census Bureau, 2017 Population Estimates

4.4
68.2
13.6
7.8
28.8
4.6
96.6
LNE
27.4
21.6
84.0
79.1
16.7
30.5

Mortality ³	
All Causes	727.5
Heart Disease	153.5
Malignant Neoplasms (Cancer)	167.1
Trachea, Bronchus, & Lung	36.9
Colon, Rectum, & Anus	22.2
Female Breast	34.4
Pancreas	13.4
Prostate	LNE
Leukemia	LNE
Chronic Lower Respiratory Diseases	39.2
Alzheimer's Disease	39.7
Cerebrovascular Disease	26.7
Accidents	78.0
Motor Vehicle Accidents	26.6
Diabetes	30.9
Influenza and Pneumonia	30.9
Intentional Self-Harm (Suicide)	LNE
 Chronic Liver Disease and Cirrhosis 	7.3
Infant Mortality	LNE

Leading Causes of Death	Total Deaths
Heart Disease Malignant Neoplasms (Cancer) Accidents Alzheimer's Disease Chronic Lower Respiratory Diseases Influenza and Pneumonia Cerebrovascular Disease Diabetes	91 83 28 27 22 21 17
 Pneumonitis Due to Solids and Liquids Unspecified Dementia Septicemia Disorders of Lipoprotein Metabolism And Other Lipidemias 	8 6 6
Percent of Deaths due to tobacco use Median age at death	10.4 82

- lower than the state average.
- oDenotes a health status indicator which is significantly higher than the state average.

•Denotes a health status indicator which is significantly

³All mortality rates except infant mortality are ageadjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of **Health Statistics**

l	Denotes	а	health	status	indicator	which	IS	significantly	lower
	than the st	tat	e avera	ige.					
ı	D 4	_	1 141-	- 4 - 4	the alternation	and the factor		- 1 1 .C 4 1 1	

oDenotes a health status indicator which is significantly higher than the state average.


¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Stanley County

Health Status Indicators 2013-2017

Demographic Information



Stanley County is located in the center of the state and averages 2.1 persons per square mile.

2017 Population Information

Subject	Number	Percent
Total population White Black or African American American Indian & Alaska Native Asian Native Hawaiian & Other Pacific Islander	3,011 2,691 23 212 10 0	100.0 89.4 0.8 7.0 0.3 0.0
Two or More Races Under 5 years Under 18 years 65 years and over	75 196 734 607	2.5 6.5 24.4 20.2

Source: United States Census Bureau, 2017 Population

Estimates

Natality	
Percent of Low Birth Weight Infants	5.3
 Percent of Mothers Receiving 	
Care in 1st Trimester	56.7
Percent of Mothers Who	
Used Tobacco While Pregnant ¹	10.0
Percent of Births Less Than 37 Wks. of Gestation	7.9
 Average Age of Mother 	28.9
Teenage Birth Rate ²	LNE
Percent White Births	84.7
Percent American Indian Births	9.5
Percent Unmarried	31.6
Percent WIC births	28.6
Percent Breastfeeding at discharge	85.2
Percent Payment-Private Insurance	72.5
Percent Payment-Medicaid	23.3
Percent C-Section	38.4

 All Causes Heart Disease Malignant Neoplasms (Cancer) Trachea, Bronchus, & Lung Colon, Rectum, & Anus Female Breast Pancreas Prostate Leukemia Chronic Lower Respiratory Diseases Alzheimer's Disease Accidents Motor Vehicle Accidents Influenza and Pneumonia Intentional Self-Harm (Suicide) Chronic Liver Disease and Cirrhosis 17.2 Chronic Lower Respiratory Disease 26.5 Influenza and Pneumonia Intentional Self-Harm (Suicide) 28.1 Infant Mortality LNE 	Mortality ³	
Malignant Neoplasms (Cancer) Trachea, Bronchus, & Lung Colon, Rectum, & Anus Pemale Breast Pancreas Prostate Leukemia Chronic Lower Respiratory Diseases Alzheimer's Disease Accidents Motor Vehicle Accidents Diabetes Influenza and Pneumonia Intentional Self-Harm (Suicide) Chronic Liver Disease and Cirrhosis IT3.8 Chronic Liver Disease and Cirrhosis	All Causes	561.7
Trachea, Bronchus, & Lung Colon, Rectum, & Anus Pemale Breast Pancreas Prostate Leukemia Chronic Lower Respiratory Diseases Alzheimer's Disease Accidents Motor Vehicle Accidents Diabetes Influenza and Pneumonia Intentional Self-Harm (Suicide) Colon, Rectum, & Lung LNE	Heart Disease	86.6
Colon, Rectum, & Anus Female Breast Pancreas Prostate Leukemia Chronic Lower Respiratory Diseases Alzheimer's Disease Cerebrovascular Disease Accidents Motor Vehicle Accidents Diabetes Influenza and Pneumonia Intentional Self-Harm (Suicide) Chronic Liver Disease and Cirrhosis Color Respiratory Disease LNE	Malignant Neoplasms (Cancer)	179.4
Female Breast LNE Pancreas 20.4 Prostate LNE Leukemia LNE Chronic Lower Respiratory Diseases 84.9 Alzheimer's Disease LNE • Cerebrovascular Disease 14.5 • Accidents 17.2 Motor Vehicle Accidents LNE Diabetes 26.5 Influenza and Pneumonia LNE Intentional Self-Harm (Suicide) 50.2 Chronic Liver Disease and Cirrhosis 28.1	Trachea, Bronchus, & Lung	57.8
Pancreas 20.4 Prostate LNE Leukemia LNE Chronic Lower Respiratory Diseases 84.9 Alzheimer's Disease LNE • Cerebrovascular Disease 14.5 • Accidents 17.2 Motor Vehicle Accidents LNE Diabetes 26.5 Influenza and Pneumonia LNE Intentional Self-Harm (Suicide) 50.2 Chronic Liver Disease and Cirrhosis 28.1	Colon, Rectum, & Anus	20.4
Prostate Leukemia LNE Chronic Lower Respiratory Diseases Alzheimer's Disease Cerebrovascular Disease Accidents Accidents Motor Vehicle Accidents LNE Diabetes Influenza and Pneumonia Intentional Self-Harm (Suicide) Chronic Liver Disease and Cirrhosis LNE	Female Breast	LNE
Leukemia LNE Chronic Lower Respiratory Diseases 84.9 Alzheimer's Disease LNE • Cerebrovascular Disease 14.5 • Accidents 17.2 Motor Vehicle Accidents LNE Diabetes 26.5 Influenza and Pneumonia LNE Intentional Self-Harm (Suicide) 50.2 Chronic Liver Disease and Cirrhosis 28.1	Pancreas	20.4
Chronic Lower Respiratory Diseases Alzheimer's Disease LNE Cerebrovascular Disease 14.5 Accidents 17.2 Motor Vehicle Accidents LNE Diabetes 16.5 Influenza and Pneumonia LNE Intentional Self-Harm (Suicide) Chronic Liver Disease and Cirrhosis 28.1	Prostate	
Alzheimer's Disease • Cerebrovascular Disease • Accidents • Motor Vehicle Accidents Diabetes Influenza and Pneumonia Intentional Self-Harm (Suicide) Chronic Liver Disease and Cirrhosis LNE LNE LNE LNE LNE LNE LNE LN	Leukemia	LNE
 Cerebrovascular Disease Accidents Motor Vehicle Accidents LNE Diabetes Influenza and Pneumonia LNE Intentional Self-Harm (Suicide) Chronic Liver Disease and Cirrhosis 14.5 LNE LNE Diabetes 26.5 Influenza and Pneumonia LNE LNE<!--</td--><td>Chronic Lower Respiratory Diseases</td><td>84.9</td>	Chronic Lower Respiratory Diseases	84.9
 Accidents Motor Vehicle Accidents LNE Diabetes Influenza and Pneumonia LNE Intentional Self-Harm (Suicide) Chronic Liver Disease and Cirrhosis 17.2 LNE Diabetes 26.5 Influenza and Pneumonia LNE LNE So.2 Chronic Liver Disease and Cirrhosis 	Alzheimer's Disease	LNE
Motor Vehicle Accidents LNE Diabetes 26.5 Influenza and Pneumonia LNE Intentional Self-Harm (Suicide) 50.2 Chronic Liver Disease and Cirrhosis 28.1	Cerebrovascular Disease	14.5
Diabetes 26.5 Influenza and Pneumonia LNE Intentional Self-Harm (Suicide) 50.2 Chronic Liver Disease and Cirrhosis 28.1	 Accidents 	17.2
Influenza and Pneumonia LNE Intentional Self-Harm (Suicide) 50.2 Chronic Liver Disease and Cirrhosis 28.1	Motor Vehicle Accidents	LNE
Intentional Self-Harm (Suicide) 50.2 Chronic Liver Disease and Cirrhosis 28.1	2.000.00	_0.0
Chronic Liver Disease and Cirrhosis 28.1	Influenza and Pneumonia	LNE
		50.2
Infant Mortality LNE		
•	Infant Mortality	LNE

Leading Causes of Death	Deaths
1. Malignant Neoplasms (Cancer)	35
T2. Heart Disease	16
T2. Chronic Lower Respiratory Diseases	16
4. Intentional Self-Harm (Suicide)	8
T5. Diabetes	5
T5. Chronic Liver Disease and Cirrhosis	5
Percent of Deaths due to tobacco use Median age at death	30.5 70

See technical notes for more information.

[•]Denotes a health status indicator which is significantly lower than the state average.

 $[\]circ \mbox{Denotes}$ a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

[•]Denotes a health status indicator which is significantly lower than the state average.


oDenotes a health status indicator which is significantly higher than the state average.

³All mortality rates except infant mortality are ageadjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

Sully County

Health Status Indicators 2013-2017

Demographic Information



Sully County is located in the central region of the state and averages 1.4 persons per square mile.

2017 Population Information

Subject	Number	Percent
Total population	1,407	100.0
White	1,337	95.0
Black or African American	6	0.4
American Indian & Alaska Native	33	2.3
Asian	2	0.1
Native Hawaiian & Other Pacific Islander	0	0.0
Two or More Races	29	2.1
Under 5 years Under 18 years	84 295	6.0 21.0
65 years and over	332	23.6

Source: United States Census Bureau, 2017 Population

Estimates

Natality	
Percent of Low Birth Weight Infants	4.8
 Percent of Mothers Receiving 	
Care in 1st Trimester	49.4
Percent of Mothers Who	
Used Tobacco While Pregnant ¹	10.8
Percent of Births Less Than 37 Wks. of Gestation	12.0
Average Age of Mother	26.9
Teenage Birth Rate ²	LNE
Percent White Births	84.3
Percent American Indian Births	12.0
Percent Unmarried	28.9
Percent WIC births	24.4
Percent Breastfeeding at discharge	77.1
Percent Payment-Private Insurance	68.3
Percent Payment-Medicaid	30.5
Percent C-Section	19.3

Mortality ³	
All Causes	409.9
Heart Disease	66.0
Malignant Neoplasms (Cancer)	133.1
Trachea, Bronchus, & Lung	LNE
Colon, Rectum, & Anus	LNE
Female Breast	LNE
Pancreas	LNE
Prostate	LNE
Leukemia	LNE
Chronic Lower Respiratory Diseases	34.7
Alzheimer's Disease	LNE
Cerebrovascular Disease	LNE
Accidents	LNE
Motor Vehicle Accidents	LNE
Diabetes	LNE
Influenza and Pneumonia	LNE
Intentional Self-Harm (Suicide)	LNE
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	LNE
Leading Causes of Death	Total Deaths

16

8

17.0

77

Denotes	a health	status	indicator	which	is	significa	ntly	lower
than the s	tate aver	age.						

 $[\]circ \text{Denotes}$ a health status indicator which is significantly higher than the state average.

193

Denotes	a health	status	indicator	which	is	significantly
lower than	the state	avera	ge.			

oDenotes a health status indicator which is significantly higher than the state average.

See technical notes for more information.

1. Malignant Neoplasms (Cancer)

Percent of Deaths due to tobacco use

2. Heart Disease

Median age at death

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

³All mortality rates except infant mortality are ageadjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

Todd County

Health Status Indicators 2013-2017

Demographic Information

Todd County is located in south central South Dakota, along the Nebraska border and averages 6.9 persons per square mile.

2017 Population Information

Subject	Number	Percent
Total population	10,065	100.0
White	882	8.8
Black or African American	50	0.5
American Indian & Alaska Native	8,855	88.0
Asian	20	0.2
Native Hawaiian & Other Pacific Islander	0	0.0
Two or More Races	258	2.6
Under 5 years	1,309	13.0
Under 18 years	4,151	41.2
65 years and over	727	7.2

Source: United States Census Bureau, 2017 Population

Estimates

Natality	
 Percent of Low Birth Weight Infants 	8.4
Percent of Mothers Receiving	
Care in 1st Trimester	39.5
 Percent of Mothers Who 	
Used Tobacco While Pregnant ¹	21.1
 Percent of Births Less Than 37 Wks. of Gestation 	11.5
 Average Age of Mother 	25.1
○ Teenage Birth Rate ²	55.6
Percent White Births	3.9
Percent American Indian Births	92.1
 Percent Unmarried 	85.6
Percent WIC births	69.2
 Percent Breastfeeding at discharge 	60.8
Percent Payment-Private Insurance	4.8
Percent Payment-Medicaid	60.8
Percent C-Section	36.7

Mortality ³	
All Causes	1,270.3
Heart Disease	202.5
 Malignant Neoplasms (Cancer) 	211.9
Trachea, Bronchus, & Lung	39.1
Colon, Rectum, & Anus	25.2
Female Breast	LNE
Pancreas	LNE
Prostate	46.9
Leukemia	24.6
Chronic Lower Respiratory Diseases	66.9
Alzheimer's Disease	23.9
Cerebrovascular Disease	36.9
 Accidents 	146.0
 Motor Vehicle Accidents 	80.9
 Diabetes 	103.7
 Influenza and Pneumonia 	51.3
 Intentional Self-Harm (Suicide) 	60.4
 Chronic Liver Disease and Cirrhosis 	75.1
 Infant Mortality 	15.92

Leading Causes of Death	lotai
Todamiy Gadooo or Dodan	Deaths
Malignant Neoplasms (Cancer)	69
2. Heart Disease	65
3. Accidents	64
4. Diabetes	34
5. Chronic Liver Disease and Cirrhosis	28
Intentional Self-Harm (Suicide)	27
7. Chronic Lower Respiratory Diseases	21
8. Influenza and Pneumonia	16
Mental and Behavioral Disorders Due	13
to Use of Alcohol	40
10. Cerebrovascular Disease	12
Percent of Deaths due to tobacco use	13.4
Median age at death	58

See technical notes for more information.

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.


[•]Denotes a health status indicator which is significantly lower than the state average.

 $[\]circ \mbox{Denotes}$ a health status indicator which is significantly higher than the state average.

³All mortality rates except infant mortality are ageadjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

Tripp County

Demographic Information



Tripp County is located along the Nebraska border in south central South Dakota and averages 3.5 persons per square mile.

2017 Population Information

Subject	Number	Percent
Total population	5,460	100.0
White	4,501	82.4
Black or African American	32	0.6
American Indian & Alaska Native	776	14.2
Asian	18	0.3
Native Hawaiian & Other Pacific Islander	0	0.0
Two or More Races	133	2.4
Under 5 years	375	6.9
Under 18 years	1,254	23.0
65 years and over	1,223	22.4

Source: United States Census Bureau, 2017 Population Estimates

Natality	
Percent of Low Birth Weight Infants	6.8
Percent of Mothers Receiving	
Care in 1st Trimester	71.4
Percent of Mothers Who	
Used Tobacco While Pregnant ¹	14.7
Percent of Births Less Than 37 Wks. of Gestation	10.0
Average Age of Mother	27.0
Teenage Birth Rate ²	9.0
Percent White Births	62.7
Percent American Indian Births	31.4
Percent Unmarried	44.9
Percent WIC births	44.5
Percent Breastfeeding at discharge	74.9
Percent Payment-Private Insurance	49.5
Percent Payment-Medicaid	47.3
Percent C-Section	37.6

Health Status Indicators 2013-2017

Mortality ³	
All Causes	741.3
Heart Disease	187.1
Malignant Neoplasms (Cancer)	144.9
Trachea, Bronchus, & Lung	32.0
Colon, Rectum, & Anus	20.5
Female Breast	LNE
Pancreas	9.5
Prostate	LNE
Leukemia	LNE
Chronic Lower Respiratory Diseases	44.2
 Alzheimer's Disease 	62.9
Cerebrovascular Disease	27.5
Accidents	40.7
Motor Vehicle Accidents	11.0
Diabetes	17.1
Influenza and Pneumonia	19.7
Intentional Self-Harm (Suicide)	LNE
Chronic Liver Disease and Cirrhosis	21.9
Infant Mortality	LNE

Leading Causes of Death	Total
Louding Oddood of Doddin	Deaths
Heart Disease	98
Malignant Neoplasms (Cancer)	72
Alzheimer's Disease	40
Chronic Lower Respiratory Diseases	23
Cerebrovascular Disease	17
6. Accidents	16
Influenza and Pneumonia	11
8. Diabetes	9
T9. Chronic Liver Disease and Cirrhosis	5
T9. Essential (Primary) Hypertension and Hypertensive Renal Disease	5
T9. Pneumonitis Due to Solids and Liquids	5
Percent of Deaths due to tobacco use Median age at death	19.0 83

- lower than the state average.
- oDenotes a health status indicator which is significantly higher than the state average.

•Denotes a health status indicator which is significantly

³All mortality rates except infant mortality are ageadjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births. See technical notes for more information.

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.


¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Turner County

Health Status Indicators 2013-2017

Demographic Information



Turner County is located in southeastern South Dakota and averages 13.5 persons per square mile.

2017 Population Information

Subject	Number	Percent
Total population	8,315	100.0
White Black or African American	8,083 42	97.2 0.5
American Indian & Alaska Native	84	1.0
Asian	22	0.3
Native Hawaiian & Other Pacific Islander	5	0.1
Two or More Races	79	1.0
Under 5 years	514	6.2
Under 18 years	1,989	23.9
65 years and over	1,733	20.8

Source: United States Census Bureau, 2017 Population Estimates

Natality	
Percent of Low Birth Weight Infants	5.1
 Percent of Mothers Receiving 	
Care in 1st Trimester	81.6
Percent of Mothers Who	
Used Tobacco While Pregnant ¹	8.9
Percent of Births Less Than 37 Wks. of Gestation	7.3
Average Age of Mother	29.3
Teenage Birth Rate ²	LNE
Percent White Births	98.0
Percent American Indian Births	0.7
Percent Unmarried	19.4
Percent WIC births	19.2
Percent Breastfeeding at discharge	81.0
Percent Payment-Private Insurance	75.7
Percent Payment-Medicaid	18.3
Percent C-Section	23.8

Mortality ³	
o All Causes	789.1
Heart Disease	168.3
Malignant Neoplasms (Cancer)	161.6
Trachea, Bronchus, & Lung	40.6
Colon, Rectum, & Anus	18.5
Female Breast	14.7
Pancreas	11.1
Prostate	12.7
Leukemia	11.0
Chronic Lower Respiratory Diseases	61.1
 Alzheimer's Disease 	68.9
Cerebrovascular Disease	23.6
Accidents	54.7
Motor Vehicle Accidents	29.1
Diabetes	13.2
Influenza and Pneumonia	10.7
Intentional Self-Harm (Suicide)	21.1
Chronic Liver Disease and Cirrhosis	19.6
Infant Mortality	13.36

Leading Causes of Death	Total Deaths
1. Heart Disease	126
2. Malignant Neoplasms (Cancer)	105
3. Alzheimer's Disease	57
4. Chronic Lower Respiratory Diseases	42
5. Accidents	31
Cerebrovascular Disease	19
T7. Diabetes	11
T7. Parkinson's Disease	11
T9. Influenza and Pneumonia	10
T9. Chronic Liver Disease and Cirrhosis	10
Percent of Deaths due to tobacco use Median age at death	13.0 82

See technical notes for more information.

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.


²Teenage Birth rate is live births per 1,000 females age 15-17.

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

³All mortality rates except infant mortality are ageadjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

Union County



Union County is located in the southeastern corner of the state and averages 31.3 persons per square mile.

2017 Population Information

Subject	Number	Percent
Total population White Black or African American	15,029 14,284 164	100.0 95.0 1.1
American Indian & Alaska Native	133	0.9
Asian	193	1.3
Native Hawaiian & Other Pacific Islander	10	0.1
Two or More Races	245	1.6
Under 5 years	860	5.7
Under 18 years	3,608	24.0
65 years and over	2,714	18.1

•Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Natality	
Percent of Low Birth Weight Infants	6.4
 Percent of Mothers Receiving 	
Care in 1st Trimester	86.4
 Percent of Mothers Who 	
Used Tobacco While Pregnant ¹	7.6
Percent of Births Less Than 37 Wks. of Gestation	8.7
 Average Age of Mother 	29.1
Teenage Birth Rate ²	LNE
Percent White Births	92.1
Percent American Indian Births	1.1
 Percent Unmarried 	20.5
Percent WIC births	13.6
Percent Breastfeeding at discharge	83.1
 Percent Payment-Private Insurance 	77.8
Percent Payment-Medicaid	18.7
 Percent C-Section 	30.1

Health Status Indicators 2013-2017

Mortality ³	
All Causes	633.9
Heart Disease	138.5
Malignant Neoplasms (Cancer)	154.3
Trachea, Bronchus, & Lung	50.0
Colon, Rectum, & Anus	10.2
Female Breast	19.4
Pancreas	10.3
 Prostate 	7.0
Leukemia	6.5
Chronic Lower Respiratory Diseases	52.9
Alzheimer's Disease	25.4
Cerebrovascular Disease	25.1
 Accidents 	28.5
 Motor Vehicle Accidents 	4.9
Diabetes	14.0
Influenza and Pneumonia	21.1
Intentional Self-Harm (Suicide)	11.2
Chronic Liver Disease and Cirrhosis	13.9
Infant Mortality	3.69

Leading Causes of Death	Deaths
1. Malignant Neoplasms (Cancer)	147
2. Heart Disease	138
Chronic Lower Respiratory Diseases	49
4. Accidents	26
T5. Alzheimer's Disease	25
T5. Cerebrovascular Disease	25
7. Influenza and Pneumonia	20
Essential (Primary) Hypertension and Hypertensive Renal Disease	15
9. Diabetes	14
10. Chronic Liver Disease and Cirrhosis	11
Percent of Deaths due to tobacco use	18.7
Median age at death	79

Total

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Source: United States Census Bureau, 2017 Population Estimates

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

³All mortality rates except infant mortality are ageadjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

Walworth County

Demographic Information

Walworth County is located in north central South Dakota, near the North Dakota border and averages 7.7 persons per square mile.

2017 Population Information

Subject	Number	Percent
Total population	5,543	100.0
White	4,440	80.1
Black or African American	29	0.5
American Indian & Alaska Native	791	14.3
Asian	104	1.9
Native Hawaiian & Other Pacific Islander	1	0.0
Two or More Races	178	3.2
Under 5 years	384	6.9
Under 18 years	1,298	23.4
65 years and over	1,311	23.7

Source: United States Census Bureau, 2017 Population

Estimates

Natality	
Percent of Low Birth Weight Infants	5.9
Percent of Mothers Receiving	
Care in 1st Trimester	65.2
Percent of Mothers Who	
Used Tobacco While Pregnant ¹	17.0
Percent of Births Less Than 37 Wks. of Gestation	9.0
 Average Age of Mother 	26.8
Teenage Birth Rate ²	5.9
Percent White Births	67.9
Percent American Indian Births	23.1
Percent Unmarried	41.3
Percent WIC births	40.3
Percent Breastfeeding at discharge	72.6
Percent Payment-Private Insurance Percent Payment Medicaid	50.7
Percent Payment-Medicaid Percent C-Section	40.4 27.9
reiteni o-settion	21.9

Health Status Indicators 2013-2017

	Mortality ³		
 All Causes 	-	809.8	
Heart Disease		150.1	
Malignant Neopl	asms (Cancer)	147.6	
Trachea, Br	onchus, & Lung	40.2	
Colon, Rect	um, & Anus	13.9	
Female Bre	ast	27.1	
Pancreas		LNE	
Prostate		LNE	
Leukemia		6.2	
Chronic Lower F	Respiratory Diseases	59.4	
 Alzheimer's D 	sease	68.9	
Cerebrovascular	Disease	36.5	
Accidents		84.0	
Motor Vehic	le Accidents	39.3	
Diabetes		27.9	
Influenza and Pr	neumonia	30.8	
Intentional Self-I		31.9	
Chronic Liver Di	sease and Cirrhosis	18.2	
Infant Mortality		LNE	

Leading Causes of Death	Total
	Deaths
1. Heart Disease	89
Malignant Neoplasms (Cancer)	72
3. Alzheimer's Disease	49
4. Chronic Lower Respiratory Diseases	31
5. Accidents	24
6. Influenza and Pneumonia	20
T7. Cerebrovascular Disease	18
T7. Diabetes	18
Unspecified Dementia	12
 Nephritis, Nephrotic Syndrome, and Nephrosis 	8
Percent of Deaths due to tobacco use Median age at death	16.6 83

- •Denotes a health status indicator which is significantly lower than the state average.
- oDenotes a health status indicator which is significantly higher than the state average.
- ¹Data for mothers who used tobacco are self-reported.

198

²Teenage Birth rate is live births per 1,000 females age 15-17.

Denote	s a	health	status	indicator	which	is	significantly
lower th	an t	he state	avera	ge.			


 $[\]circ \mbox{Denotes}$ a health status indicator which is significantly higher than the state average.

See technical notes for more information.

³All mortality rates except infant mortality are ageadjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

Yankton County

Demographic Information



Yankton County is located in southeastern South Dakota on the Nebraska border and averages 43.1 persons per square mile.

2017 Population Information

Subject Total population White Black or African American American Indian & Alaska Native Asian Native Hawaiian & Other Pacific Islander Two or More Races	Number 22,662 20,873 485 686 195 11 412	Percent 100.0 92.1 2.1 3.0 0.9 0.0 1.8
Under 5 years	1,368	6.0
Under 18 years	4,784	21.1
65 years and over	4,212	18.6

Source: United States Census Bureau, 2017 Population

Estimates

Health Status Indicators 2013-2017

Natality	
Percent of Low Birth Weight Infants	7.6
 Percent of Mothers Receiving 	
Care in 1st Trimester	84.8
 Percent of Mothers Who 	
Used Tobacco While Pregnant ¹	19.8
Percent of Births Less Than 37 Wks. of Gestation	7.1
Average Age of Mother	27.5
Teenage Birth Rate ²	10.5
Percent White Births	87.0
Percent American Indian Births	4.3
Percent Unmarried	40.2
Percent WIC births	33.9
Percent Breastfeeding at discharge	76.1
 Percent Payment-Private Insurance 	66.0
Percent Payment-Medicaid	32.3
Percent C-Section	28.9

Mortality ³	
All Causes	701.4
Heart Disease	166.5
Malignant Neoplasms (Cancer)	137.3
Trachea, Bronchus, & Lung	39.2
Colon, Rectum, & Anus	14.2
Female Breast	14.6
 Pancreas 	3.3
Prostate	16.7
Leukemia	11.1
Chronic Lower Respiratory Diseases	53.3
Alzheimer's Disease	33.1
Cerebrovascular Disease	34.9
Accidents	53.0
Motor Vehicle Accidents	18.7
Diabetes	27.9
Influenza and Pneumonia	19.9
Intentional Self-Harm (Suicide)	15.5
 Chronic Liver Disease and Cirrhosis 	7.0
Infant Mortality	9.37

Leading Causes of Death	Total Deaths
1. Heart Disease	295
Malignant Neoplasms (Cancer)	221
Chronic Lower Respiratory Diseases	93
4. Accidents	74
5. Alzheimer's Disease	70
Cerebrovascular Disease	67
7. Diabetes	47
Influenza and Pneumonia	37
Intentional Self-Harm (Suicide)	20
 Essential (Primary) Hypertension and Hypertensive Renal Disease 	19
Percent of Deaths due to tobacco use Median age at death	20.8 80

•Denotes a health status indicator which is significantly lower than the state average.

See technical notes for more information.

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.


²Teenage Birth rate is live births per 1,000 females age 15-17.

oDenotes a health status indicator which is significantly higher than the state average.

³All mortality rates except infant mortality are ageadjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

Ziebach County

Demographic Information



Ziebach County is located in north central South Dakota and averages 1.4 persons per square mile.

2017 Population Information

Subject	Number	Percent
Total population	2,756	100.0
White	689	25.0
Black or African American	13	0.5
American Indian & Alaska Native	1,933	70.1
Asian	10	0.4
Native Hawaiian & Other Pacific Islander	1	0.0
Two or More Races	110	4.0
Under 5 years	166	6.0
Under 18 years	830	30.1
65 years and over	261	9.5

Source: United States Census Bureau, 2017 Population

Estimates

Natality	
Percent of Low Birth Weight Infants	6.9
 Percent of Mothers Receiving 	
Care in 1st Trimester	53.2
Percent of Mothers Who	
Used Tobacco While Pregnant ¹	20.1
Percent of Births Less Than 37 Wks. of Gestation	14.1
 Average Age of Mother 	26.5
Teenage Birth Rate ²	13.0
Percent White Births	11.8
Percent American Indian Births	81.3
 Percent Unmarried 	70.8
 Percent WIC births 	76.2
Percent Breastfeeding at discharge	66.9
Percent Payment-Private Insurance	18.8
Percent Payment-Medicaid	68.1
Percent C-Section	24.3

Health Status Indicators 2013-2017

Mortality ³	
All Causes	778.4
Heart Disease	171.8
 Malignant Neoplasms (Cancer) 	64.9
Trachea, Bronchus, & Lung	LNE
Colon, Rectum, & Anus	25.4
Female Breast	LNE
Pancreas	LNE
Prostate	LNE
Leukemia	LNE
Chronic Lower Respiratory Diseases	69.2
Alzheimer's Disease	LNE
Cerebrovascular Disease	LNE
o Accidents	120.8
Motor Vehicle Accidents	46.1
Diabetes	40.0
Influenza and Pneumonia	23.0
Intentional Self-Harm (Suicide)	34.2
Chronic Liver Disease and Cirrhosis	44.5
Infant Mortality	LNE

Leading Causes of Death	Total Deaths
T1. Heart Disease	15
T1. Accidents	15
Malignant Neoplasms (Cancer)	7
T4. Chronic Lower Respiratory Diseases	5
T4. Intentional Self-Harm (Suicide)	5
T4. Chronic Liver Disease and Cirrhosis	5
T4. Septicemia	5
Percent of Deaths due to tobacco use Median age at death	10.0 59

See technical notes for more information.

[•]Denotes a health status indicator which is significantly lower than the state average.

 $[\]circ \mbox{Denotes}$ a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

[•]Denotes a health status indicator which is significantly lower than the state average.

oDenotes a health status indicator which is significantly higher than the state average.

³All mortality rates except infant mortality are ageadjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.