

# Alcohol Use

## DRANK IN PAST 30 DAYS

**Definition:** Respondents who report drinking alcohol in the past 30 days.


### Prevalence of Drinking in Past 30 Days

- South Dakota 59%
- Nationwide median 57%

### Healthy People 2020 Objective


There was no comparable Healthy People 2020 Objective for drinking alcohol in the past 30 days.

**Figure 64**  
**Percent of Respondents Who Drank Alcohol in the Past 30 Days, 1995, 1997, 1999, and 2001-2010**



Source: The Behavioral Risk Factor Surveillance System, South Dakota Department of Health, 1995, 1997, 1999, and 2001-2010

**Figure 65**  
**Percent of Respondents Who Drank Alcohol in the Past 30 Days, 2011**



Note: Beginning in 2011, the CDC began using a different methodology to weight the data, therefore current data estimates cannot be compared to previous years.

Source: The Behavioral Risk Factor Surveillance System, South Dakota Department of Health, 2011

**Table 43**  
**Respondents Who Drank Alcohol in Past 30 Days, 2011**


		# Resp.	% Drank Alcohol Past 30 Days	95% CI	
				Lower	Upper
<b>Total</b>		<b>7,941</b>	<b>59%</b>	<b>57%</b>	<b>61%</b>
<b>Gender</b>	Male	3,170	64%	60%	67%
	Female	4,771	54%	51%	57%
<b>Age</b>	18-44	2,111	67%	63%	70%
	45-64	3,136	60%	57%	63%
	65+	2,652	38%	34%	41%
<b>Race</b>	White	6,863	60%	58%	62%
	American Indian	792	44%	36%	53%
<b>Ethnicity</b>	Hispanic	102	57%	42%	71%
	Non-Hispanic	7,827	59%	57%	61%
<b>Household Income</b>	Less than \$25,000	1,922	47%	42%	51%
	\$25,000-\$49,999	2,041	57%	53%	62%
	\$50,000+	2,840	75%	72%	78%
<b>Education</b>	High School, G.E.D., or Less	3,067	48%	44%	51%
	Some Post-High School	2,341	66%	63%	70%
	College Graduate	2,522	69%	65%	72%
<b>Employment Status</b>	Employed for Wages	3,517	67%	64%	70%
	Self-employed	865	67%	61%	73%
	Unemployed	317	55%	45%	65%
	Homemaker	441	48%	39%	57%
	Student	180	61%	49%	72%
	Retired	2,227	41%	38%	45%
	Unable to Work	383	31%	23%	40%
<b>Marital Status</b>	Married/Unmarried Couple	4,642	63%	60%	65%
	Divorced/Separated	1,053	53%	48%	59%
	Widowed	1,199	30%	25%	35%
	Never Married	1,038	62%	56%	67%
<b>Home Ownership Status</b>	Own Home	6,042	61%	59%	63%
	Rent Home	1,601	55%	51%	59%
<b>Children Status</b>	Children in Household (Ages 18-44)	1,441	66%	62%	71%
	No Children in Household (Ages 18-44)	668	68%	62%	74%
<b>Phone Status</b>	Landline	6,991	57%	55%	60%
	Cell Phone	950	67%	63%	70%

Source: The Behavioral Risk Factor Surveillance System, South Dakota Department of Health, 2011

### **National Statistics**

The national median for respondents who drank in the past 30 days was 57.1 percent, as shown in Figure 66 on the next page. South Dakota had 58.8 percent of respondents who drank in the past 30 days. Utah had the lowest percent of respondents who drank in the past 30 days with 29.5 percent, while the District of Columbia had the highest with 67.4 percent.

**Figure 66**  
**Nationally, Respondents Who Drank Alcohol in Past 30 Days, 2011**



Source: The Behavioral Risk Factor Surveillance System, South Dakota Department of Health, 2011

**Further Analysis**

The following data show the percent of those who drank alcohol in the past 30 days for various health behaviors and conditions. For example, 52 percent of respondents who have high blood pressure drank alcohol in the past 30 days, while 62 percent of respondents who do not have high blood pressure drank alcohol in the past 30 days.

Health Behavior or Condition	# Respondents	% Drank Alcohol in Past 30 Days	95% CI	
			Lower	Upper
Underweight (BMI < 18.5)	*	*	*	*
Recommended (BMI = 18.5-24.9)	2,381	61%	57%	64%
Overweight (BMI = 25.0-29.9)	2,837	62%	58%	65%
Obese - Class I (BMI = 30.0-34.9)	1,461	59%	55%	64%
Obese - Classes II & III (BMI = 35.0+)	755	54%	47%	61%
Leisure Time Exercise	5,716	62%	60%	65%
No Leisure Time Exercise	2,217	49%	45%	54%
Meet Aerobic Recommendations	3,702	62%	59%	65%
Did Not Meet Aerobic Recommendations	4,039	56%	53%	59%
At Least Five Servings of Fruits and Vegetables	1,030	53%	47%	58%
Less Than Five Servings of Fruits and Vegetables	6,875	60%	57%	62%
Current Smoker	1,409	64%	60%	69%
Former Smoker	2,383	58%	54%	62%
Never Smoked	4,120	57%	54%	59%
Smokeless Tobacco	375	69%	59%	78%
No Smokeless Tobacco	7,566	58%	56%	60%

**Table 44 (continued)**  
**Drank Alcohol in Past 30 Days for Selected Health Behaviors and Conditions, 2011**

Health Behavior or Condition	# Respondents	% Drank Alcohol in Past 30 Days	95% CI	
			Lower	Upper
Diabetes	960	36%	30%	42%
No Diabetes	6,972	61%	59%	63%
High Blood Pressure	3,103	52%	49%	55%
No High Blood Pressure	4,829	62%	59%	64%
High Blood Cholesterol	2,781	54%	51%	58%
No High Blood Cholesterol	3,894	60%	57%	63%
Flu Shot (65+)	1,864	37%	33%	41%
No Flu Shot (65+)	774	41%	34%	48%
Pneumonia Shot (65+)	1,741	35%	31%	39%
No Pneumonia Shot (65+)	811	45%	39%	52%
Always/Almost Always Wear Seat Belt	6,732	58%	56%	60%
Sometimes/Seldom/Never Wear Seat Belt	1,187	63%	58%	68%
Previously Had Heart Attack	545	33%	27%	41%
Never Had Heart Attack	7,362	60%	58%	63%
Have Angina or Coronary Heart Disease	520	36%	29%	44%
Do Not Have Angina or Coronary Heart Disease	7,360	60%	58%	62%
Previously Had a Stroke	295	20%	14%	28%
Never Had a Stroke	7,629	60%	58%	62%
Cancer	1,319	44%	39%	49%
No Cancer	6,592	61%	59%	63%
Skin Cancer	734	42%	36%	48%
No Skin Cancer	7,188	60%	58%	62%
Current Asthma	619	47%	39%	54%
Former Asthma	245	52%	40%	64%
Never Had Asthma	7,030	60%	58%	62%
Arthritis	2,501	45%	41%	49%
No Arthritis	5,402	63%	61%	66%
COPD	536	40%	32%	48%
No COPD	7,379	60%	58%	62%
Depression	1,283	51%	46%	56%
No Depression	6,632	60%	58%	63%
Kidney Disease	247	38%	28%	49%
No Kidney Disease	7,675	59%	57%	61%
Vision Impairment	1,703	45%	41%	50%
No Vision Impairment	6,200	62%	59%	64%
Good/Very Good/Excellent Health Status	6,614	62%	60%	65%
Fair/Poor Health Status	1,318	38%	33%	43%
Physical Health Not Good for 0-29 Days of the Past 30	7,272	61%	58%	63%
Physical Health Not Good for 30 Days of the Past 30	528	33%	26%	41%
Mental Health Not Good for 0-19 Days of the Past 30	7,415	60%	57%	62%
Mental Health Not Good for 20-30 Days of the Past 30	406	50%	41%	59%
Usual Activities Unattainable for 0-9 Days of the Past 30	7,304	60%	58%	62%
Usual Activities Unattainable for 10-30 Days of the Past 30	561	41%	34%	49%
Physical, Mental, or Emotional Problems	2,296	46%	42%	50%
No Physical, Mental, or Emotional Problems	5,628	63%	61%	65%
Disability with Special Equipment Needed	816	32%	26%	39%
No Disability with Special Equipment Needed	7,123	61%	59%	63%
Been Tested for HIV (18-64)	1,577	62%	58%	67%
Never Been Tested for HIV (18-64)	6,129	58%	56%	60%

Note: \*Results based on sample sizes less than 100 have been suppressed.

Source: The Behavioral Risk Factor Surveillance System, South Dakota Department of Health, 2011

## **BINGE DRINKING**

**Definition:** Male respondents who report having five or more alcoholic drinks on one occasion or female respondents who have four or more alcoholic drinks on one occasion, one or more times in the past month.


### **Prevalence of Binge Drinking**

- South Dakota 22%
- Nationwide median 18%

### **Healthy People 2020 Objective**


There was no comparable Healthy People 2020 Objective for binge drinking.

**Figure 67**  
**Percent of Respondents Who Engage in Binge Drinking,**  
**1987-1993, 1995, 1997, 1999, and 2001-2010**



Source: The Behavioral Risk Factor Surveillance System, South Dakota Department of Health, 1987-1993, 1995, 1997, 1999, and 2001-2010

**Figure 68**  
**Percent of Respondents Who Engage in Binge Drinking, 2011**



Note: Beginning in 2011, the CDC began using a different methodology to weight the data, therefore current data estimates cannot be compared to previous years.

Source: The Behavioral Risk Factor Surveillance System, South Dakota Department of Health, 2011

**Table 45  
Respondents Who Engage in Binge Drinking, 2011**


		# Resp.	% Binge Drinkers	95% CI	
				Lower	Upper
<b>Total</b>		<b>7,900</b>	<b>22%</b>	<b>20%</b>	<b>24%</b>
<b>Gender</b>	Male	3,148	29%	26%	32%
	Female	4,752	16%	14%	18%
<b>Age</b>	18-44	2,086	36%	32%	39%
	45-64	3,125	15%	13%	18%
	65+	2,647	3%	2%	5%
<b>Race</b>	White	6,838	22%	20%	24%
	American Indian	781	28%	20%	37%
<b>Ethnicity</b>	Hispanic	100	24%	13%	40%
	Non-Hispanic	7,788	22%	20%	24%
<b>Household Income</b>	Less than \$25,000	1,910	21%	17%	24%
	\$25,000-\$49,999	2,033	20%	16%	24%
	\$50,000+	2,830	27%	24%	30%
<b>Education</b>	High School, G.E.D., or Less	3,046	18%	15%	21%
	Some Post-High School	2,326	26%	23%	30%
	College Graduate	2,517	23%	20%	27%
<b>Employment Status</b>	Employed for Wages	3,497	27%	24%	30%
	Self-employed	864	26%	20%	32%
	Unemployed	312	30%	21%	39%
	Homemaker	439	14%	7%	26%
	Student	178	38%	28%	49%
	Retired	2,221	5%	3%	6%
<b>Marital Status</b>	Unable to Work	378	12%	7%	20%
	Married/Unmarried Couple	4,622	20%	17%	22%
	Divorced/Separated	1,048	20%	16%	24%
	Widowed	1,196	4%	2%	7%
	Never Married	1,025	36%	31%	42%
	<b>Home Ownership Status</b>	Own Home	6,021	19%	17%
Rent Home		1,586	30%	26%	35%
<b>Children Status</b>	Children in Household (Ages 18-44)	1,429	34%	29%	38%
	No Children in Household (Ages 18-44)	655	39%	33%	45%
<b>Phone Status</b>	Landline	6,964	19%	17%	22%
	Cell Phone	936	37%	34%	42%

Source: The Behavioral Risk Factor Surveillance System, South Dakota Department of Health, 2011

### **National Statistics**

The national median for respondents who binge drink is 15.7 percent, as shown in Figure 69 on the next page. South Dakota had 22.1 percent of respondents who binge drink. Tennessee had the lowest percent of respondents who binge drink with 10 percent, while the District of Columbia had the highest with 25 percent.

**Figure 69**  
**Nationally, Respondents Who Engage in Binge Drinking, 2011**



Source: The Behavioral Risk Factor Surveillance System, South Dakota Department of Health, 2011

**Further Analysis**

The following data show the percent of those who engage in binge drinking for various health behaviors and conditions. For example, 49 percent of respondents who use smokeless tobacco also engage in binge drinking, while 20 percent of respondents who do not use smokeless tobacco engage in binge drinking.

Health Behavior or Condition	# Respondents	% Binge Drinkers	95% CI	
			Lower	Upper
Underweight (BMI < 18.5)	*	*	*	*
Recommended (BMI = 18.5-24.9)	2,359	22%	18%	26%
Overweight (BMI = 25.0-29.9)	2,829	25%	22%	28%
Obese - Class I (BMI = 30.0-34.9)	1,455	20%	16%	24%
Obese - Classes II & III (BMI = 35.0+)	753	26%	19%	34%
Leisure Time Exercise	5,688	24%	21%	26%
No Leisure Time Exercise	2,204	18%	15%	22%
Meet Aerobic Recommendations	3,685	22%	19%	25%
Did Not Meet Aerobic Recommendations	4,020	23%	20%	26%
At Least Five Servings of Fruits and Vegetables	1,028	8%	6%	11%
Less Than Five Servings of Fruits and Vegetables	6,837	24%	22%	26%
Current Smoker	1,389	38%	33%	44%
Former Smoker	2,372	17%	14%	20%
Never Smoked	4,111	17%	15%	20%

**Table 46 (continued)**  
**Binge Drinking for Selected Health Behaviors and Conditions, 2011**

Health Behavior or Condition	# Respondents	% Binge Drinkers	95% CI	
			Lower	Upper
Smokeless Tobacco	368	49%	38%	59%
No Smokeless Tobacco	7,532	20%	18%	22%
Diabetes	959	12%	8%	16%
No Diabetes	6,932	23%	21%	25%
High Blood Pressure	3,088	17%	14%	20%
No High Blood Pressure	4,803	25%	22%	27%
High Blood Cholesterol	2,775	15%	12%	18%
No High Blood Cholesterol	3,875	19%	17%	22%
Flu Shot (65+)	1,860	3%	2%	5%
No Flu Shot (65+)	773	4%	2%	7%
Pneumonia Shot (65+)	1,737	3%	2%	5%
No Pneumonia Shot (65+)	810	4%	2%	7%
Heavy Drinker	393	88%	82%	92%
Not a Heavy Drinker	7,460	17%	16%	19%
Always/Almost Always Wear Seat Belt	6,706	19%	17%	21%
Sometimes/Seldom/Never Wear Seat Belt	1,172	38%	32%	44%
Previously Had Heart Attack	544	6%	3%	11%
Never Had Heart Attack	7,322	23%	21%	25%
Have Angina or Coronary Heart Disease	518	6%	3%	11%
Do Not Have Angina or Coronary Heart Disease	7,321	23%	21%	25%
Previously Had a Stroke	294	5%	2%	12%
Never Had a Stroke	7,589	23%	21%	25%
Cancer	1,316	11%	8%	15%
No Cancer	6,554	24%	22%	26%
Skin Cancer	733	6%	4%	9%
No Skin Cancer	7,148	23%	21%	25%
Current Asthma	615	18%	12%	25%
Former Asthma	243	25%	15%	37%
Never Had Asthma	6,995	22%	20%	24%
Arthritis	2,489	10%	8%	13%
No Arthritis	5,373	26%	24%	28%
COPD	532	13%	8%	20%
No COPD	7,342	23%	21%	25%
Depression	1,275	21%	17%	26%
No Depression	6,599	22%	20%	24%
Kidney Disease	247	15%	8%	27%
No Kidney Disease	7,634	22%	20%	24%
Vision Impairment	1,696	13%	10%	17%
No Vision Impairment	6,166	24%	22%	26%
Good/Very Good/Excellent Health Status	6,580	24%	22%	26%
Fair/Poor Health Status	1,311	13%	10%	18%
Physical Health Not Good for 0-29 Days of the Past 30	7,234	23%	21%	25%
Physical Health Not Good for 30 Days of the Past 30	526	10%	6%	16%
Mental Health Not Good for 0-19 Days of the Past 30	7,380	22%	20%	24%
Mental Health Not Good for 20-30 Days of the Past 30	401	25%	16%	36%
Usual Activities Unattainable for 0-9 Days of the Past 30	7,268	23%	21%	25%
Usual Activities Unattainable for 10-30 Days of the Past 30	558	16%	11%	22%
Physical, Mental, or Emotional Problems	2,282	13%	10%	16%
No Physical, Mental, or Emotional Problems	5,601	25%	23%	28%
Disability with Special Equipment Needed	814	7%	4%	12%
No Disability with Special Equipment Needed	7,084	23%	21%	25%
Been Tested for HIV (18-64)	1,563	28%	24%	32%
Never Been Tested for HIV (18-64)	6,108	20%	18%	23%

Note: \*Results based on sample sizes less than 100 have been suppressed.

Source: The Behavioral Risk Factor Surveillance System, South Dakota Department of Health, 2011

## HEAVY DRINKING

**Definition:** Male respondents who report having more than 2 drinks per day, or female respondents who report having more than 1 drink per day.


### Prevalence of Heavy Drinking

- South Dakota 6%
- Nationwide median 7%

### Healthy People 2020 Objective


There was no comparable Healthy People 2020 Objective for heavy drinking.

**Figure 70**  
**Percent of Respondents Who Engage in Heavy Drinking,**  
**1987-1993, 1995, 1997, 1999, and 2001-2010**



Source: The Behavioral Risk Factor Surveillance System, South Dakota Department of Health, 1987-1993, 1995, 1997, 1999, and 2001-2010

**Figure 71**  
**Percent of Respondents Who Engage in Heavy Drinking, 2011**



Note: Beginning in 2011, the CDC began using a different methodology to weight the data, therefore current data estimates cannot be compared to previous years.

Source: The Behavioral Risk Factor Surveillance System, South Dakota Department of Health, 2011


<b>Table 47</b>					
<b>Respondents Who Engage in Heavy Drinking, 2011</b>					
		<b># Resp.</b>	<b>% Heavy Drinkers</b>	<b>95% CI</b>	
				<b>Lower</b>	<b>Upper</b>
<b>Total</b>		<b>7,874</b>	<b>6%</b>	<b>5%</b>	<b>7%</b>
<b>Gender</b>	Male	3,129	7%	6%	9%
	Female	4,745	5%	4%	6%
<b>Age</b>	18-44	2,067	8%	7%	11%
	45-64	3,119	5%	4%	6%
	65+	2,646	2%	1%	3%
<b>Race</b>	White	6,825	6%	5%	7%
	American Indian	768	5%	3%	7%
<b>Ethnicity</b>	Hispanic	101	12%	4%	30%
	Non-Hispanic	7,761	6%	5%	7%
<b>Household Income</b>	Less than \$25,000	1,899	7%	5%	9%
	\$25,000-\$49,999	2,029	6%	4%	9%
	\$50,000+	2,825	6%	5%	8%
<b>Education</b>	High School, G.E.D., or Less	3,025	5%	4%	7%
	Some Post-High School	2,326	7%	5%	9%
	College Graduate	2,512	5%	4%	7%
<b>Employment Status</b>	Employed for Wages	3,488	6%	5%	8%
	Self-employed	857	7%	4%	10%
	Unemployed	308	8%	4%	16%
	Homemaker	435	3%	1%	7%
	Student	177	14%	7%	24%
	Retired	2,223	3%	2%	4%
<b>Marital Status</b>	Unable to Work	375	3%	1%	12%
	Married/Unmarried Couple	4,617	5%	4%	6%
	Divorced/Separated	1,044	6%	4%	9%
	Widowed	1,191	2%	1%	3%
	Never Married	1,013	11%	8%	15%
	<b>Home Ownership Status</b>	Own Home	6,012	6%	5%
Rent Home		1,572	6%	4%	8%
<b>Children Status</b>	Children in Household (Ages 18-44)	1,420	6%	5%	9%
	No Children in Household (Ages 18-44)	645	11%	8%	16%
<b>Phone Status</b>	Landline	6,941	5%	4%	6%
	Cell Phone	933	13%	10%	16%

Source: The Behavioral Risk Factor Surveillance System, South Dakota Department of Health, 2011

### **National Statistics**

The national median for respondents who engage in heavy drinking is 6.6 percent, as shown in Figure 72 on the next page. South Dakota had 5.9 percent of respondents who engage in heavy drinking. Tennessee had the lowest percent of respondents who engage in heavy drinking with 3.4 percent, while Wisconsin had the highest with 9.8 percent.

**Figure 72**  
**Nationally, Respondents Who Engage in Heavy Drinking, 2011**



Source: The Behavioral Risk Factor Surveillance System, South Dakota Department of Health, 2011

**Further Analysis**

The following data show the percent of those who engage in heavy drinking for various health behaviors and conditions. For example, 12 percent of respondents who currently smoke are heavy drinkers, while 4 percent of respondents who have never smoked are heavy drinkers.

<b>Table 48</b>				
<b>Heavy Drinking for Selected Health Behaviors and Conditions, 2011</b>				
<b>Health Behavior or Condition</b>	<b># Respondents</b>	<b>% Heavy Drinkers</b>	<b>95% CI</b>	
			<b>Lower</b>	<b>Upper</b>
Underweight (BMI < 18.5)	*	*	*	*
Recommended (BMI = 18.5-24.9)	2,355	7%	5%	10%
Overweight (BMI = 25.0-29.9)	2,823	6%	5%	8%
Obese - Class I (BMI = 30.0-34.9)	1,449	5%	3%	7%
Obese - Classes II & III (BMI = 35.0+)	747	4%	2%	9%
Leisure Time Exercise	5,668	6%	5%	7%
No Leisure Time Exercise	2,198	6%	4%	8%
Meet Aerobic Recommendations	3,673	6%	5%	8%
Did Not Meet Aerobic Recommendations	4,008	6%	5%	7%
At Least Five Servings of Fruits and Vegetables	1,025	2%	1%	4%
Less Than Five Servings of Fruits and Vegetables	6,814	6%	5%	8%
Current Smoker	1,372	12%	9%	16%
Former Smoker	2,372	5%	4%	7%
Never Smoked	4,101	4%	3%	5%
Smokeless Tobacco	368	11%	7%	18%
No Smokeless Tobacco	7,506	6%	5%	7%

**Table 48 (continued)  
Heavy Drinking for Selected Health Behaviors and Conditions, 2011**

Health Behavior or Condition	# Respondents	% Heavy Drinkers	95% CI	
			Lower	Upper
Diabetes	955	2%	1%	5%
No Diabetes	6,910	6%	5%	7%
High Blood Pressure	3,080	5%	4%	6%
No High Blood Pressure	4,785	6%	5%	8%
High Blood Cholesterol	2,773	5%	3%	7%
No High Blood Cholesterol	3,860	5%	4%	6%
Flu Shot (65+)	1,859	1%	1%	2%
No Flu Shot (65+)	773	4%	2%	7%
Pneumonia Shot (65+)	1,737	1%	1%	2%
No Pneumonia Shot (65+)	809	4%	2%	7%
Binge Drinker	1,176	23%	19%	28%
Not a Binge Drinker	6,677	1%	1%	1%
Always/Almost Always Wear Seat Belt	6,689	5%	4%	6%
Sometimes/Seldom/Never Wear Seat Belt	1,164	11%	8%	16%
Previously Had Heart Attack	543	3%	1%	9%
Never Had Heart Attack	7,297	6%	5%	7%
Have Angina or Coronary Heart Disease	520	1%	0%	2%
Do Not Have Angina or Coronary Heart Disease	7,294	6%	5%	7%
Previously Had a Stroke	295	2%	1%	6%
Never Had a Stroke	7,562	6%	5%	7%
Cancer	1,315	3%	2%	5%
No Cancer	6,530	6%	5%	7%
Skin Cancer	733	2%	1%	4%
No Skin Cancer	7,123	6%	5%	7%
Current Asthma	613	7%	3%	13%
Former Asthma	241	9%	4%	19%
Never Had Asthma	6,974	6%	5%	7%
Arthritis	2,488	4%	3%	5%
No Arthritis	5,349	7%	5%	8%
COPD	530	6%	2%	13%
No COPD	7,319	6%	5%	7%
Depression	1,273	7%	5%	11%
No Depression	6,576	6%	5%	7%
Kidney Disease	246	4%	2%	12%
No Kidney Disease	7,609	6%	5%	7%
Vision Impairment	1,694	6%	4%	10%
No Vision Impairment	6,143	6%	5%	7%
Good/Very Good/Excellent Health Status	6,559	6%	5%	7%
Fair/Poor Health Status	1,307	5%	3%	8%
Physical Health Not Good for 0-29 Days of the Past 30	7,210	6%	5%	7%
Physical Health Not Good for 30 Days of the Past 30	524	2%	1%	4%
Mental Health Not Good for 0-19 Days of the Past 30	7,357	6%	5%	7%
Mental Health Not Good for 20-30 Days of the Past 30	398	7%	3%	17%
Usual Activities Unattainable for 0-9 Days of the Past 30	7,244	6%	5%	7%
Usual Activities Unattainable for 10-30 Days of the Past 30	555	6%	3%	11%
Physical, Mental, or Emotional Problems	2,277	4%	3%	6%
No Physical, Mental, or Emotional Problems	5,580	7%	5%	8%
Disability with Special Equipment Needed	811	1%	1%	3%
No Disability with Special Equipment Needed	7,061	6%	5%	7%
Been Tested for HIV (18-64)	1,556	7%	5%	10%
Never Been Tested for HIV (18-64)	6,087	5%	4%	7%

Source: The Behavioral Risk Factor Surveillance System, South Dakota Department of Health, 2011