

# Overweight and Obese

## OVERWEIGHT OR OBESE

**Definition:** Overweight or obese is defined as having a Body Mass Index (BMI) of 25.0 or above. Body Mass Index (BMI) is calculated by taking a person's body weight in pounds, divided by their height in inches, divided by height in inches (again) times 703. The mathematical equation for BMI is:  $\text{weight (lb)}/\text{height (in)}^2 \times 703$ .


### Prevalence of Overweight or Obese

- South Dakota 64%
- Nationwide median 64%

### Healthy People 2020 Objective


There is no comparable Healthy People 2020 objective for overweight or obese.

**Figure 1**  
**Percent of Respondents Who Are Overweight or Obese Based on Body Mass Index, 1993-2010**



Source: The Behavioral Risk Factor Surveillance System, South Dakota Department of Health, 1993-2010

**Figure 2**  
**Percent of Respondents Who Are Overweight or Obese Based on Body Mass Index, 2011**



Note: Beginning in 2011, the CDC began using a different methodology to weight the data, therefore current data estimates cannot be compared to previous years.

Source: The Behavioral Risk Factor Surveillance System, South Dakota Department of Health, 2011


<b>Table 4</b>					
<b>Respondents Who Are Overweight or Obese, 2011</b>					
		<b>#</b>	<b>%</b>	<b>95% CI</b>	
		<b>Resp.</b>	<b>Overweight</b>	<b>Lower</b>	<b>Upper</b>
<b>Total</b>		<b>7,819</b>	<b>64%</b>	<b>62%</b>	<b>67%</b>
<b>Gender</b>	Male	3,280	73%	70%	76%
	Female	4,539	55%	52%	58%
<b>Age</b>	18-44	2,071	58%	54%	61%
	45-64	3,083	72%	69%	75%
	65+	2,640	68%	64%	71%
<b>Race</b>	White	6,717	64%	61%	66%
	American Indian	817	77%	70%	83%
<b>Ethnicity</b>	Hispanic	101	70%	54%	82%
	Non-Hispanic	7,705	64%	62%	66%
<b>Household Income</b>	Less than \$25,000	1,918	65%	61%	69%
	\$25,000-\$49,999	2,016	65%	61%	69%
	\$50,000+	2,792	69%	65%	72%
<b>Education</b>	High School, G.E.D., or Less	3,052	66%	62%	69%
	Some Post-High School	2,294	62%	58%	66%
	College Graduate	2,463	65%	62%	69%
<b>Employment Status</b>	Employed for Wages	3,436	64%	61%	67%
	Self-employed	864	70%	64%	75%
	Unemployed	328	69%	59%	77%
	Homemaker	416	53%	44%	62%
	Student	176	41%	31%	53%
	Retired	2,226	69%	66%	72%
<b>Marital Status</b>	Unable to Work	367	67%	57%	76%
	Married/Unmarried Couple	4,535	69%	66%	71%
	Divorced/Separated	1,049	65%	59%	70%
	Widowed	1,198	63%	58%	67%
	Never Married	1,028	54%	48%	59%
	<b>Home Ownership Status</b>	Own Home	5,958	68%	66%
Rent Home		1,574	61%	57%	66%
<b>Children Status</b>	Children in Household (Ages 18-44)	1,411	63%	58%	67%
	No Children in Household (Ages 18-44)	658	50%	44%	56%
<b>Phone Status</b>	Landline	6,875	66%	63%	68%
	Cell Phone	944	56%	52%	60%

Source: The Behavioral Risk Factor Surveillance System, South Dakota Department of Health, 2011

### **National Statistics**

The national median for respondents who are overweight or obese was 64 percent, as shown in Figure 2 on the next page. South Dakota had 64 percent of respondents who are overweight or obese. The District of Columbia had the lowest percent of respondents who are overweight or obese with 53 percent, while West Virginia had the highest percent with 69 percent.

**Figure 3**  
**Nationally, Respondents Who Are Overweight or Obese, 2011**



Source: The Behavioral Risk Factor Surveillance System, South Dakota Department of Health, 2011

**Further Analysis**

The following data show the percent of those who are overweight or obese for various health behaviors and conditions. For example, 84 percent of respondents who have diabetes are overweight or obese, while 62 percent of respondents who do not have diabetes are overweight or obese.

Health Behavior or Condition	# Respondents	% Overweight or Obese	95% CI	
			Lower	Upper
Leisure Time Exercise	5,533	62%	59%	64%
No Leisure Time Exercise	2,099	72%	68%	75%
Meet Aerobic Recommendations	3,612	61%	58%	64%
Did Not Meet Aerobic Recommendations	3,848	67%	64%	70%
At Least Five Servings of Fruits and Vegetables	983	57%	50%	62%
Less Than Five Servings of Fruits and Vegetables	6,655	65%	63%	68%
Current Smoker	1,421	59%	54%	64%
Former Smoker	2,367	73%	70%	76%
Never Smoked	4,005	62%	59%	65%
Smokeless Tobacco	398	73%	63%	81%
No Smokeless Tobacco	7,421	64%	62%	66%
Diabetes	953	84%	79%	88%
No Diabetes	6,858	62%	60%	65%

**Table 5 (continued)**  
**Overweight or Obese for Selected Health Behaviors and Conditions, 2011**

Health Behavior or Condition	# Respondents	% Overweight or Obese	95% CI	
			Lower	Upper
High Blood Pressure	3,064	79%	77%	82%
No High Blood Pressure	4,742	58%	55%	60%
High Blood Cholesterol	2,754	77%	74%	80%
No High Blood Cholesterol	3,820	64%	61%	67%
Flu Shot (65+)	1,802	71%	67%	75%
No Flu Shot (65+)	749	61%	54%	68%
Pneumonia Shot (65+)	1,692	69%	65%	73%
No Pneumonia Shot (65+)	774	64%	57%	70%
Drank Alcohol in Past 30 Days	4,100	65%	62%	67%
Did Not Drink Alcohol in Past 30 Days	3,432	64%	61%	68%
Binge Drinker	1,186	67%	62%	72%
Not a Binge Drinker	6,308	64%	62%	66%
Heavy Drinker	395	59%	49%	68%
Not a Heavy Drinker	7,077	65%	63%	67%
Always/Almost Always Wear Seat Belt	6,434	63%	61%	66%
Sometimes/Seldom/Never Wear Seat Belt	1,135	68%	63%	74%
Previously Had Heart Attack	553	78%	71%	83%
Never Had Heart Attack	7,234	64%	61%	66%
Have Angina or Coronary Heart Disease	528	77%	70%	83%
Do Not Have Angina or Coronary Heart Disease	7,225	64%	61%	66%
Previously Had a Stroke	294	64%	52%	74%
Never Had a Stroke	7,508	64%	62%	67%
Cancer	1,300	65%	60%	70%
No Cancer	6,491	64%	62%	67%
Skin Cancer	728	68%	62%	73%
No Skin Cancer	7,073	64%	62%	66%
Current Asthma	603	71%	64%	77%
Former Asthma	244	65%	52%	76%
Never Had Asthma	6,925	64%	61%	66%
Arthritis	2,454	73%	69%	76%
No Arthritis	5,329	62%	59%	64%
COPD	536	69%	61%	76%
No COPD	7,256	64%	62%	66%
Depression	1,253	64%	59%	69%
No Depression	6,540	65%	62%	67%
Kidney Disease	243	76%	66%	83%
No Kidney Disease	7,557	64%	62%	66%
Vision Impairment	1,699	63%	58%	68%
No Vision Impairment	6,082	65%	62%	67%
Good/Very Good/Excellent Health Status	6,499	63%	61%	66%
Fair/Poor Health Status	1,311	71%	66%	75%
Physical Health Not Good for 0-29 Days of the Past 30	7,161	64%	62%	67%
Physical Health Not Good for 30 Days of the Past 30	515	69%	61%	76%
Mental Health Not Good for 0-19 Days of the Past 30	7,286	64%	62%	67%
Mental Health Not Good for 20-30 Days of the Past 30	412	64%	55%	73%
Usual Activities Unattainable for 0-9 Days of the Past 30	7,190	64%	62%	66%
Usual Activities Unattainable for 10-30 Days of the Past 30	554	66%	58%	73%
Physical, Mental, or Emotional Problems	2,182	69%	65%	73%
No Physical, Mental, or Emotional Problems	5,405	63%	60%	65%
Disability with Special Equipment Needed	780	67%	59%	74%
No Disability with Special Equipment Needed	6,819	64%	62%	66%
Been Tested for HIV (18-64)	1,514	63%	58%	68%
Never Been Tested for HIV (18-64)	5,830	65%	62%	67%

Source: The Behavioral Risk Factor Surveillance System, South Dakota Department of Health, 2011

## OBESITY, CLASS I-III

**Definition: Obesity, Class I-III is defined as having a Body Mass Index (BMI) of 30.0 or greater. Body Mass Index (BMI) is calculated by taking a person's body weight in pounds divided by height in inches, divided by height in inches (again) times 703. The mathematical equation for BMI is:  $\text{weight (lb)}/\text{height (in)}^2 \times 703$ .**


### Prevalence of Obesity, Class I-III

- South Dakota 28%
- Nationwide median 28%

### South Dakota Department of Health 2020 Initiative


*Reduce the proportion of adults aged 18 years and older who are obese to 23 percent.*

**Figure 4**  
**Percent of Respondents Who Are Class I-III Obese Based on Body Mass Index, 1993-2010**



Source: The Behavioral Risk Factor Surveillance System, South Dakota Department of Health, 1993-2010

**Figure 5**  
**Percent of Respondents Who Are Class I-III Obese Based on Body Mass Index, 2011**



Note: Beginning in 2011, the CDC began using a different methodology to weight the data, therefore current data estimates cannot be compared to previous years.

Source: The Behavioral Risk Factor Surveillance System, South Dakota Department of Health, 2011

**Table 6  
Respondents Who Are Class I-III Obese, 2011**


		# Resp.	% Obese	95% CI	
				Lower	Upper
<b>Total</b>		<b>7,819</b>	<b>28%</b>	<b>26%</b>	<b>30%</b>
<b>Gender</b>	Male	3,280	30%	27%	33%
	Female	4,539	26%	24%	29%
<b>Age</b>	18-44	2,071	25%	23%	29%
	45-64	3,083	34%	31%	37%
	65+	2,640	24%	21%	27%
<b>Race</b>	White	6,717	27%	25%	29%
	American Indian	817	45%	37%	53%
<b>Ethnicity</b>	Hispanic	101	38%	25%	54%
	Non-Hispanic	7,705	28%	26%	30%
<b>Household Income</b>	Less than \$25,000	1,918	33%	29%	37%
	\$25,000-\$49,999	2,016	28%	24%	32%
	\$50,000+	2,792	27%	24%	30%
<b>Education</b>	High School, G.E.D., or Less	3,052	30%	27%	33%
	Some Post-High School	2,294	27%	24%	31%
	College Graduate	2,463	26%	23%	30%
<b>Employment Status</b>	Employed for Wages	3,436	29%	26%	32%
	Self-employed	864	27%	22%	33%
	Unemployed	328	43%	34%	53%
	Homemaker	416	25%	17%	35%
	Student	176	16%	10%	23%
	Retired	2,226	24%	21%	27%
	Unable to Work	367	40%	31%	49%
<b>Marital Status</b>	Married/Unmarried Couple	4,535	29%	26%	31%
	Divorced/Separated	1,049	32%	27%	37%
	Widowed	1,198	25%	21%	31%
	Never Married	1,028	26%	21%	31%
<b>Home Ownership Status</b>	Own Home	5,958	28%	26%	30%
	Rent Home	1,574	31%	27%	35%
<b>Children Status</b>	Children in Household (Ages 18-44)	1,411	27%	23%	31%
	No Children in Household (Ages 18-44)	658	23%	19%	29%
<b>Phone Status</b>	Landline	6,875	29%	27%	31%
	Cell Phone	944	24%	21%	27%

Source: The Behavioral Risk Factor Surveillance System, South Dakota Department of Health, 2011

### **National Statistics**

The national median for respondents who are obese was 28 percent, as shown in Figure 6 on the next page. South Dakota had 28 percent of respondents who are obese. Colorado had the lowest percent of respondents who are obese with 21 percent, while Mississippi had the highest percent with 35 percent.

**Figure 6**  
**Nationally, Respondents Who Are Class I-III Obese, 2011**



Source: The Behavioral Risk Factor Surveillance System, South Dakota Department of Health, 2011

**Further Analysis**

Following are data illustrating the percent of those who are Class I-III obese for various health behaviors and conditions. For example, 42 percent of respondents who have high blood pressure are Class I-III obese, while 22 percent of respondents who do not have high blood pressure are Class I-III obese.

Health Behavior or Condition	# Respondents	% Obese	95% CI	
			Lower	Upper
Leisure Time Exercise	5,533	27%	24%	29%
No Leisure Time Exercise	2,099	32%	28%	36%
Meet Aerobic Recommendations	3,612	25%	23%	28%
Did Not Meet Aerobic Recommendations	3,848	30%	28%	33%
At Least Five Servings of Fruits and Vegetables	983	24%	19%	30%
Less Than Five Servings of Fruits and Vegetables	6,655	28%	26%	31%
Current Smoker	1,421	27%	23%	32%
Former Smoker	2,367	34%	31%	38%
Never Smoked	4,005	26%	23%	28%
Smokeless Tobacco	398	31%	23%	40%
No Smokeless Tobacco	7,421	28%	26%	30%
Diabetes	953	53%	47%	60%
No Diabetes	6,858	25%	24%	27%
High Blood Pressure	3,064	42%	39%	45%
No High Blood Pressure	4,742	22%	20%	24%
High Blood Cholesterol	2,754	37%	34%	41%
No High Blood Cholesterol	3,820	28%	26%	31%

**Table 7 (continued)**  
**Class I-III Obese Status for Selected Health Behaviors and Conditions, 2011**

Health Behavior or Condition	# Respondents	% Obese	95% CI	
			Lower	Upper
Flu Shot (65+)	1,802	25%	21%	29%
No Flu Shot (65+)	749	20%	15%	26%
Pneumonia Shot (65+)	1,692	24%	20%	28%
No Pneumonia Shot (65+)	774	19%	14%	25%
Drank Alcohol in Past 30 Days	4,100	27%	24%	30%
Did Not Drink Alcohol in Past 30 Days	3,432	30%	27%	33%
Binge Drinker	1,186	27%	23%	32%
Not a Binge Drinker	6,308	28%	26%	31%
Heavy Drinker	395	21%	15%	28%
Not a Heavy Drinker	7,077	28%	26%	30%
Always/Almost Always Wear Seat Belt	6,434	27%	25%	29%
Sometimes/Seldom/Never Wear Seat Belt	1,135	33%	28%	38%
Previously Had Heart Attack	553	37%	30%	45%
Never Had Heart Attack	7,234	27%	25%	29%
Have Angina or Coronary Heart Disease	528	38%	31%	46%
Do Not Have Angina or Coronary Heart Disease	7,225	28%	26%	30%
Previously Had a Stroke	294	30%	21%	41%
Never Had a Stroke	7,508	28%	26%	30%
Cancer	1,300	26%	22%	31%
No Cancer	6,491	28%	26%	31%
Skin Cancer	728	25%	20%	31%
No Skin Cancer	7,073	28%	26%	30%
Current Asthma	603	40%	33%	47%
Former Asthma	244	37%	26%	50%
Never Had Asthma	6,925	27%	25%	29%
Arthritis	2,454	36%	32%	40%
No Arthritis	5,329	26%	23%	28%
COPD	536	36%	28%	45%
No COPD	7,256	28%	26%	30%
Depression	1,253	36%	31%	41%
No Depression	6,540	27%	25%	29%
Kidney Disease	243	44%	34%	55%
No Kidney Disease	7,557	28%	26%	30%
Vision Impairment	1,699	28%	24%	32%
No Vision Impairment	6,082	28%	26%	30%
Good/Very Good/Excellent Health Status	6,499	25%	23%	27%
Fair/Poor Health Status	1,311	44%	39%	49%
Physical Health Not Good for 0-29 Days of the Past 30	7,161	28%	26%	30%
Physical Health Not Good for 30 Days of the Past 30	515	38%	31%	46%
Mental Health Not Good for 0-19 Days of the Past 30	7,286	27%	25%	29%
Mental Health Not Good for 20-30 Days of the Past 30	412	40%	32%	50%
Usual Activities Unattainable for 0-9 Days of the Past 30	7,190	28%	26%	30%
Usual Activities Unattainable for 10-30 Days of the Past 30	554	35%	29%	43%
Physical, Mental, or Emotional Problems	2,182	37%	33%	41%
No Physical, Mental, or Emotional Problems	5,405	25%	23%	27%
Disability with Special Equipment Needed	780	38%	32%	45%
No Disability with Special Equipment Needed	6,819	27%	25%	29%
Been Tested for HIV (18-64)	1,514	30%	26%	35%
Never Been Tested for HIV (18-64)	5,830	27%	25%	30%

Source: The Behavioral Risk Factor Surveillance System, South Dakota Department of Health, 2011

## **OBESITY, CLASSES II-III**

**Definition: Obesity, Classes II-III is defined as having a Body Mass Index (BMI) of 35.0 or greater. Body Mass Index (BMI) is calculated by taking a person's body weight in pounds divided by height in inches, divided by height in inches (again) times 703. The mathematical equation for BMI is:  $\text{weight (lb)}/\text{height (in)}^2 \times 703$ .**


### **Prevalence of Obesity, Classes II-III**

- South Dakota 9%
- There is no nationwide median for Obesity classes II-III

### **Healthy People 2020 Objective**


There is no comparable Healthy People 2020 objective for obesity, classes II-III.

**Figure 7**  
**Percent of Respondents Who Are Class II-III Obese Based on Body Mass Index, 2002-2010**



Source: The Behavioral Risk Factor Surveillance System, South Dakota Department of Health, 2002-2010

**Figure 8**  
**Percent of Respondents Who Are Class II-III Obese Based on Body Mass Index, 2011**



Note: Beginning in 2011, the CDC began using a different methodology to weight the data, therefore current data estimates cannot be compared to previous years.

Source: The Behavioral Risk Factor Surveillance System, South Dakota Department of Health, 2011

		# Resp.	% Obese	95% CI	
				Lower	Upper
<b>Total</b>		<b>7,819</b>	<b>9%</b>	<b>8%</b>	<b>11%</b>
<b>Gender</b>	Male	3,280	9%	7%	10%
	Female	4,539	10%	8%	12%
<b>Age</b>	18-44	2,071	9%	7%	11%
	45-64	3,083	11%	9%	13%
	65+	2,640	7%	5%	10%
<b>Race</b>	White	6,717	9%	7%	10%
	American Indian	817	16%	10%	24%
<b>Ethnicity</b>	Hispanic	101	13%	5%	29%
	Non-Hispanic	7,705	9%	8%	11%
<b>Household Income</b>	Less than \$25,000	1,918	11%	9%	14%
	\$25,000-\$49,999	2,016	11%	9%	14%
	\$50,000+	2,792	7%	6%	9%
<b>Education</b>	High School, G.E.D., or Less	3,052	11%	9%	13%
	Some Post-High School	2,294	9%	7%	11%
	College Graduate	2,463	7%	6%	9%
<b>Employment Status</b>	Employed for Wages	3,436	9%	8%	11%
	Self-employed	864	7%	4%	11%
	Unemployed	328	16%	10%	24%
	Homemaker	416	12%	6%	22%
	Student	176	4%	2%	9%
	Retired	2,226	8%	6%	10%
	Unable to Work	367	18%	13%	25%
<b>Marital Status</b>	Married/Unmarried Couple	4,535	9%	8%	11%
	Divorced/Separated	1,049	11%	8%	15%
	Widowed	1,198	6%	4%	10%
	Never Married	1,028	10%	7%	13%
<b>Home Ownership Status</b>	Own Home	5,958	9%	8%	10%
	Rent Home	1,574	11%	9%	15%
<b>Children Status</b>	Children in Household (Ages 18-44)	1,411	9%	7%	13%
	No Children in Household (Ages 18-44)	658	9%	6%	12%
<b>Phone Status</b>	Landline	6,875	9%	8%	11%
	Cell Phone	944	9%	7%	11%

Source: The Behavioral Risk Factor Surveillance System, South Dakota Department of Health, 2011

### **Further Analysis**

Following are data illustrating the percent of those who are obese for various health behaviors and conditions. For example, 21 percent of respondents who have diabetes are Class II-III obese, while 8 percent of respondents who do not have diabetes are Class II-III obese.

<b>Health Behavior or Condition</b>	<b># Respondents</b>	<b>% Obese</b>	95% CI	
			<b>Lower</b>	<b>Upper</b>
Leisure Time Exercise	5,533	9%	7%	10%
No Leisure Time Exercise	2,099	11%	9%	14%
Meet Aerobic Recommendations	3,612	8%	6%	10%
Did Not Meet Aerobic Recommendations	3,848	11%	9%	13%
At Least Five Servings of Fruits and Vegetables	983	8%	6%	12%
Less Than Five Servings of Fruits and Vegetables	6,655	9%	8%	11%
Current Smoker	1,421	11%	8%	15%
Former Smoker	2,367	11%	9%	13%
Never Smoked	4,005	8%	6%	9%
Smokeless Tobacco	398	8%	4%	13%
No Smokeless Tobacco	7,421	9%	8%	11%

**Table 9 (continued)**  
**Class II-III Obese Status for Selected Health Behaviors and Conditions, 2011**

Health Behavior or Condition	# Respondents	% Obese	95% CI	
			Lower	Upper
Diabetes	953	21%	17%	26%
No Diabetes	6,858	8%	7%	9%
High Blood Pressure	3,064	15%	12%	17%
No High Blood Pressure	4,742	7%	6%	9%
High Blood Cholesterol	2,754	13%	10%	15%
No High Blood Cholesterol	3,820	9%	7%	11%
Flu Shot (65+)	1,802	8%	6%	11%
No Flu Shot (65+)	749	6%	3%	11%
Pneumonia Shot (65+)	1,692	8%	6%	11%
No Pneumonia Shot (65+)	774	7%	4%	12%
Drank Alcohol in Past 30 Days	4,100	8%	7%	10%
Did Not Drink Alcohol in Past 30 Days	3,432	11%	9%	13%
Binge Drinker	1,186	11%	8%	15%
Not a Binge Drinker	6,308	9%	8%	10%
Heavy Drinker	395	6%	3%	13%
Not a Heavy Drinker	7,077	9%	8%	11%
Always/Almost Always Wear Seat Belt	6,434	9%	7%	10%
Sometimes/Seldom/Never Wear Seat Belt	1,135	13%	9%	17%
Previously Had Heart Attack	553	14%	9%	21%
Never Had Heart Attack	7,234	9%	8%	10%
Have Angina or Coronary Heart Disease	528	13%	9%	19%
Do Not Have Angina or Coronary Heart Disease	7,225	9%	8%	11%
Previously Had a Stroke	294	8%	4%	15%
Never Had a Stroke	7,508	9%	8%	11%
Cancer	1,300	9%	6%	12%
No Cancer	6,491	9%	8%	11%
Skin Cancer	728	7%	5%	11%
No Skin Cancer	7,073	9%	8%	11%
Current Asthma	603	15%	11%	20%
Former Asthma	244	19%	10%	33%
Never Had Asthma	6,925	8%	7%	10%
Arthritis	2,454	12%	10%	15%
No Arthritis	5,329	8%	7%	10%
COPD	536	15%	10%	20%
No COPD	7,256	9%	8%	10%
Depression	1,253	13%	11%	16%
No Depression	6,540	9%	7%	10%
Kidney Disease	243	23%	14%	34%
No Kidney Disease	7,557	9%	8%	10%
Vision Impairment	1,699	10%	8%	13%
No Vision Impairment	6,082	9%	8%	11%
Good/Very Good/Excellent Health Status	6,499	8%	7%	10%
Fair/Poor Health Status	1,311	16%	13%	20%
Physical Health Not Good for 0-29 Days of the Past 30	7,161	9%	8%	10%
Physical Health Not Good for 30 Days of the Past 30	515	15%	11%	21%
Mental Health Not Good for 0-19 Days of the Past 30	7,286	9%	8%	10%
Mental Health Not Good for 20-30 Days of the Past 30	412	19%	12%	27%
Usual Activities Unattainable for 0-9 Days of the Past 30	7,190	9%	8%	10%
Usual Activities Unattainable for 10-30 Days of the Past 30	554	15%	11%	21%
Physical, Mental, or Emotional Problems	2,182	14%	11%	17%
No Physical, Mental, or Emotional Problems	5,405	8%	7%	9%
Disability with Special Equipment Needed	780	19%	15%	25%
No Disability with Special Equipment Needed	6,819	9%	7%	10%
Been Tested for HIV (18-64)	1,514	12%	9%	16%
Never Been Tested for HIV (18-64)	5,830	9%	7%	10%

Source: The Behavioral Risk Factor Surveillance System, South Dakota Department of Health, 2011