


# Map 1

## Percent of Low Birth Weight Infants by County, 2008-2012

U.S. = 8.1%                      South Dakota = 6.3%



- Denotes that the county's percent is significantly lower than the state percent
- Denotes that the county's percent is significantly higher than the state percent

Note: "Significantly" refers to statistical significance at the 0.05 level. Although a county's actual percent may be higher or lower than the state percent, the difference may not be statistically significant due to the small number of people in the county. The percent of low birth weight infants is calculated based on the first weight of the newborn obtained after birth. Low birth weight infants are those born alive who weigh less than 2,500 grams (about 5 pounds 9 ounces). The U.S. percent of low birth weight infants is from 2011. See technical notes for more complete explanations.


Source: South Dakota Department of Health, Office of Health Statistics.



### Map 3

## Percent of Mothers Who Used Tobacco While Pregnant by County, 2008-2012

**U.S. = 9.7%\***                      **South Dakota = 17.5%**




● Denotes that the county's percent is significantly lower than the state percent.  
 ○ Denotes that the county's percent is significantly higher than the state percent.  
 Note: "Significantly" refers to statistical significance at the 0.05 level. Although a county's actual percent may be higher or lower than the state percent, the difference may not be statistically significant due to the small number of people in the county. The data for mothers who used tobacco while pregnant are self-reported on the birth certificates. The U.S. percent of mothers who used tobacco while pregnant is from 2008 and is based on 24 states using the 2003 standard birth certificate, including South Dakota. See technical notes for more complete explanations.  
 \*Due to lack of comparable data from all 50 states, U.S. and South Dakota cannot be compared.  
 Source: South Dakota Department of Health, Office of Health Statistics.

## Map 4

### Percent of Births Less Than 37 Weeks Gestation by County, 2008-2012

**U.S. = 11.7%\***                      **South Dakota = 8.2%**




● Denotes that the county's rate is significantly lower than the state rate.  
 ○ Denotes that the county's rate is significantly higher than the state rate.  
 Note: "Significantly" refers to statistical significance at the 0.05 level. Although a county's actual percent may be higher or lower than the state percent, the difference may not be statistically significant due to the small number of people in the county. The U.S. percent of births less than 37 weeks gestation is from 2011. See technical notes for more complete explanations.  
 \*U.S. and South Dakota data cannot be compared.  
 Source: South Dakota Department of Health, Office of Health Statistics.

# Map 5

## Average Age of Mother by Resident County, 2008-2012

U.S. = 25.6\*      South Dakota = 27.1




● Denotes that the county's age is significantly lower than the state age.  
 ○ Denotes that the county's age is significantly higher than the state age.  
 Note: "Significantly" refers to statistical significance at the 0.05 level. Although a county's actual age may be higher or lower than the state age, the difference may not be statistically significant due to the small number of people in the county. The U.S. average age of mothers is from 2011.  
 \*Due to lack of comparable data from all 50 states, U.S. and South Dakota can not be compared. See technical notes for more complete explanations.  
 Source: South Dakota Department of Health, Office of Health Statistics.



# Map 7

## Death Rate Due to All Causes by County, 2008-2012

**U.S. = 747.0**                      **South Dakota = 707.3**




● Denotes that the county's rate is significantly lower than the state rate.  
 ○ Denotes that the county's rate is significantly higher than the state rate.  
 Note: "Significantly" refers to statistical significance at the 0.05 level. Although a county's actual rate may be higher or lower than the state rate, the difference may not be statistically significant due to the small number of people in the county. The death rate is age-adjusted per 100,000 population. This eliminates age difference between populations, making them easier to compare. The U.S. age-adjusted death rate is from 2010. See technical notes for more complete explanations.  
 Source: South Dakota Department of Health, Office of Health Statistics.

## Map 8

### Death Rate Due to Heart Disease by County, 2008-2012

**U.S. = 179.1**                      **South Dakota = 158.4**




• Denotes that the county's rate is significantly lower than the state rate.  
 ○ Denotes that the county's rate is significantly higher than the state rate.  
 Note: "Significantly" refers to statistical significance at the 0.05 level. Although a county's actual rate may be higher or lower than the state rate, the difference may not be statistically significant due to the small number of people in the county. The death rate is age-adjusted per 100,000 population. This eliminates age difference between populations, making them easier to compare. Heart disease is defined as ICD-10 codes I00-I09, I11, I13, and I20-I51. The U.S. age-adjusted Heart Disease death rate is from 2010. See technical notes for more complete explanations.  
 Source: South Dakota Department of Health, Office of Health Statistics.

## Map 9

### Death Rate Due to All Malignant Neoplasms (Cancer) by County, 2008-2012

U.S. = 172.8      South Dakota = 164.5




• Denotes that the county's rate is significantly lower than the state rate.  
 ○ Denotes that the county's rate is significantly higher than the state rate.  
 Note: "Significantly" refers to statistical significance at the 0.05 level. Although a county's actual rate may be higher or lower than the state rate, the difference may not be statistically significant due to the small number of people in the county. The death rate is age-adjusted per 100,000 population. This eliminates age difference between populations, making them easier to compare. Malignant Neoplasms (Cancer) are defined as ICD-10 codes C00-C97. The U.S. age-adjusted Malignant Neoplasms (Cancer) death rate is from 2010. See technical notes for more complete explanations.  
 Source: South Dakota Department of Health, Office of Health Statistics.



# Map 11

## Death Rate Due to Colorectal Cancer by County, 2008-2012

**U.S. = 15.8**                      **South Dakota = 15.8**




● Denotes that the county's rate is significantly lower than the state rate.  
 ○ Denotes that the county's rate is significantly higher than the state rate.  
 Note: "Significantly" refers to statistical significance at the 0.05 level. Although a county's actual rate may be higher or lower than the state rate, the difference may not be statistically significant due to the small number of people in the county. The death rate is age-adjusted per 100,000 population. This eliminates age difference between populations, making them easier to compare. Colorectal Cancer is defined as ICD-10 codes C18-C21. The U.S. age-adjusted Colorectal Cancer death rate is from 2010. See technical notes for more complete explanations.  
 Source: South Dakota Department of Health, Office of Health Statistics.

## Map 12

### Death Rate Due to Female Breast Cancer by County, 2008-2012

**U.S. = 22.1**                      **South Dakota = 20.7**



- Denotes that the county's rate is significantly lower than the state rate.
- Denotes that the county's rate is significantly higher than the state rate.

Note: "Significantly" refers to statistical significance at the 0.05 level. Although a county's actual rate may be higher or lower than the state rate, the difference may not be statistically significant due to the small number of people in the county. The death rate is age-adjusted per 100,000 population. This eliminates age difference between populations, making them easier to compare. Female Breast Cancer is defined as ICD-10 code C50. The U.S. age-adjusted Female Breast Cancer death rate is from 2010. See technical notes for more complete explanations.

Source: South Dakota Department of Health, Office of Health Statistics.










# Map 17

## Death Rate Due to Motor Vehicle Accidents by County, 2008-2012

**U.S. = 11.3**                      **South Dakota = 15.5**




● Denotes that the county's rate is significantly lower than the state rate.  
 ○ Denotes that the county's rate is significantly higher than the state rate.  
 Note: "Significantly" refers to statistical significance at the 0.05 level. Although a county's actual rate may be higher or lower than the state rate, the difference may not be statistically significant due to the small number of people in the county. The death rate is age-adjusted per 100,000 population. This eliminates age difference between populations, making them easier to compare. Motor Vehicle Accidents are defined as ICD-10 codes (V02-V04, V09.0, V09.2, V12-V14, V19.0-V19.2, V19.4-V19.6, V20-V79, V80.3-V80.5, V81.0-V81.1, V82.0-V82.1, V83-V86, V87.0-V87.8, V88.0-V88.8, V89.0, and V89.2) The U.S. age-adjusted Motor Vehicle Accident death rate is from 2010. See technical notes for more complete explanations.  
 Source: South Dakota Department of Health, Office of Health Statistics.



# Map 19

## Death Rate Due to Diabetes Mellitus by County, 2008-2012

U.S. = 20.8 South Dakota = 23.0



● Denotes that the county's rate is significantly lower than the state rate.  
 ○ Denotes that the county's rate is significantly higher than the state rate.  
 Note: "Significantly" refers to statistical significance at the 0.05 level. Although a county's actual rate may be higher or lower than the state rate, the difference may not be statistically significant due to the small number of people in the county. The death rate is age-adjusted per 100,000 population. This eliminates age difference between populations, making them easier to compare. Diabetes Mellitus is defined as ICD-10 codes E10-E14. The U.S. age-adjusted Diabetes Mellitus death rate is from 2010. See technical notes for more complete explanations.  
 Source: South Dakota Department of Health, Office of Health Statistics.








# Map 23

## Infant Mortality Rate by County, 2008-2012

**U.S. = 6.15**                      **South Dakota = 7.40**



● Denotes that the county's rate is significantly lower than the state rate.  
 ○ Denotes that the county's rate is significantly higher than the state rate.  
 Note: "Significantly" refers to statistical significance at the 0.05 level. Although a county's actual rate may be higher or lower than the state rate, the difference may not be statistically significant due to the small number of people in the county. Infant mortality is calculated as the number of deaths to babies (less than 1 year old) per 1,000 live births. The U.S. infant mortality rate is from 2010. See technical notes for more complete explanations. Source: South Dakota Department of Health, Office of Health Statistics.

