

# Tobacco Use

## CIGARETTE SMOKING

**Definition:** Respondents who report having smoked at least 100 cigarettes in their lifetime and now smoke every day or smoke some days.


### Prevalence of Current Cigarette Smoking

- South Dakota 20%
- Nationwide median 19%

### South Dakota Department of Health 2020 Initiative


Reduce the percent of adults who smoke cigarettes to 19 percent.

**Figure 18**  
**Percent of Respondents Who Currently Smoke Cigarettes, 1987-2010**



Source: The Behavioral Risk Factor Surveillance System, South Dakota Department of Health, 1987-2010

**Figure 19**  
**Percent of Respondents Who Currently Smoke Cigarettes, 2011-2013**



Note: Beginning in 2011, the CDC began using a different methodology to weight the data, therefore current data estimates cannot be compared to previous years.

Source: The Behavioral Risk Factor Surveillance System, South Dakota Department of Health, 2011-2013

**Table 20**  
**Respondents Who Currently Smoke Cigarettes, 2011-2013**

		2011-2013	2011	2012	2013
<b>Total</b>		<b>21%</b>	<b>23%</b>	<b>22%</b>	<b>20%</b>
<b>Gender</b>	Male	23%	24%	25%	20%
	Female	20%	22%	19%	19%
<b>Age</b>	18-34	31%	*	*	*
	35-44	22%	*	*	*
	45-54	24%	*	*	*
	55-64	17%	19%	20%	*
	65-74	13%	*	*	*
	75+	5%	*	*	*
<b>Race</b>	White	19%	21%	19%	17%
	American Indian	48%	*	*	*
<b>Ethnicity</b>	Hispanic	24%	*	*	*
	Non-Hispanic	21%	23%	22%	20%
<b>Household Income</b>	Less than \$15,000	38%	*	*	*
	\$15,000-\$24,999	31%	*	*	*
	\$25,000-\$34,999	26%	*	*	*
	\$35,000-\$49,999	21%	*	*	*
	\$50,000-\$74,999	17%	*	*	*
	\$75,000+	11%	10%	13%	*
<b>Education</b>	Less than High School, G.E.D.	33%	*	*	*
	High School, G.E.D.	27%	28%	29%	23%
	Some Post-High School	21%	24%	20%	19%
	College Graduate	10%	10%	10%	9%
<b>Employment Status</b>	Employed for Wages	23%	24%	24%	20%
	Self-employed	17%	*	*	*
	Unemployed	51%	*	*	*
	Homemaker	20%	*	*	*
	Student	17%	*	*	*
	Retired	8%	9%	9%	7%
	Unable to Work	44%	*	*	*
<b>Marital Status</b>	Married/Unmarried Couple	17%	18%	17%	15%
	Divorced/Separated	38%	*	*	*
	Widowed	14%	*	*	*
	Never Married	30%	*	*	*
<b>Home Ownership Status</b>	Own Home	16%	18%	17%	14%
	Rent Home	36%	37%	36%	*
<b>Children Status</b>	Children in Household (Ages 18-44)	27%	29%	26%	*
	No Children in Household (Ages 18-44)	29%	*	*	*
<b>Phone Status</b>	Landline	17%	21%	17%	12%
	Cell Phone	29%	*	30%	27%
<b>Pregnancy Status</b>	Pregnant (Ages 18-44)	18%	*	*	*
	Not Pregnant (Ages 18-44)	28%	*	*	*


Note: \*Results based on small sample sizes have been suppressed.

Source: The Behavioral Risk Factor Surveillance System, South Dakota Department of Health, 2011-2013

### **National Statistics**

The national median for respondents who currently smoke cigarettes was 19.0 percent, as shown in Figure 20 on the next page. South Dakota had 19.6 percent of respondents who currently smoke cigarettes. Utah had the lowest percent of respondents who currently smoke cigarettes with 10.3 percent, while West Virginia had the highest with 27.3 percent.

**Figure 20**  
**Nationally, Respondents Who Currently Smoke Cigarettes, 2013**



Source: The Behavioral Risk Factor Surveillance System, South Dakota Department of Health, 2013

**Further Analysis**

The following data show the percent of those who currently smoke cigarettes for various health behaviors and conditions. For example, 44 percent of respondents who are heavy drinkers currently smoke cigarettes, while 20 percent of respondents who are not heavy drinkers currently smoke cigarettes.

<b>Health Behavior or Condition</b>	<b>% Currently Smoke Cigarettes</b>
Underweight (BMI < 18.5)	34%
Recommended (BMI = 18.5-24.9)	25%
Overweight (BMI = 25.0-29.9)	19%
Obese - Class I (BMI = 30.0-34.9)	19%
Obese - Classes II & III (BMI = 35.0+)	23%
Leisure Time Exercise	19%
No Leisure Time Exercise	27%
Met Physical Activity Recommendations	19%
Did Not Meet Physical Activity Recommendations	23%
2+ Servings of Fruit per Day	13%
< 2 Servings of Fruit per Day	24%
3+ Servings of Vegetables per Day	17%
< 3 Servings of Vegetables per Day	22%
5+ Servings of Fruits and Vegetables per Day	13%
< 5 Servings of Fruits and Vegetables per Day	22%

**Table 21 (continued)  
Currently Smoke Cigarettes for Selected Health Behaviors and  
Conditions, 2011-2013**

<b>Health Behavior or Condition</b>	<b>% Currently Smoke Cigarettes</b>
Smokeless Tobacco	29%
No Smokeless Tobacco	21%
Heard of "South Dakota QuitLine"	21%
Not Heard of "South Dakota QuitLine"	11%
Diabetes	20%
No Diabetes	22%
High Blood Pressure	18%
No High Blood Pressure	23%
High Blood Cholesterol	17%
No High Blood Cholesterol	18%
Health Insurance (18-64)	21%
No Health Insurance (18-64)	44%
Employer Based Coverage (18-64)	17%
Private Health Plan (18-64)	12%
Medicare (18-64)	38%
Medicaid or Medical Assistance (18-64)	49%
The Military, CHAMPUS, TriCare, or the VA (18-64)	28%
The Indian Health Service (18-64)	54%
Flu Shot (65+)	8%
No Flu Shot (65+)	11%
Pneumonia Shot (65+)	8%
No Pneumonia Shot (65+)	10%
Previously Had Heart Attack	20%
Never Had Heart Attack	21%
Have Angina or Coronary Heart Disease	18%
Do Not Have Angina or Coronary Heart Disease	22%
Previously Had a Stroke	22%
Never Had a Stroke	21%
Cancer	16%
No Cancer	22%
Skin Cancer	11%
No Skin Cancer	22%
Current Asthma	25%
Former Asthma	25%
Never Had Asthma	21%
Arthritis	21%
No Arthritis	22%
COPD	36%
No COPD	21%
Depression	35%
No Depression	19%
Kidney Disease	22%
No Kidney Disease	21%
Severe Vision Impairment	24%
No Severe Vision Impairment	19%
Recent Increased Confusion/Memory Loss	32%
No Recent Increased Confusion/Memory Loss	18%
< 6 Hours of Sleep	45%
6+ Hours of Sleep	17%

<b>Table 21 (continued)</b>	
<b>Currently Smoke Cigarettes for Selected Health Behaviors and Conditions, 2011-2013</b>	
<b>Health Behavior or Condition</b>	<b>% Currently Smoke Cigarettes</b>
Always/Almost Always Wear Seat Belt	18%
Sometimes/Seldom/Never Wear Seat Belt	35%
Drank Alcohol in Past 30 Days	23%
Did Not Drink Alcohol in Past 30 Days	19%
Binge Drinker	37%
Not a Binge Drinker	17%
Heavy Drinker	44%
Not a Heavy Drinker	20%
Good/Very Good/Excellent Health Status	20%
Fair/Poor Health Status	30%
Physical Health Not Good for 0-29 Days of the Past 30	21%
Physical Health Not Good for 30 Days of the Past 30	30%
Mental Health Not Good for 0-19 Days of the Past 30	20%
Mental Health Not Good for 20-30 Days of the Past 30	45%
Usual Activities Unattainable for 0-9 Days of the Past 30	20%
Usual Activities Unattainable for 10-30 Days of the Past 30	36%
Physical, Mental, or Emotional Problems	27%
No Physical, Mental, or Emotional Problems	20%
Disability with Special Equipment Needed	21%
No Disability with Special Equipment Needed	22%
< 2 Hours of TV per Day	17%
2+ Hours of TV per Day	20%
Been Tested for HIV (18-64)	33%
Never Been Tested for HIV (18-64)	17%

Source: The Behavioral Risk Factor Surveillance System, South Dakota Department of Health, 2011-2013

In 2012-2013, 55 percent of respondents tried to stop smoking for one day or longer because they were trying to quit smoking as shown below in Table 22.

<b>Table 22</b>		
<b>Respondents Who Tried to Stop Smoking, Within the Past 12 Months, for One Day or Longer Because They Were Trying to Quit Smoking, 2011-2013</b>		
<b>Survey Year</b>	<b># of Respondents</b>	<b>Percent</b>
2012-2013	2,638	55%
2011-2012	2,909	56%

Source: The Behavioral Risk Factor Surveillance System, South Dakota Department of Health, 2011-2013


Table 23, on the next page, shows the percent of current cigarette smokers for 2011-2013 by the type of health insurance they have. Those with Indian Health Service coverage had the highest percent of current smokers with 54 percent. This was followed by Medicaid or medical assistance with 49 percent and Medicare with 38 percent.

<b>Table 23 Percent of Current Cigarette Smoking by Type of Health Insurance, Ages 18-64, 2011-2013</b>	
<b>Type of Health Insurance</b>	<b>2011-2013</b>
The Indian Health Service	54%
Medicaid or Medical Assistance	49%
Medicare	38%
The Military, CHAMPUS, TriCare, or the VA	28%
Employer Based Coverage	17%
Private Health Insurance Plan	12%
None	44%

Source: The Behavioral Risk Factor Surveillance System, South Dakota Department of Health, 2011-2013

Figure 21, below, shows the percent of respondents who have been advised to quit smoking in the past 12 months by a health professional. In 2011-2013, 70 percent of respondents said they have been advised to quit smoking by a health professional.


**Figure 21  
Percent of Respondents Who Have Been Advised by a Doctor, Nurse, or Other Health Professional to Quit Smoking in the Past 12 Months, 2011-2013**



Source: The Behavioral Risk Factor Surveillance System, South Dakota Department of Health, 2011-2013

Figure 22, on the next page, shows the respondents' place of work official smoking policy for work areas. The majority of respondents for all three years stated that smoking was not allowed in any work areas.


**Figure 22**  
**Respondents' Place of Work Smoking Policy, 2011-2013**



Source: The Behavioral Risk Factor Surveillance System, South Dakota Department of Health, 2011-2013

Figure 23, below, shows the respondents' rules about smoking inside their homes. The majority of respondents for all three years stated that smoking was not allowed anywhere in their homes.

**Figure 23**  
**Respondents' Rules About Smoking Inside the Home, 2011-2013**



Source: The Behavioral Risk Factor Surveillance System, South Dakota Department of Health, 2011-2013

## **SMOKELESS TOBACCO**

**Definition:** Respondents who report that they use chewing tobacco or snuff every day or some days.


### **Prevalence of Smokeless Tobacco**

- South Dakota 7%
- There is no nationwide median for smokeless tobacco.

### **Healthy People 2020 Objective**


There was no stated Healthy People 2020 Objective for smokeless tobacco use.

**Figure 24**  
**Percent of Respondents Who Use Smokeless Tobacco,**  
**1990, 1993-1996, 2001, 2003, and 2005-2010**



Source: The Behavioral Risk Factor Surveillance System, South Dakota Department of Health, 1990, 1993-1996, 2001, 2003, and 2005-2010

**Figure 25**  
**Percent of Respondents Who Use Smokeless Tobacco, 2011-2013**



Note: Beginning in 2011, the CDC began using a different methodology to weight the data, therefore current data estimates cannot be compared to previous years.

Source: The Behavioral Risk Factor Surveillance System, South Dakota Department of Health, 2011-2013

**Table 24**  
**Respondents Who Use Smokeless Tobacco, 2011-2013**

		2011-2013	2011	2012	2013
<b>Total</b>		<b>7%</b>	<b>7%</b>	<b>6%</b>	<b>7%</b>
<b>Gender</b>	Male	12%	12%	12%	12%
	Female	1%	1%	1%	1%
<b>Age</b>	18-34	10%	*	*	*
	35-44	10%	*	*	*
	45-54	5%	*	*	*
	55-64	4%	3%	2%	*
	65-74	3%	*	*	*
	75+	2%	*	*	*
<b>Race</b>	White	6%	6%	6%	7%
	American Indian	9%	*	*	*
<b>Ethnicity</b>	Hispanic	5%	*	*	*
	Non-Hispanic	7%	7%	6%	7%
<b>Household Income</b>	Less than \$15,000	5%	*	*	*
	\$15,000-\$24,999	7%	*	*	*
	\$25,000-\$34,999	6%	*	*	*
	\$35,000-\$49,999	7%	*	*	*
	\$50,000-\$74,999	8%	*	*	*
	\$75,000+	6%	6%	7%	*
<b>Education</b>	Less than High School, G.E.D.	9%	*	*	*
	High School, G.E.D.	7%	7%	7%	8%
	Some Post-High School	7%	8%	6%	7%
	College Graduate	4%	5%	4%	4%
<b>Employment Status</b>	Employed for Wages	8%	8%	8%	8%
	Self-employed	9%	*	*	*
	Unemployed	8%	*	*	*
	Homemaker	1%	*	*	*
	Student	8%	*	*	*
	Retired	2%	2%	2%	2%
	Unable to Work	5%	*	*	*
<b>Marital Status</b>	Married/Unmarried Couple	6%	6%	6%	6%
	Divorced/Separated	8%	*	*	*
	Widowed	1%	*	*	*
	Never Married	10%	*	*	*
<b>Home Ownership Status</b>	Own Home	6%	6%	6%	6%
	Rent Home	8%	9%	7%	*
<b>Children Status</b>	Children in Household (Ages 18-44)	9%	8%	9%	*
	No Children in Household (Ages 18-44)	12%	*	*	*
<b>Phone Status</b>	Landline	5%	6%	4%	5%
	Cell Phone	9%	*	10%	8%
<b>Pregnancy Status</b>	Pregnant (Ages 18-44)	0.2%	*	*	*
	Not Pregnant (Ages 18-44)	2%	*	*	*

Note: \*Results based on small sample sizes have been suppressed.

Source: The Behavioral Risk Factor Surveillance System, South Dakota Department of Health, 2011-2013

### **Further Analysis**

The following data show the percent of those who currently use smokeless tobacco for various health behaviors and conditions. For example, 13 percent of respondents who are binge drinkers use smokeless tobacco, while five percent of respondents who are not binge drinkers use smokeless tobacco.

**Table 25  
Smokeless Tobacco Use for Selected Health Behaviors and  
Conditions, 2011-2013**

<b>Health Behavior or Condition</b>	<b>% Use Smokeless Tobacco</b>
Underweight (BMI < 18.5)	3%
Recommended (BMI = 18.5-24.9)	5%
Overweight (BMI = 25.0-29.9)	8%
Obese - Class I (BMI = 30.0-34.9)	9%
Obese - Classes II & III (BMI = 35.0+)	7%
Leisure Time Exercise	6%
No Leisure Time Exercise	8%
Met Physical Activity Recommendations	6%
Did Not Meet Physical Activity Recommendations	7%
2+ Servings of Fruit per Day	3%
< 2 Servings of Fruit per Day	8%
3+ Servings of Vegetables per Day	3%
< 3 Servings of Vegetables per Day	7%
5+ Servings of Fruits and Vegetables per Day	2%
< 5 Servings of Fruits and Vegetables per Day	7%
Current Smoker	9%
Former Smoker	7%
Never Smoked	5%
Heard of "South Dakota QuitLine"	7%
Not Heard of "South Dakota QuitLine"	5%
Diabetes	4%
No Diabetes	7%
High Blood Pressure	6%
No High Blood Pressure	7%
High Blood Cholesterol	5%
No High Blood Cholesterol	6%
Health Insurance (18-64)	7%
No Health Insurance (18-64)	9%
Employer Based Coverage (18-64)	6%
Private Health Plan (18-64)	10%
Medicare (18-64)	7%
Medicaid or Medical Assistance (18-64)	5%
The Military, CHAMPUS, TriCare, or the VA (18-64)	9%
The Indian Health Service (18-64)	13%
Flu Shot (65+)	1%
No Flu Shot (65+)	4%
Pneumonia Shot (65+)	2%
No Pneumonia Shot (65+)	2%
Previously Had Heart Attack	4%
Never Had Heart Attack	7%
Have Angina or Coronary Heart Disease	4%
Do Not Have Angina or Coronary Heart Disease	7%
Previously Had a Stroke	3%
Never Had a Stroke	7%
Cancer	3%
No Cancer	7%
Skin Cancer	4%
No Skin Cancer	7%
Current Asthma	9%
Former Asthma	8%
Never Had Asthma	6%


**Table 25 (continued)  
Smokeless Tobacco Use for Selected Health Behaviors and  
Conditions, 2011-2013**

<b>Health Behavior or Condition</b>	<b>% Use Smokeless Tobacco</b>
Arthritis	6%
No Arthritis	7%
COPD	7%
No COPD	7%
Depression	7%
No Depression	6%
Kidney Disease	5%
No Kidney Disease	7%
Severe Vision Impairment	5%
No Severe Vision Impairment	7%
Recent Increased Confusion/Memory Loss	6%
No Recent Increased Confusion/Memory Loss	6%
< 6 Hours of Sleep	12%
6+ Hours of Sleep	6%
Always/Almost Always Wear Seat Belt	5%
Sometimes/Seldom/Never Wear Seat Belt	14%
Drank Alcohol in Past 30 Days	8%
Did Not Drink Alcohol in Past 30 Days	5%
Binge Drinker	13%
Not a Binge Drinker	5%
Heavy Drinker	14%
Not a Heavy Drinker	6%
Good/Very Good/Excellent Health Status	7%
Fair/Poor Health Status	7%
Physical Health Not Good for 0-29 Days of the Past 30	7%
Physical Health Not Good for 30 Days of the Past 30	6%
Mental Health Not Good for 0-19 Days of the Past 30	7%
Mental Health Not Good for 20-30 Days of the Past 30	7%
Usual Activities Unattainable for 0-9 Days of the Past 30	7%
Usual Activities Unattainable for 10-30 Days of the Past 30	8%
Physical, Mental, or Emotional Problems	5%
No Physical, Mental, or Emotional Problems	7%
Disability with Special Equipment Needed	5%
No Disability with Special Equipment Needed	7%
< 2 Hours of TV per Day	6%
2+ Hours of TV per Day	7%
Been Tested for HIV (18-64)	7%
Never Been Tested for HIV (18-64)	6%

Source: The Behavioral Risk Factor Surveillance System, South Dakota Department of Health, 2011-2013

Figure 26, below, shows the percent of respondents whose doctor, nurse, or other health professional advised them to stop using smokeless tobacco. A little under half, 49 percent, of the respondents stated they were advised to quit using smokeless tobacco by a health professional.

**Figure 26**  
**Percent of Respondents Advised to Quit Using Smokeless Tobacco by a Doctor, Nurse, or Other Health Professional, 2011-2013**



Source: The Behavioral Risk Factor Surveillance System, South Dakota Department of Health, 2011-2013

## QUITLINE PROGRAM

**Definition:** Respondents who have heard of the South Dakota Quitline program.

### Prevalence of Respondents Who Have Heard of QuitLine

- South Dakota 80%
- There is no nationwide median for the South Dakota QuitLine program.

### Healthy People 2020 Objective

There was no stated Healthy People 2020 Objective for respondents who have heard of the South Dakota QuitLine program.

<b>Table 26</b>		
<b>Respondents Who Have Heard of the QuitLine Program, 2013</b>		
		<b>2013</b>
<b>Total</b>		<b>80%</b>
<b>Gender</b>	Male	77%
	Female	82%
<b>Age</b>	18-34	85%
	35-44	87%
	45-54	87%
	55-64	76%
	65-74	74%
	75+	51%
<b>Race</b>	White	81%
	American Indian	74%
<b>Ethnicity</b>	Hispanic	*
	Non-Hispanic	80%
<b>Household Income</b>	Less than \$15,000	79%
	\$15,000-\$24,999	76%
	\$25,000-\$34,999	79%
	\$35,000-\$49,999	84%
	\$50,000-\$74,999	82%
	\$75,000+	85%
<b>Education</b>	Less than High School, G.E.D.	71%
	High School, G.E.D.	76%
	Some Post-High School	83%
	College Graduate	84%
<b>Employment Status</b>	Employed for Wages	86%
	Self-employed	74%
	Unemployed	80%
	Homemaker	80%
	Student	83%
	Retired	65%
	Unable to Work	72%
<b>Marital Status</b>	Married/Unmarried Couple	81%
	Divorced/Separated	85%
	Widowed	60%
	Never Married	79%
<b>Home Ownership Status</b>	Own Home	80%
	Rent Home	81%
<b>Children Status</b>	Children in Household (Ages 18-44)	87%
	No Children in Household (Ages 18-44)	83%
<b>Phone Status</b>	Landline	75%
	Cell Phone	85%
<b>Pregnancy Status</b>	Pregnant (Ages 18-44)	*
	Not Pregnant (Ages 18-44)	87%

Note: \*Results based on small sample sizes have been suppressed.

Source: The Behavioral Risk Factor Surveillance System, South Dakota Department of Health, 2013

## Further Analysis

The following data show the percent of those who have heard of the *QuitLine* program for various health behaviors and conditions. For example, 88 percent of respondents who currently smoke have heard of the *QuitLine* program, while 77 percent of respondents who have never smoked have heard of the *QuitLine* program.

<b>Health Behavior or Condition</b>	<b>% Heard of <i>QuitLine</i></b>
Underweight (BMI < 18.5)	67%
Recommended (BMI = 18.5-24.9)	82%
Overweight (BMI = 25.0-29.9)	78%
Obese - Class I (BMI = 30.0-34.9)	78%
Obese - Classes II & III (BMI = 35.0+)	83%
Leisure Time Exercise	82%
No Leisure Time Exercise	71%
Met Physical Activity Recommendations	82%
Did Not Meet Physical Activity Recommendations	77%
2+ Servings of Fruit per Day	80%
< 2 Servings of Fruit per Day	80%
3+ Servings of Vegetables per Day	81%
< 3 Servings of Vegetables per Day	80%
5+ Servings of Fruits and Vegetables per Day	83%
< 5 Servings of Fruits and Vegetables per Day	80%
Current Smoker	88%
Former Smoker	80%
Never Smoked	77%
Smokeless Tobacco	84%
No Smokeless Tobacco	79%
Diabetes	76%
No Diabetes	80%
High Blood Pressure	73%
No High Blood Pressure	83%
High Blood Cholesterol	78%
No High Blood Cholesterol	81%
Health Insurance (18-64)	85%
No Health Insurance (18-64)	80%
Employer Based Coverage (18-64)	88%
Private Health Plan (18-64)	78%
Medicare (18-64)	79%
Medicaid or Medical Assistance (18-64)	84%
The Military, CHAMPUS, TriCare, or the VA (18-64)	86%
The Indian Health Service (18-64)	73%
Flu Shot (65+)	64%
No Flu Shot (65+)	61%
Pneumonia Shot (65+)	63%
No Pneumonia Shot (65+)	64%
Previously Had Heart Attack	74%
Never Had Heart Attack	80%
Have Angina or Coronary Heart Disease	75%
Do Not Have Angina or Coronary Heart Disease	80%
Previously Had a Stroke	62%
Never Had a Stroke	80%

**Table 27 (continued)  
 Heard of the *QuitLine* Program for Selected Health Behaviors and  
 Conditions, 2013**

<b>Health Behavior or Condition</b>	<b>% Heard of <i>QuitLine</i></b>
Cancer	76%
No Cancer	80%
Skin Cancer	72%
No Skin Cancer	80%
Current Asthma	79%
Former Asthma	83%
Never Had Asthma	80%
Arthritis	76%
No Arthritis	81%
COPD	77%
No COPD	80%
Depression	88%
No Depression	79%
Kidney Disease	78%
No Kidney Disease	80%
Severe Vision Impairment	73%
No Severe Vision Impairment	80%
Recent Increased Confusion/Memory Loss	77%
No Recent Increased Confusion/Memory Loss	80%
< 6 Hours of Sleep	82%
6+ Hours of Sleep	80%
Always/Almost Always Wear Seat Belt	80%
Sometimes/Seldom/Never Wear Seat Belt	77%
Drank Alcohol in Past 30 Days	85%
Did Not Drink Alcohol in Past 30 Days	73%
Binge Drinker	88%
Not a Binge Drinker	78%
Heavy Drinker	85%
Not a Heavy Drinker	80%
Good/Very Good/Excellent Health Status	81%
Fair/Poor Health Status	73%
Physical Health Not Good for 0-29 Days of the Past 30	80%
Physical Health Not Good for 30 Days of the Past 30	71%
Mental Health Not Good for 0-19 Days of the Past 30	80%
Mental Health Not Good for 20-30 Days of the Past 30	82%
Usual Activities Unattainable for 0-9 Days of the Past 30	80%
Usual Activities Unattainable for 10-30 Days of the Past 30	76%
Physical, Mental, or Emotional Problems	76%
No Physical, Mental, or Emotional Problems	81%
Disability with Special Equipment Needed	67%
No Disability with Special Equipment Needed	81%
< 2 Hours of TV per Day	84%
2+ Hours of TV per Day	78%
Been Tested for HIV (18-64)	85%
Never Been Tested for HIV (18-64)	78%

Source: The Behavioral Risk Factor Surveillance System, South Dakota Department of Health, 2013