


South Dakota

Demographic Information



South Dakota is located in the north central portion of the United States and averages 10.7 persons per square mile.

2013 Population Information

| Subject | Number | Percent |
|--|---------|---------|
| Total population | 844,877 | 100.0 |
| White | 725,386 | 85.9 |
| Black or African American | 15,735 | 1.9 |
| American Indian & Alaska Native | 75,233 | 8.9 |
| Asian | 9,863 | 1.2 |
| Native Hawaiian & Other Pacific Islander | 622 | 0.1 |
| Two or More Races | 18,038 | 2.1 |
| Under 5 years | 59,957 | 7.1 |
| Under 18 years | 207,959 | 24.6 |
| 65 years and over | 126,163 | 14.9 |

Source: United States Census Bureau, 2013 Population Estimates

Health Status Indicators 2009-2013

Nativity

| | |
|---|------|
| ● Percent of Low Birth Weight Infants | 6.3 |
| * Percent of Mothers Receiving Care in 1st Trimester | 70.0 |
| * Percent of Mothers Who Used Tobacco While Pregnant ¹ | 16.8 |
| Percent of Births Less Than 37 Wks. of Gestation | 8.1 |
| Average Age of Mother | 27.2 |
| Teenage Birth Rate ² | 15.8 |
| Percent White Births | 77.2 |
| Percent American Indian Births | 16.1 |
| Percent Unmarried | 38.3 |
| Percent WIC births | 38.7 |
| Percent Breastfeeding at discharge | 74.6 |
| Percent Payment-Private Insurance | 55.8 |
| Percent Payment-Medicaid | 34.9 |
| Percent C-Section | 25.8 |

●Denotes a health status indicator which is significantly lower than the national average.

○Denotes a health status indicator which is significantly higher than the national average.

*Due to lack of comparable data from all 50 states, significant differences cannot be calculated.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

| | |
|---------------------------------------|-------|
| ● All Causes | 700.1 |
| ● Heart Disease | 155.7 |
| ● Malignant Neoplasms (Cancer) | 162.3 |
| Trachea, Bronchus, & Lung | 43.5 |
| Colon, Rectum, & Anus | 15.9 |
| Pancreas | 10.5 |
| Female Breast | 20.0 |
| ○ Leukemia | 7.8 |
| Prostate | 19.9 |
| ○ Accidents | 44.4 |
| ○ Motor Vehicle Accidents | 16.0 |
| ○ Alzheimer's Disease | 36.5 |
| ○ Cerebrovascular Disease | 39.0 |
| ○ Chronic Lower Respiratory Diseases | 44.3 |
| ○ Diabetes Mellitus | 23.3 |
| Influenza & Pneumonia | 15.5 |
| ○ Intentional Self-Harm (Suicide) | 16.5 |
| ○ Chronic Liver Disease and Cirrhosis | 11.2 |
| ○ Infant Mortality | 7.05 |

Leading Causes of Death

| Leading Causes of Death | Total Deaths |
|---------------------------------------|--------------|
| 1. Heart Disease | 8,273 |
| 2. Malignant Neoplasms (Cancer) | 8,010 |
| 3. Chronic Lower Respiratory Diseases | 2,268 |
| 4. Alzheimer's Disease | 2,108 |
| 5. Cerebrovascular Diseases | 2,094 |
| 3. Accidents | 1,987 |
| 7. Diabetes | 1,166 |
| 8. Influenza and Pneumonia | 853 |
| 9. Suicide | 674 |
| 10. Unspecified Dementia | 508 |
| Percent of Deaths due to tobacco use | 17.4 |
| Median age at death | 80 |

●Denotes a health status indicator which is significantly lower than the national average.

○Denotes a health status indicator which is significantly higher than the national average.


³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Aurora County

Demographic Information



Aurora County is located in the south central portion of the state and averages 3.8 persons per square mile.

2013 Population Information

| Subject | Number | Percent |
|--|--------|---------|
| Total population | 2,720 | 100.0 |
| White | 2,617 | 96.2 |
| Black or African American | 12 | 0.4 |
| American Indian & Alaska Native | 50 | 1.8 |
| Asian | 17 | 0.6 |
| Native Hawaiian & Other Pacific Islander | 0 | 0.0 |
| Two or More Races | 24 | 0.9 |
| Under 5 years | 182 | 6.7 |
| Under 18 years | 720 | 26.5 |
| 65 years and over | 543 | 20.0 |

Source: United States Census Bureau, 2013 Population Estimates

Health Status Indicators 2009-2013

Natality

| | |
|---|------|
| Percent of Low Birth Weight Infants | 4.7 |
| Percent of Mothers Receiving Care in 1st Trimester | 68.3 |
| Percent of Mothers Who Used Tobacco While Pregnant ¹ | 13.4 |
| Percent of Births Less Than 37 Wks. of Gestation | 8.7 |
| ○ Average Age of Mother | 28.6 |
| Teenage Birth Rate ² | LNE |
| Percent White Births | 98.3 |
| Percent American Indian Births | LNE |
| ● Percent Unmarried | 23.8 |
| Percent WIC births | 36.3 |
| Percent Breastfeeding at discharge | 73.3 |
| ○ Percent Payment-Private Insurance | 72.7 |
| ● Percent Payment-Medicaid | 21.5 |
| ○ Percent C-Section | 37.8 |

Mortality³

| | |
|-------------------------------------|-------|
| All Causes | 632.9 |
| Heart Disease | 179.1 |
| Malignant Neoplasms (Cancer) | 157.4 |
| Trachea, Bronchus, & Lung | 55.9 |
| Colon, Rectum, & Anus | 17.9 |
| Pancreas | LNE |
| Female Breast | 51.6 |
| Leukemia | LNE |
| Prostate | 30.4 |
| Accidents | 34.4 |
| Motor Vehicle Accidents | LNE |
| Alzheimer's Disease | 26.8 |
| Cerebrovascular Disease | 49.5 |
| Chronic Lower Respiratory Diseases | 24.0 |
| Diabetes Mellitus | 28.8 |
| Influenza & Pneumonia | 15.8 |
| Intentional Self-Harm (Suicide) | LNE |
| Chronic Liver Disease and Cirrhosis | LNE |
| Infant Mortality | LNE |

Leading Causes of Death

| Leading Causes of Death | Total Deaths |
|--|--------------|
| 1. Heart Disease | 44 |
| 2. Malignant Neoplasms (Cancer) | 35 |
| 3. Cerebrovascular Diseases | 11 |
| 4. Alzheimer's Disease | 8 |
| T5. Chronic Lower Respiratory Diseases | 6 |
| T5. Accidents | 6 |
| T5. Diabetes | 6 |
| 8. Influenza and pneumonia | 5 |
| Percent of Deaths due to tobacco use | 15.0 |
| Median age at death | 83 |

● Denotes a health status indicator which is significantly lower than the state average.

○ Denotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

● Denotes a health status indicator which is significantly lower than the state average.

○ Denotes a health status indicator which is significantly higher than the state average.


³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Beadle County

Demographic Information



Beadle County is located in the center of eastern South Dakota and averages 13.8 persons per square mile.

2013 Population Information

| Subject | Number | Percent |
|--|--------|---------|
| Total population | 18,080 | 100.0 |
| White | 16,158 | 89.4 |
| Black or African American | 308 | 1.7 |
| American Indian & Alaska Native | 227 | 1.3 |
| Asian | 1,055 | 5.8 |
| Native Hawaiian & Other Pacific Islander | 32 | 0.2 |
| Two or More Races | 300 | 1.7 |
| Under 5 years | 1,484 | 8.2 |
| Under 18 years | 4,561 | 25.2 |
| 65 years and over | 3,021 | 16.7 |

Source: United States Census Bureau, 2013 Population Estimates

Health Status Indicators 2009-2013

Natality

| | |
|---|------|
| Percent of Low Birth Weight Infants | 6.9 |
| ● Percent of Mothers Receiving Care in 1st Trimester | 55.2 |
| Percent of Mothers Who Used Tobacco While Pregnant ¹ | 17.0 |
| Percent of Births Less Than 37 Wks. of Gestation | 7.9 |
| ● Average Age of Mother | 26.3 |
| Teenage Birth Rate ² | 21.3 |
| Percent White Births | 83.1 |
| Percent American Indian Births | 0.9 |
| Percent Unmarried | 38.5 |
| ○ Percent WIC births | 52.7 |
| Percent Breastfeeding at discharge | 72.8 |
| ● Percent Payment-Private Insurance | 48.4 |
| ○ Percent Payment-Medicaid | 39.8 |
| ○ Percent C-Section | 33.7 |

●Denotes a health status indicator which is significantly lower than the state average.
○Denotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

| | |
|-------------------------------------|-------|
| All Causes | 726.4 |
| Heart Disease | 164.0 |
| Malignant Neoplasms (Cancer) | 158.3 |
| ● Trachea, Bronchus, & Lung | 30.8 |
| Colon, Rectum, & Anus | 13.8 |
| Pancreas | 10.3 |
| Female Breast | 22.1 |
| Leukemia | 9.3 |
| Prostate | 16.5 |
| Accidents | 45.6 |
| Motor Vehicle Accidents | 15.5 |
| Alzheimer's Disease | 38.5 |
| Cerebrovascular Disease | 43.8 |
| Chronic Lower Respiratory Diseases | 40.6 |
| ○ Diabetes Mellitus | 43.9 |
| Influenza & Pneumonia | 20.7 |
| Intentional Self-Harm (Suicide) | 8.8 |
| Chronic Liver Disease and Cirrhosis | 6.3 |
| Infant Mortality | 13.33 |

Leading Causes of Death

| | Total Deaths |
|---|--------------|
| 1. Heart Disease | 231 |
| 2. Malignant Neoplasms (Cancer) | 197 |
| 3. Alzheimer's Disease | 64 |
| 4. Cerebrovascular Diseases | 62 |
| 5. Diabetes | 56 |
| 6. Chronic Lower Respiratory Diseases | 55 |
| 7. Accidents | 44 |
| 8. Influenza and Pneumonia | 29 |
| 9. Certain Conditions Originating in the Perinatal Period | 13 |
| 10. Essential (Primary) Hypertension and Hypertensive Renal Disease | 12 |

Percent of Deaths due to tobacco use 17.7

Median age at death 81

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Bennett County

Demographic Information



Bennett County is located on the southern border in western South Dakota and averages 2.9 persons per square mile.

2013 Population Information

| Subject | Number | Percent |
|--|--------|---------|
| Total population | 3,452 | 100.0 |
| White | 1,174 | 34.0 |
| Black or African American | 13 | 0.4 |
| American Indian & Alaska Native | 2,060 | 59.7 |
| Asian | 23 | 0.7 |
| Native Hawaiian & Other Pacific Islander | 1 | 0.0 |
| Two or More Races | 181 | 5.2 |
| Under 5 years | 351 | 10.2 |
| Under 18 years | 1,161 | 33.6 |
| 65 years and over | 404 | 11.7 |

Source: United States Census Bureau, 2013 Population Estimates

Health Status Indicators 2009-2013

Nativity

| | |
|---|------|
| Percent of Low Birth Weight Infants | 5.5 |
| ● Percent of Mothers Receiving Care in 1st Trimester | 41.6 |
| ○ Percent of Mothers Who Used Tobacco While Pregnant ¹ | 22.3 |
| Percent of Births Less Than 37 Wks. of Gestation | 8.2 |
| ● Average Age of Mother | 24.6 |
| ○ Teenage Birth Rate ² | 44.6 |
| Percent White Births | 16.7 |
| Percent American Indian Births | 77.6 |
| ○ Percent Unmarried | 74.8 |
| ○ Percent WIC births | 76.1 |
| ● Percent Breastfeeding at discharge | 49.7 |
| ● Percent Payment-Private Insurance | 12.5 |
| ○ Percent Payment-Medicaid | 62.8 |
| Percent C-Section | 21.5 |

●Denotes a health status indicator which is significantly lower than the state average.
○Denotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

| | |
|-------------------------------------|-------|
| ○ All Causes | 931.5 |
| Heart Disease | 187.2 |
| Malignant Neoplasms (Cancer) | 162.1 |
| Trachea, Bronchus, & Lung | 30.5 |
| Colon, Rectum, & Anus | 33.9 |
| Pancreas | 18.7 |
| Female Breast | LNE |
| Leukemia | LNE |
| Prostate | LNE |
| Accidents | 79.2 |
| ○ Motor Vehicle Accidents | 52.5 |
| Alzheimer's Disease | LNE |
| Cerebrovascular Disease | 46.5 |
| Chronic Lower Respiratory Diseases | 51.7 |
| Diabetes Mellitus | 36.6 |
| ○ Influenza & Pneumonia | 50.5 |
| Intentional Self-Harm (Suicide) | 21.9 |
| Chronic Liver Disease and Cirrhosis | 35.3 |
| Infant Mortality | 22.92 |

Leading Causes of Death

| Leading Causes of Death | Total Deaths |
|--|--------------|
| 1. Heart Disease | 31 |
| 2. Malignant Neoplasms (Cancer) | 27 |
| 3. Accidents | 14 |
| T4. Chronic Lower Respiratory Diseases | 9 |
| T4. Influenza and Pneumonia | 9 |
| 6. Cerebrovascular Diseases | 8 |
| 7. Diabetes | 6 |
| 8. Chronic Liver Disease and Cirrhosis | 5 |
| Percent of Deaths due to tobacco use | 21.0 |
| Median age at death | 74 |

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Bon Homme County

Demographic Information



Bon Homme County is located on the Nebraska border in eastern South Dakota and averages 12.5 persons per square mile.

2013 Population Information

| Subject | Number | Percent |
|--|--------|---------|
| Total population | 7,046 | 100.0 |
| White | 6,319 | 89.7 |
| Black or African American | 81 | 1.1 |
| American Indian & Alaska Native | 524 | 7.4 |
| Asian | 19 | 0.3 |
| Native Hawaiian & Other Pacific Islander | 0 | 0.0 |
| Two or More Races | 103 | 1.5 |
| Under 5 years | 324 | 4.6 |
| Under 18 years | 1,368 | 19.4 |
| 65 years and over | 1,374 | 19.5 |

Source: United States Census Bureau, 2013 Population Estimates

Health Status Indicators 2009-2013

Nativity

| | |
|---|------|
| Percent of Low Birth Weight Infants | 6.0 |
| Percent of Mothers Receiving Care in 1st Trimester | 79.0 |
| ● Percent of Mothers Who Used Tobacco While Pregnant ¹ | 11.4 |
| Percent of Births Less Than 37 Wks. of Gestation | 7.9 |
| ○ Average Age of Mother | 28.2 |
| Teenage Birth Rate ² | LNE |
| Percent White Births | 94.9 |
| Percent American Indian Births | 3.5 |
| ● Percent Unmarried | 19.6 |
| Percent WIC births | 40.1 |
| Percent Breastfeeding at discharge | 76.1 |
| ○ Percent Payment-Private Insurance | 74.9 |
| ● Percent Payment-Medicaid | 20.6 |
| Percent C-Section | 27.2 |

●Denotes a health status indicator which is significantly lower than the state average.
○Denotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

| | |
|-------------------------------------|-------|
| ● All Causes | 595.2 |
| Heart Disease | 130.6 |
| Malignant Neoplasms (Cancer) | 141.5 |
| Trachea, Bronchus, & Lung | 48.5 |
| Colon, Rectum, & Anus | 14.8 |
| Pancreas | LNE |
| Female Breast | 11.1 |
| Leukemia | 11.4 |
| Prostate | 12.3 |
| Accidents | 34.3 |
| Motor Vehicle Accidents | 23.6 |
| ● Alzheimer's Disease | 24.4 |
| Cerebrovascular Disease | 41.4 |
| Chronic Lower Respiratory Diseases | 44.4 |
| Diabetes Mellitus | 19.4 |
| Influenza & Pneumonia | 9.0 |
| Intentional Self-Harm (Suicide) | 14.7 |
| Chronic Liver Disease and Cirrhosis | LNE |
| Infant Mortality | LNE |

Leading Causes of Death

| Leading Causes of Death | Total Deaths |
|---|--------------|
| 1. Heart Disease | 88 |
| 2. Malignant Neoplasms (Cancer) | 80 |
| 3. Cerebrovascular Diseases | 29 |
| 4. Chronic Lower Respiratory Diseases | 28 |
| 5. Alzheimer's Disease | 21 |
| 6. Accidents | 15 |
| 7. Diabetes | 12 |
| 8. Neoplasms of Uncertain or Unknown Behavior | 10 |
| T9. Influenza and Pneumonia | 7 |
| T9. Unspecified Dementia | 7 |
| T9. Vascular Dementia | 7 |

| | |
|--------------------------------------|------|
| Percent of Deaths due to tobacco use | 14.1 |
| Median age at death | 84 |

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Brookings County

Demographic Information



Brookings County is located on the eastern border of the state and averages 40.3 persons per square mile.

2013 Population Information

| Subject | Number | Percent |
|--|--------|---------|
| Total population | 32,968 | 100.0 |
| White | 30,683 | 93.1 |
| Black or African American | 485 | 1.5 |
| American Indian & Alaska Native | 429 | 1.3 |
| Asian | 925 | 2.8 |
| Native Hawaiian & Other Pacific Islander | 22 | 0.1 |
| Two or More Races | 424 | 1.3 |
| Under 5 years | 1,972 | 6.0 |
| Under 18 years | 6,422 | 19.5 |
| 65 years and over | 3,397 | 10.3 |

Source: United States Census Bureau, 2013 Population Estimates

Health Status Indicators 2009-2013

Nativity

| | |
|---|------|
| ● Percent of Low Birth Weight Infants | 4.9 |
| ○ Percent of Mothers Receiving Care in 1st Trimester | 76.5 |
| ● Percent of Mothers Who Used Tobacco While Pregnant ¹ | 11.9 |
| ● Percent of Births Less Than 37 Wks. of Gestation | 6.6 |
| ○ Average Age of Mother | 28.1 |
| ● Teenage Birth Rate ² | 4.7 |
| Percent White Births | 88.9 |
| Percent American Indian Births | 1.9 |
| ● Percent Unmarried | 24.0 |
| ● Percent WIC births | 27.8 |
| ○ Percent Breastfeeding at discharge | 81.9 |
| ○ Percent Payment-Private Insurance | 75.1 |
| ● Percent Payment-Medicaid | 19.9 |
| ● Percent C-Section | 21.3 |

●Denotes a health status indicator which is significantly lower than the state average.
○Denotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

| | |
|---------------------------------------|-------|
| All Causes | 654.1 |
| Heart Disease | 144.3 |
| Malignant Neoplasms (Cancer) | 168.1 |
| Trachea, Bronchus, & Lung | 40.0 |
| Colon, Rectum, & Anus | 15.5 |
| Pancreas | 11.3 |
| Female Breast | 11.6 |
| Leukemia | 12.3 |
| Prostate | 22.4 |
| ● Accidents | 29.2 |
| ● Motor Vehicle Accidents | 9.4 |
| Alzheimer's Disease | 35.1 |
| ○ Cerebrovascular Disease | 52.7 |
| Chronic Lower Respiratory Diseases | 28.6 |
| ● Diabetes Mellitus | 14.1 |
| Influenza & Pneumonia | 13.3 |
| ● Intentional Self-Harm (Suicide) | 8.2 |
| ● Chronic Liver Disease and Cirrhosis | 4.2 |
| Infant Mortality | 6.56 |

Leading Causes of Death

| | Total Deaths |
|---------------------------------------|--------------|
| 1. Malignant Neoplasms (Cancer) | 231 |
| 2. Heart Disease | 217 |
| 3. Cerebrovascular Diseases | 82 |
| 4. Alzheimer's Disease | 57 |
| 5. Accidents | 44 |
| 6. Chronic Lower Respiratory Diseases | 42 |
| 7. Diabetes | 21 |
| 8. Influenza and Pneumonia | 20 |
| 9. Unspecified Dementia | 17 |
| 10. Intentional Self-Harm (Suicide) | 12 |

| | |
|--------------------------------------|------|
| Percent of Deaths due to tobacco use | 16.7 |
| Median age at death | 83 |

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Brown County

Demographic Information



Brown County is located on the North Dakota border in the eastern part of the state and averages 21.3 persons per square mile.

2013 Population Information

| Subject | Number | Percent |
|--|--------|---------|
| Total population | 37,907 | 100.0 |
| White | 34,892 | 92.0 |
| Black or African American | 525 | 1.4 |
| American Indian & Alaska Native | 1,204 | 3.2 |
| Asian | 563 | 1.5 |
| Native Hawaiian & Other Pacific Islander | 69 | 0.2 |
| Two or More Races | 654 | 1.7 |
| Under 5 years | 2,537 | 6.7 |
| Under 18 years | 8,873 | 23.4 |
| 65 years and over | 6,019 | 15.9 |

Source: United States Census Bureau, 2013 Population Estimates

Health Status Indicators 2009-2013

Nativity

| | |
|---|------|
| Percent of Low Birth Weight Infants | 5.7 |
| ○ Percent of Mothers Receiving Care in 1st Trimester | 76.3 |
| Percent of Mothers Who Used Tobacco While Pregnant ¹ | 17.7 |
| Percent of Births Less Than 37 Wks. of Gestation | 7.6 |
| Average Age of Mother | 27.3 |
| ● Teenage Birth Rate ² | 9.3 |
| Percent White Births | 90.2 |
| Percent American Indian Births | 5.4 |
| ● Percent Unmarried | 34.1 |
| ● Percent WIC births | 34.3 |
| ○ Percent Breastfeeding at discharge | 79.2 |
| ○ Percent Payment-Private Insurance | 65.1 |
| ● Percent Payment-Medicaid | 29.5 |
| ○ Percent C-Section | 28.4 |

●Denotes a health status indicator which is significantly lower than the state average.
○Denotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

| | |
|-------------------------------------|-------|
| All Causes | 706.2 |
| Heart Disease | 166.8 |
| Malignant Neoplasms (Cancer) | 155.6 |
| Trachea, Bronchus, & Lung | 45.1 |
| Colon, Rectum, & Anus | 16.2 |
| Pancreas | 12.7 |
| Female Breast | 22.0 |
| Leukemia | 7.2 |
| Prostate | 15.3 |
| ● Accidents | 33.1 |
| ● Motor Vehicle Accidents | 10.2 |
| ○ Alzheimer's Disease | 46.8 |
| Cerebrovascular Disease | 37.8 |
| Chronic Lower Respiratory Diseases | 41.8 |
| Diabetes Mellitus | 27.4 |
| Influenza & Pneumonia | 17.8 |
| Intentional Self-Harm (Suicide) | 15.2 |
| Chronic Liver Disease and Cirrhosis | 7.6 |
| Infant Mortality | 5.80 |

Leading Causes of Death

| | Total Deaths |
|---------------------------------------|--------------|
| 1. Heart Disease | 477 |
| 2. Malignant Neoplasms (Cancer) | 367 |
| 3. Alzheimer's Disease | 143 |
| 4. Chronic Lower Respiratory Diseases | 110 |
| 5. Cerebrovascular Diseases | 108 |
| 6. Accidents | 73 |
| 7. Diabetes | 70 |
| 8. Influenza and Pneumonia | 53 |
| 9. Unspecified Dementia | 32 |
| 10. Intentional Self-Harm (Suicide) | 28 |

| | |
|--------------------------------------|------|
| Percent of Deaths due to tobacco use | 16.1 |
| Median age at death | 82 |

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Brule County

Demographic Information



Brule County is located in the south central part of the state and averages 6.4 persons per square mile.

2013 Population Information

| Subject | Number | Percent |
|--|--------|---------|
| Total population | 5,366 | 100.0 |
| White | 4,659 | 86.8 |
| Black or African American | 25 | 0.5 |
| American Indian & Alaska Native | 500 | 9.3 |
| Asian | 14 | 0.3 |
| Native Hawaiian & Other Pacific Islander | 1 | 0.0 |
| Two or More Races | 167 | 3.1 |
| Under 5 years | 379 | 7.1 |
| Under 18 years | 1,385 | 25.8 |
| 65 years and over | 903 | 16.8 |

Source: United States Census Bureau, 2013 Population Estimates

Health Status Indicators 2009-2013

Nativity

| | |
|---|------|
| Percent of Low Birth Weight Infants | 6.3 |
| Percent of Mothers Receiving Care in 1st Trimester | 65.5 |
| Percent of Mothers Who Used Tobacco While Pregnant ¹ | 16.9 |
| Percent of Births Less Than 37 Wks. of Gestation | 10.4 |
| Average Age of Mother | 27.7 |
| ● Teenage Birth Rate ² | 7.7 |
| Percent White Births | 78.1 |
| Percent American Indian Births | 18.8 |
| Percent Unmarried | 34.0 |
| Percent WIC births | 38.3 |
| Percent Breastfeeding at discharge | 71.4 |
| ○ Percent Payment-Private Insurance | 66.2 |
| Percent Payment-Medicaid | 30.8 |
| Percent C-Section | 24.9 |

●Denotes a health status indicator which is significantly lower than the state average.
○Denotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

| | |
|-------------------------------------|-------|
| All Causes | 698.2 |
| Heart Disease | 159.4 |
| Malignant Neoplasms (Cancer) | 173.2 |
| Trachea, Bronchus, & Lung | 53.5 |
| Colon, Rectum, & Anus | 22.6 |
| Pancreas | 7.3 |
| Female Breast | 17.0 |
| Leukemia | LNE |
| Prostate | 32.8 |
| Accidents | 50.8 |
| Motor Vehicle Accidents | 11.0 |
| Alzheimer's Disease | 29.6 |
| Cerebrovascular Disease | 32.5 |
| Chronic Lower Respiratory Diseases | 24.5 |
| Diabetes Mellitus | 39.9 |
| Influenza & Pneumonia | 15.6 |
| Intentional Self-Harm (Suicide) | 14.1 |
| Chronic Liver Disease and Cirrhosis | 10.1 |
| Infant Mortality | 13.70 |

Leading Causes of Death

| Leading Causes of Death | Total Deaths |
|--|--------------|
| 1. Heart Disease | 66 |
| 2. Malignant Neoplasms (Cancer) | 64 |
| 3. Accidents | 18 |
| 4. Alzheimer's Disease | 17 |
| T5. Cerebrovascular Diseases | 14 |
| T5. Diabetes | 14 |
| 7. Chronic Lower Respiratory Diseases | 10 |
| 8. Pneumonitis due to solids and liquids | 8 |
| 9. Influenza and Pneumonia | 7 |
| 10. Parkinson's Disease | 5 |

| | |
|--------------------------------------|------|
| Percent of Deaths due to tobacco use | 20.7 |
| Median age at death | 81 |

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Buffalo County

Demographic Information



Buffalo County is located in the central portion of the state and averages 4.1 persons per square mile.

2013 Population Information

| Subject | Number | Percent |
|--|--------|---------|
| Total population | 2,024 | 100.0 |
| White | 363 | 17.9 |
| Black or African American | 11 | 0.5 |
| American Indian & Alaska Native | 1,608 | 79.4 |
| Asian | 1 | 0.0 |
| Native Hawaiian & Other Pacific Islander | 0 | 0.0 |
| Two or More Races | 41 | 2.0 |
| Under 5 years | 264 | 13.0 |
| Under 18 years | 805 | 39.8 |
| 65 years and over | 149 | 7.4 |

Source: United States Census Bureau, 2013 Population Estimates

Health Status Indicators 2009-2013

Nativity

| | |
|---|------|
| Percent of Low Birth Weight Infants | 7.4 |
| ● Percent of Mothers Receiving Care in 1st Trimester | 40.0 |
| ○ Percent of Mothers Who Used Tobacco While Pregnant ¹ | 34.7 |
| Percent of Births Less Than 37 Wks. of Gestation | 10.5 |
| ● Average Age of Mother | 25.8 |
| ○ Teenage Birth Rate ² | 67.7 |
| Percent White Births | 15.6 |
| Percent American Indian Births | 79.0 |
| ○ Percent Unmarried | 75.5 |
| ○ Percent WIC births | 74.5 |
| ● Percent Breastfeeding at discharge | 46.3 |
| ● Percent Payment-Private Insurance | 18.3 |
| ○ Percent Payment-Medicaid | 73.9 |
| Percent C-Section | 24.9 |

● Denotes a health status indicator which is significantly lower than the state average.
○ Denotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

| | |
|---------------------------------------|---------|
| ○ All Causes | 1,304.8 |
| Heart Disease | 231.9 |
| Malignant Neoplasms (Cancer) | 242.6 |
| Trachea, Bronchus, & Lung | LNE |
| Colon, Rectum, & Anus | 47.4 |
| Female Breast | LNE |
| Pancreas | LNE |
| Leukemia | LNE |
| Prostate | LNE |
| Accidents | 88.0 |
| Motor Vehicle Accidents | LNE |
| Alzheimer's Disease | 60.2 |
| Cerebrovascular Disease | 51.9 |
| Chronic Lower Respiratory Diseases | LNE |
| Diabetes Mellitus | 65.9 |
| Influenza & Pneumonia | LNE |
| Intentional Self-Harm (Suicide) | 36.2 |
| ○ Chronic Liver Disease and Cirrhosis | 121.1 |
| Infant Mortality | 23.35 |

Leading Causes of Death

| Leading Causes of Death | Total Deaths |
|--|--------------|
| T1. Heart Disease | 15 |
| T1. Malignant Neoplasms (Cancer) | 15 |
| 3. Chronic Liver Disease and Cirrhosis | 9 |
| 4. Accidents | 8 |
| 5. Diabetes | 5 |

| | |
|--------------------------------------|------|
| Percent of Deaths due to tobacco use | 22.8 |
| Median age at death | 57 |

● Denotes a health status indicator which is significantly lower than the state average.

○ Denotes a health status indicator which is significantly higher than the state average.


³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Butte County

Demographic Information



Butte County is located on the western border of the state and averages 4.5 people per square mile.

2013 Population Information

| Subject | Number | Percent |
|--|--------|---------|
| Total population | 10,330 | 100.0 |
| White | 9,749 | 94.4 |
| Black or African American | 80 | 0.8 |
| American Indian & Alaska Native | 202 | 2.0 |
| Asian | 35 | 0.3 |
| Native Hawaiian & Other Pacific Islander | 9 | 0.1 |
| Two or More Races | 255 | 2.5 |
| Under 5 years | 663 | 6.4 |
| Under 18 years | 2,513 | 24.3 |
| 65 years and over | 1,820 | 17.6 |

Source: United States Census Bureau, 2013 Population Estimates

Health Status Indicators 2009-2013

Natality

| | |
|---|------|
| Percent of Low Birth Weight Infants | 5.6 |
| Percent of Mothers Receiving Care in 1st Trimester | 77.5 |
| ○ Percent of Mothers Who Used Tobacco While Pregnant ¹ | 25.1 |
| Percent of Births Less Than 37 Wks. of Gestation | 8.0 |
| ● Average Age of Mother | 26.6 |
| Teenage Birth Rate ² | 19.5 |
| Percent White Births | 93.6 |
| Percent American Indian Births | 2.0 |
| Percent Unmarried | 37.9 |
| ○ Percent WIC births | 49.1 |
| Percent Breastfeeding at discharge | 79.3 |
| ● Percent Payment-Private Insurance | 48.4 |
| ○ Percent Payment-Medicaid | 43.8 |
| Percent C-Section | 22.7 |

●Denotes a health status indicator which is significantly lower than the state average.
○Denotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

| | |
|-------------------------------------|-------|
| ○ All Causes | 775.4 |
| Heart Disease | 187.4 |
| Malignant Neoplasms (Cancer) | 191.2 |
| Trachea, Bronchus, & Lung | 59.6 |
| Colon, Rectum, & Anus | 16.7 |
| Pancreas | 10.8 |
| Female Breast | 22.7 |
| Leukemia | 6.0 |
| Prostate | 35.4 |
| Accidents | 43.4 |
| Motor Vehicle Accidents | 22.7 |
| Alzheimer's Disease | 36.0 |
| Cerebrovascular Disease | 34.1 |
| Chronic Lower Respiratory Diseases | 45.2 |
| Diabetes Mellitus | 26.5 |
| Influenza & Pneumonia | 24.1 |
| Intentional Self-Harm (Suicide) | 24.8 |
| Chronic Liver Disease and Cirrhosis | 7.3 |
| Infant Mortality | 9.32 |

Leading Causes of Death

| Leading Causes of Death | Total Deaths |
|---------------------------------------|--------------|
| 1. Heart Disease | 131 |
| 2. Malignant Neoplasms (Cancer) | 129 |
| 3. Chronic Lower Respiratory Diseases | 31 |
| 4. Alzheimer's Disease | 25 |
| 5. Cerebrovascular Diseases | 23 |
| 6. Accidents | 22 |
| 7. Diabetes | 19 |
| 8. Influenza and Pneumonia | 17 |
| 9. Intentional Self-Harm (Suicide) | 13 |
| T10. Unspecified Dementia | 6 |
| T10. Septicemia | 6 |

| | |
|--------------------------------------|------|
| Percent of Deaths due to tobacco use | 19.1 |
| Median age at death | 79 |

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Campbell County

Demographic Information



Campbell County is located in the north central portion of the state and averages 2.0 persons per square mile.

2013 Population Information

| Subject | Number | Percent |
|--|--------|---------|
| Total population | 1,334 | 100.0 |
| White | 1,307 | 98.0 |
| Black or African American | 2 | 0.1 |
| American Indian & Alaska Native | 7 | 0.5 |
| Asian | 7 | 0.5 |
| Native Hawaiian & Other Pacific Islander | 0 | 0.0 |
| Two or More Races | 11 | 0.8 |
| Under 5 years | 49 | 3.7 |
| Under 18 years | 214 | 16.0 |
| 65 years and over | 338 | 25.3 |

Source: United States Census Bureau, 2013 Population Estimates

Health Status Indicators 2009-2013

Natality

| | |
|---|------|
| Percent of Low Birth Weight Infants | LNE |
| Percent of Mothers Receiving Care in 1st Trimester | 72.6 |
| ● Percent of Mothers Who Used Tobacco While Pregnant ¹ | 4.8 |
| Percent of Births Less Than 37 Wks. of Gestation | 6.6 |
| ○ Average Age of Mother | 28.7 |
| Teenage Birth Rate ² | LNE |
| Percent White Births | 95.0 |
| Percent American Indian Births | LNE |
| ● Percent Unmarried | 20.6 |
| Percent WIC births | 36.7 |
| Percent Breastfeeding at discharge | 88.5 |
| ○ Percent Payment-Private Insurance | 81.4 |
| ● Percent Payment-Medicaid | 8.5 |
| Percent C-Section | 41.9 |

●Denotes a health status indicator which is significantly lower than the state average.
○Denotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

| | |
|-------------------------------------|-------|
| All Causes | 581.6 |
| Heart Disease | 190.6 |
| Malignant Neoplasms (Cancer) | 106.6 |
| Trachea, Bronchus, & Lung | LNE |
| Colon, Rectum, & Anus | 36.2 |
| Pancreas | LNE |
| Female Breast | LNE |
| Leukemia | LNE |
| Prostate | LNE |
| Accidents | 84.8 |
| Motor Vehicle Accidents | LNE |
| Alzheimer's Disease | 23.8 |
| Cerebrovascular Disease | 52.8 |
| Chronic Lower Respiratory Diseases | LNE |
| Diabetes Mellitus | 20.5 |
| Influenza & Pneumonia | LNE |
| Intentional Self-Harm (Suicide) | LNE |
| Chronic Liver Disease and Cirrhosis | LNE |
| Infant Mortality | LNE |

Leading Causes of Death

| Leading Causes of Death | Total Deaths |
|--------------------------------------|--------------|
| 1. Heart Disease | 23 |
| 2. Malignant Neoplasms (Cancer) | 15 |
| 3. Cerebrovascular Diseases | 8 |
| 4. Accidents | 5 |
| Percent of Deaths due to tobacco use | 20.5 |
| Median age at death | 80 |

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Charles Mix County

Demographic Information



Charles Mix County is located in the south central area of the state and averages 8.3 persons per square mile.

2013 Population Information

| Subject | Number | Percent |
|--|--------|---------|
| Total population | 9,241 | 100.0 |
| White | 5,950 | 64.4 |
| Black or African American | 35 | 0.4 |
| American Indian & Alaska Native | 2,968 | 32.1 |
| Asian | 24 | 0.3 |
| Native Hawaiian & Other Pacific Islander | 1 | 0.0 |
| Two or More Races | 263 | 2.8 |
| Under 5 years | 801 | 8.7 |
| Under 18 years | 2,716 | 29.4 |
| 65 years and over | 1,668 | 18.0 |

Source: United States Census Bureau, 2013 Population Estimates

Health Status Indicators 2009-2013

Nativity

| | |
|---|------|
| Percent of Low Birth Weight Infants | 5.2 |
| ● Percent of Mothers Receiving Care in 1st Trimester | 63.3 |
| ○ Percent of Mothers Who Used Tobacco While Pregnant ¹ | 23.1 |
| Percent of Births Less Than 37 Wks. of Gestation | 8.7 |
| ● Average Age of Mother | 25.7 |
| ○ Teenage Birth Rate ² | 26.4 |
| Percent White Births | 45.0 |
| Percent American Indian Births | 48.8 |
| ○ Percent Unmarried | 56.8 |
| ○ Percent WIC births | 61.8 |
| ● Percent Breastfeeding at discharge | 67.5 |
| ● Percent Payment-Private Insurance | 37.1 |
| ○ Percent Payment-Medicaid | 57.2 |
| Percent C-Section | 26.8 |

● Denotes a health status indicator which is significantly lower than the state average.
○ Denotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

| | |
|-------------------------------------|-------|
| All Causes | 747.8 |
| Heart Disease | 188.6 |
| Malignant Neoplasms (Cancer) | 180.6 |
| Trachea, Bronchus, & Lung | 35.1 |
| Colon, Rectum, & Anus | 23.6 |
| Pancreas | 17.2 |
| Female Breast | 18.0 |
| Leukemia | LNE |
| Prostate | 23.7 |
| Accidents | 56.6 |
| Motor Vehicle Accidents | 20.9 |
| Alzheimer's Disease | 49.7 |
| Cerebrovascular Disease | 26.5 |
| Chronic Lower Respiratory Diseases | 34.9 |
| Diabetes Mellitus | 40.1 |
| Influenza & Pneumonia | 18.8 |
| Intentional Self-Harm (Suicide) | 15.9 |
| Chronic Liver Disease and Cirrhosis | 18.5 |
| Infant Mortality | 5.08 |

Leading Causes of Death

| Leading Causes of Death | Total Deaths |
|---|--------------|
| 1. Heart Disease | 129 |
| 2. Malignant Neoplasms (Cancer) | 117 |
| 3. Alzheimer's Disease | 40 |
| 4. Accidents | 29 |
| T5. Chronic Lower Respiratory Diseases | 24 |
| T5. Diabetes | 24 |
| 7. Cerebrovascular Diseases | 20 |
| 8. Influenza and Pneumonia | 14 |
| T9. Chronic Liver Disease and Cirrhosis | 9 |
| T9. Essential (Primary) Hypertension and Hypertensive Renal Disease | 9 |

| | |
|--------------------------------------|------|
| Percent of Deaths due to tobacco use | 13.8 |
| Median age at death | 80 |

● Denotes a health status indicator which is significantly lower than the state average.

○ Denotes a health status indicator which is significantly higher than the state average.


³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Clark County

Demographic Information



Clark County is located in east central South Dakota and averages 3.9 persons per square mile.

2013 Population Information

| Subject | Number | Percent |
|--|--------|---------|
| Total population | 3,610 | 100.0 |
| White | 3,540 | 98.1 |
| Black or African American | 34 | 0.9 |
| American Indian & Alaska Native | 4 | 0.1 |
| Asian | 3 | 0.1 |
| Native Hawaiian & Other Pacific Islander | 0 | 0.0 |
| Two or More Races | 29 | 0.8 |
| Under 5 years | 286 | 7.9 |
| Under 18 years | 835 | 23.1 |
| 65 years and over | 778 | 21.6 |

Source: United States Census Bureau, 2013 Population Estimates

Health Status Indicators 2009-2013

Natality

| | |
|---|------|
| ● Percent of Low Birth Weight Infants | 3.6 |
| ● Percent of Mothers Receiving Care in 1st Trimester | 57.9 |
| Percent of Mothers Who Used Tobacco While Pregnant ¹ | 13.7 |
| Percent of Births Less Than 37 Wks. of Gestation | 5.2 |
| ○ Average Age of Mother | 28.6 |
| Teenage Birth Rate ² | LNE |
| Percent White Births | 98.0 |
| Percent American Indian Births | 1.6 |
| ● Percent Unmarried | 19.1 |
| ● Percent WIC births | 24.5 |
| Percent Breastfeeding at discharge | 86.2 |
| ○ Percent Payment-Private Insurance | 77.3 |
| ● Percent Payment-Medicaid | 18.2 |
| Percent C-Section | 21.5 |

●Denotes a health status indicator which is significantly lower than the state average.
○Denotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

| | |
|-------------------------------------|-------|
| All Causes | 740.0 |
| Heart Disease | 185.1 |
| Malignant Neoplasms (Cancer) | 138.0 |
| Trachea, Bronchus, & Lung | 40.4 |
| Colon, Rectum, & Anus | 12.0 |
| Pancreas | 7.2 |
| Female Breast | 17.0 |
| Leukemia | LNE |
| Prostate | LNE |
| Accidents | 68.7 |
| Motor Vehicle Accidents | 18.1 |
| Alzheimer's Disease | 41.1 |
| Cerebrovascular Disease | 46.8 |
| Chronic Lower Respiratory Diseases | 51.0 |
| Diabetes Mellitus | 24.2 |
| Influenza & Pneumonia | 27.5 |
| Intentional Self-Harm (Suicide) | LNE |
| Chronic Liver Disease and Cirrhosis | LNE |
| Infant Mortality | 11.95 |

Leading Causes of Death

| Leading Causes of Death | Total Deaths |
|---|--------------|
| 1. Heart Disease | 70 |
| 2. Malignant Neoplasms (Cancer) | 48 |
| T3. Chronic Lower Respiratory Diseases | 18 |
| T3. Alzheimer's Disease | 18 |
| T3. Accidents | 18 |
| 6. Cerebrovascular Diseases | 17 |
| 7. Influenza and Pneumonia | 10 |
| 8. Diabetes | 8 |
| 9. Gastrointestinal Hemorrhage, Unspecified | 5 |
| Percent of Deaths due to tobacco use | 15.8 |
| Median age at death | 84 |

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Clay County

Demographic Information



Clay County is located in the southeastern part of the state and averages 33.6 persons per square mile.

2013 Population Information

| Subject | Number | Percent |
|--|--------|---------|
| Total population | 13,935 | 100.0 |
| White | 12,653 | 90.8 |
| Black or African American | 226 | 1.6 |
| American Indian & Alaska Native | 449 | 3.2 |
| Asian | 293 | 2.1 |
| Native Hawaiian & Other Pacific Islander | 9 | 0.1 |
| Two or More Races | 305 | 2.2 |
| Under 5 years | 784 | 5.6 |
| Under 18 years | 2,477 | 17.8 |
| 65 years and over | 1,484 | 10.6 |

Source: United States Census Bureau, 2013 Population Estimates

Health Status Indicators 2009-2013

Natality

| | |
|---|------|
| Percent of Low Birth Weight Infants | 6.3 |
| ○ Percent of Mothers Receiving Care in 1st Trimester | 78.3 |
| ● Percent of Mothers Who Used Tobacco While Pregnant ¹ | 10.6 |
| Percent of Births Less Than 37 Wks. of Gestation | 7.7 |
| ○ Average Age of Mother | 27.8 |
| ● Teenage Birth Rate ² | 4.1 |
| Percent White Births | 90.4 |
| Percent American Indian Births | 4.5 |
| ● Percent Unmarried | 28.3 |
| Percent WIC births | 36.2 |
| Percent Breastfeeding at discharge | 77.3 |
| Percent Payment-Private Insurance | 60.7 |
| Percent Payment-Medicaid | 36.3 |
| ○ Percent C-Section | 33.5 |

●Denotes a health status indicator which is significantly lower than the state average.
○Denotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

| | |
|-------------------------------------|-------|
| All Causes | 763.6 |
| ○ Heart Disease | 217.5 |
| Malignant Neoplasms (Cancer) | 173.7 |
| Trachea, Bronchus, & Lung | 49.5 |
| Colon, Rectum, & Anus | 19.7 |
| Pancreas | 8.5 |
| Female Breast | 25.0 |
| Leukemia | 7.8 |
| Prostate | 27.1 |
| Accidents | 41.6 |
| Motor Vehicle Accidents | 12.3 |
| ● Alzheimer's Disease | 19.2 |
| Cerebrovascular Disease | 51.6 |
| Chronic Lower Respiratory Diseases | 45.8 |
| Diabetes Mellitus | 32.0 |
| Influenza & Pneumonia | 14.8 |
| Intentional Self-Harm (Suicide) | 16.8 |
| Chronic Liver Disease and Cirrhosis | 5.5 |
| Infant Mortality | 7.67 |

Leading Causes of Death

| | Total Deaths |
|--|--------------|
| 1. Heart Disease | 147 |
| 2. Malignant Neoplasms (Cancer) | 106 |
| 3. Cerebrovascular Diseases | 34 |
| 4. Chronic Lower Respiratory Diseases | 30 |
| 5. Accidents | 24 |
| 6. Diabetes | 19 |
| 7. Alzheimer's Disease | 15 |
| 8. Influenza and Pneumonia | 10 |
| 9. Intentional Self-Harm (Suicide) | 9 |
| T10. Parkinson's Disease | 7 |
| T10. Congenital Malformations, Deformations, and Chromosomal Abnormalities | 7 |

| | |
|--------------------------------------|------|
| Percent of Deaths due to tobacco use | 15.8 |
| Median age at death | 81 |

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Codington County

Demographic Information



Codington County is located in eastern South Dakota and averages 39.5 persons per square mile.

2013 Population Information

| Subject | Number | Percent |
|--|--------|---------|
| Total population | 27,853 | 100.0 |
| White | 26,434 | 94.9 |
| Black or African American | 222 | 0.8 |
| American Indian & Alaska Native | 638 | 2.3 |
| Asian | 208 | 0.7 |
| Native Hawaiian & Other Pacific Islander | 1 | 0.0 |
| Two or More Races | 350 | 1.3 |
| Under 5 years | 1,974 | 7.1 |
| Under 18 years | 6,785 | 24.4 |
| 65 years and over | 4,268 | 15.3 |

Source: United States Census Bureau, 2013 Population Estimates

Health Status Indicators 2009-2013

Nativity

| | |
|---|------|
| Percent of Low Birth Weight Infants | 5.5 |
| ○ Percent of Mothers Receiving Care in 1st Trimester | 79.2 |
| ○ Percent of Mothers Who Used Tobacco While Pregnant ¹ | 24.8 |
| ● Percent of Births Less Than 37 Wks. of Gestation | 6.6 |
| Average Age of Mother | 27.0 |
| ● Teenage Birth Rate ² | 7.9 |
| Percent White Births | 92.1 |
| Percent American Indian Births | 4.2 |
| Percent Unmarried | 37.7 |
| Percent WIC births | 40.0 |
| Percent Breastfeeding at discharge | 75.2 |
| ○ Percent Payment-Private Insurance | 63.4 |
| Percent Payment-Medicaid | 32.9 |
| ● Percent C-Section | 22.2 |

●Denotes a health status indicator which is significantly lower than the state average.
○Denotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

| | |
|---------------------------------------|-------|
| All Causes | 662.7 |
| Heart Disease | 141.5 |
| Malignant Neoplasms (Cancer) | 167.8 |
| Trachea, Bronchus, & Lung | 45.0 |
| Colon, Rectum, & Anus | 19.8 |
| Pancreas | 9.5 |
| Female Breast | 14.9 |
| Leukemia | 9.8 |
| Prostate | 20.4 |
| Accidents | 39.7 |
| Motor Vehicle Accidents | 10.9 |
| Alzheimer's Disease | 38.5 |
| Cerebrovascular Disease | 37.0 |
| Chronic Lower Respiratory Diseases | 43.1 |
| Diabetes Mellitus | 20.5 |
| Influenza & Pneumonia | 17.5 |
| Intentional Self-Harm (Suicide) | 17.1 |
| ● Chronic Liver Disease and Cirrhosis | 5.2 |
| Infant Mortality | 5.18 |

Leading Causes of Death

| Leading Causes of Death | Total Deaths |
|---------------------------------------|--------------|
| 1. Malignant Neoplasms (Cancer) | 282 |
| 2. Heart Disease | 263 |
| 3. Chronic Lower Respiratory Diseases | 77 |
| 4. Alzheimer's Disease | 73 |
| 5. Cerebrovascular Diseases | 69 |
| 6. Accidents | 60 |
| 7. Diabetes | 36 |
| 8. Influenza and Pneumonia | 32 |
| 9. Intentional Self-Harm (Suicide) | 23 |
| 10. Parkinson's Disease | 16 |
| Percent of Deaths due to tobacco use | 21.8 |
| Median age at death | 81 |

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Corson County

Demographic Information



Corson County is located in the north central portion of the state and averages 1.6 persons per square mile.

2013 Population Information

| Subject | Number | Percent |
|--|--------|---------|
| Total population | 4,215 | 100.0 |
| White | 1,324 | 31.4 |
| Black or African American | 9 | 0.2 |
| American Indian & Alaska Native | 2,733 | 64.8 |
| Asian | 14 | 0.3 |
| Native Hawaiian & Other Pacific Islander | 4 | 0.1 |
| Two or More Races | 131 | 3.1 |
| Under 5 years | 365 | 8.7 |
| Under 18 years | 1,427 | 33.9 |
| 65 years and over | 453 | 10.7 |

Source: United States Census Bureau, 2013 Population Estimates

Health Status Indicators 2009-2013

Nativity

| | |
|---|------|
| Percent of Low Birth Weight Infants | 6.3 |
| ● Percent of Mothers Receiving Care in 1st Trimester | 46.2 |
| Percent of Mothers Who Used Tobacco While Pregnant ¹ | 20.3 |
| Percent of Births Less Than 37 Wks. of Gestation | 11.1 |
| ● Average Age of Mother | 25.3 |
| ○ Teenage Birth Rate ² | 34.1 |
| Percent White Births | 22.2 |
| Percent American Indian Births | 75.1 |
| ○ Percent Unmarried | 68.3 |
| ○ Percent WIC births | 77.9 |
| ● Percent Breastfeeding at discharge | 55.2 |
| ● Percent Payment-Private Insurance | 17.0 |
| ○ Percent Payment-Medicaid | 76.9 |
| Percent C-Section | 26.9 |

●Denotes a health status indicator which is significantly lower than the state average.
○Denotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

| | |
|---------------------------------------|---------|
| ○ All Causes | 1,292.3 |
| ○ Heart Disease | 289.2 |
| ○ Malignant Neoplasms (Cancer) | 238.4 |
| Trachea, Bronchus, & Lung | 46.7 |
| Colon, Rectum, & Anus | 35.7 |
| Pancreas | 15.7 |
| Female Breast | 54.7 |
| Leukemia | LNE |
| Prostate | LNE |
| ○ Accidents | 158.7 |
| ○ Motor Vehicle Accidents | 55.2 |
| Alzheimer's Disease | 40.7 |
| Cerebrovascular Disease | 32.3 |
| Chronic Lower Respiratory Diseases | 51.6 |
| Diabetes Mellitus | 56.2 |
| Influenza & Pneumonia | 26.5 |
| ○ Intentional Self-Harm (Suicide) | 77.5 |
| ○ Chronic Liver Disease and Cirrhosis | 58.3 |
| Infant Mortality | 14.42 |

Leading Causes of Death

| Leading Causes of Death | Total Deaths |
|--|--------------|
| 1. Heart Disease | 51 |
| 2. Malignant Neoplasms (Cancer) | 43 |
| 3. Accidents | 31 |
| 4. Intentional Self-Harm (Suicide) | 16 |
| 5. Chronic Liver Disease and Cirrhosis | 12 |
| T6. Chronic Lower Respiratory Diseases | 9 |
| T6. Diabetes | 9 |
| 8. Alzheimer's Disease | 6 |
| T9. Cerebrovascular Diseases | 5 |
| T9. Septicemia | 5 |
| Percent of Deaths due to tobacco use | 17.3 |
| Median age at death | 63 |

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Custer County

Demographic Information



Custer County is located in the southern Black Hills and averages 5.3 persons per square mile.

2013 Population Information

| Subject | Number | Percent |
|--|--------|---------|
| Total population | 8,468 | 100.0 |
| White | 7,891 | 93.2 |
| Black or African American | 60 | 0.7 |
| American Indian & Alaska Native | 292 | 3.4 |
| Asian | 50 | 0.6 |
| Native Hawaiian & Other Pacific Islander | 3 | 0.0 |
| Two or More Races | 172 | 2.0 |
| Under 5 years | 355 | 4.2 |
| Under 18 years | 1,481 | 17.5 |
| 65 years and over | 2,012 | 23.8 |

Source: United States Census Bureau, 2013 Population Estimates

Health Status Indicators 2009-2013

Nativity

| | |
|---|------|
| Percent of Low Birth Weight Infants | 7.8 |
| Percent of Mothers Receiving Care in 1st Trimester | 71.3 |
| Percent of Mothers Who Used Tobacco While Pregnant ¹ | 18.8 |
| Percent of Births Less Than 37 Wks. of Gestation | 8.9 |
| Average Age of Mother | 27.4 |
| Teenage Birth Rate ² | 12.5 |
| Percent White Births | 88.3 |
| Percent American Indian Births | 7.0 |
| • Percent Unmarried | 31.9 |
| Percent WIC births | 36.7 |
| Percent Breastfeeding at discharge | 81.5 |
| Percent Payment-Private Insurance | 51.0 |
| Percent Payment-Medicaid | 39.3 |
| • Percent C-Section | 16.1 |

● Denotes a health status indicator which is significantly lower than the state average.
○ Denotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

| | |
|-------------------------------------|-------|
| All Causes | 634.9 |
| ● Heart Disease | 124.2 |
| Malignant Neoplasms (Cancer) | 163.8 |
| Trachea, Bronchus, & Lung | 35.4 |
| Colon, Rectum, & Anus | 18.2 |
| ● Pancreas | 3.7 |
| Female Breast | 18.4 |
| Leukemia | 7.7 |
| Prostate | 15.1 |
| Accidents | 52.9 |
| Motor Vehicle Accidents | 22.2 |
| ● Alzheimer's Disease | 18.7 |
| Cerebrovascular Disease | 42.0 |
| Chronic Lower Respiratory Diseases | 38.0 |
| ● Diabetes Mellitus | 10.0 |
| Influenza & Pneumonia | 15.4 |
| Intentional Self-Harm (Suicide) | 10.9 |
| Chronic Liver Disease and Cirrhosis | 7.1 |
| Infant Mortality | LNE |

Leading Causes of Death

| Leading Causes of Death | Total Deaths |
|--|--------------|
| 1. Malignant Neoplasms (Cancer) | 111 |
| 2. Heart Disease | 87 |
| 3. Cerebrovascular Diseases | 26 |
| 4. Chronic Lower Respiratory Diseases | 25 |
| 5. Accidents | 24 |
| 6. Alzheimer's Disease | 12 |
| 7. Influenza and Pneumonia | 10 |
| 8. Diabetes | 7 |
| T9. Intentional Self-Harm (Suicide) | 6 |
| T9. Parkinson's Disease | 6 |
| T9. Disorders of Lipoprotein Metabolism and Other Lipidemias | 6 |

| | |
|--------------------------------------|------|
| Percent of Deaths due to tobacco use | 19.9 |
| Median age at death | 77 |

● Denotes a health status indicator which is significantly lower than the state average.

○ Denotes a health status indicator which is significantly higher than the state average.


³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Davison County

Demographic Information



Davison County is located in eastern South Dakota and averages 44.8 persons per square mile.

2013 Population Information

| Subject | Number | Percent |
|--|--------|---------|
| Total population | 19,823 | 100.0 |
| White | 18,680 | 94.2 |
| Black or African American | 208 | 1.0 |
| American Indian & Alaska Native | 523 | 2.6 |
| Asian | 117 | 0.6 |
| Native Hawaiian & Other Pacific Islander | 18 | 0.1 |
| Two or More Races | 277 | 1.4 |
| Under 5 years | 1,262 | 6.4 |
| Under 18 years | 4,584 | 23.1 |
| 65 years and over | 3,411 | 17.2 |

Source: United States Census Bureau, 2013 Population Estimates

Health Status Indicators 2009-2013

Natality

| | |
|---|------|
| Percent of Low Birth Weight Infants | 5.2 |
| ○ Percent of Mothers Receiving Care in 1st Trimester | 76.9 |
| ○ Percent of Mothers Who Used Tobacco While Pregnant ¹ | 20.0 |
| Percent of Births Less Than 37 Wks. of Gestation | 8.3 |
| ● Average Age of Mother | 26.5 |
| ● Teenage Birth Rate ² | 11.4 |
| Percent White Births | 89.7 |
| Percent American Indian Births | 7.5 |
| Percent Unmarried | 40.1 |
| Percent WIC births | 41.9 |
| Percent Breastfeeding at discharge | 69.8 |
| ○ Percent Payment-Private Insurance | 61.7 |
| Percent Payment-Medicaid | 35.3 |
| Percent C-Section | 26.9 |

Mortality³

| | |
|---------------------------------------|-------|
| ● All Causes | 642.3 |
| Heart Disease | 179.4 |
| Malignant Neoplasms (Cancer) | 152.9 |
| Trachea, Bronchus, & Lung | 48.9 |
| Colon, Rectum, & Anus | 13.4 |
| Pancreas | 12.2 |
| ● Female Breast | 9.5 |
| Leukemia | 8.7 |
| Prostate | 12.4 |
| ● Accidents | 31.5 |
| ● Motor Vehicle Accidents | 7.6 |
| ● Alzheimer's Disease | 19.4 |
| Cerebrovascular Disease | 41.3 |
| Chronic Lower Respiratory Diseases | 33.6 |
| Diabetes Mellitus | 15.4 |
| Influenza & Pneumonia | 20.1 |
| Intentional Self-Harm (Suicide) | 14.4 |
| ● Chronic Liver Disease and Cirrhosis | 5.5 |
| Infant Mortality | 5.96 |

Leading Causes of Death

| | Total Deaths |
|--|--------------|
| 1. Heart Disease | 270 |
| 2. Malignant Neoplasms (Cancer) | 205 |
| 3. Cerebrovascular Diseases | 69 |
| 4. Chronic Lower Respiratory Diseases | 51 |
| 5. Influenza and Pneumonia | 40 |
| 6. Alzheimer's Disease | 38 |
| 7. Accidents | 37 |
| 8. Diabetes | 20 |
| 9. Unspecified Dementia | 18 |
| T10. Intentional Self-Harm (Suicide) | 13 |
| T10. Parkinson's Disease | 13 |
| T10. Pneumonitis due to Solids and Liquids | 13 |

| | |
|--------------------------------------|------|
| Percent of Deaths due to tobacco use | 16.9 |
| Median age at death | 82 |

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Day County

Demographic Information



Day County is located in the northeastern part of the state and averages 5.6 persons per square mile.

2013 Population Information

| Subject | Number | Percent |
|--|--------|---------|
| Total population | 5,596 | 100.0 |
| White | 4,953 | 88.5 |
| Black or African American | 30 | 0.5 |
| American Indian & Alaska Native | 497 | 8.9 |
| Asian | 16 | 0.3 |
| Native Hawaiian & Other Pacific Islander | 0 | 0.0 |
| Two or More Races | 100 | 1.8 |
| Under 5 years | 316 | 5.6 |
| Under 18 years | 1,230 | 22.0 |
| 65 years and over | 1,325 | 23.7 |

Source: United States Census Bureau, 2013 Population Estimates

Health Status Indicators 2009-2013

Nativity

| | |
|---|------|
| Percent of Low Birth Weight Infants | 6.9 |
| Percent of Mothers Receiving Care in 1st Trimester | 67.1 |
| ○ Percent of Mothers Who Used Tobacco While Pregnant ¹ | 29.7 |
| Percent of Births Less Than 37 Wks. of Gestation | 8.8 |
| Average Age of Mother | 27.4 |
| Teenage Birth Rate ² | 9.6 |
| Percent White Births | 71.7 |
| Percent American Indian Births | 22.0 |
| Percent Unmarried | 42.2 |
| Percent WIC births | 43.1 |
| Percent Breastfeeding at discharge | 71.2 |
| Percent Payment-Private Insurance | 57.4 |
| Percent Payment-Medicaid | 37.0 |
| Percent C-Section | 25.9 |

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

| | |
|-------------------------------------|-------|
| All Causes | 719.0 |
| ○ Heart Disease | 198.8 |
| Malignant Neoplasms (Cancer) | 152.5 |
| Trachea, Bronchus, & Lung | 44.1 |
| Colon, Rectum, & Anus | 13.9 |
| Pancreas | 11.2 |
| Female Breast | 21.6 |
| Leukemia | LNE |
| Prostate | 47.7 |
| Accidents | 50.6 |
| Motor Vehicle Accidents | 18.0 |
| Alzheimer's Disease | 34.5 |
| Cerebrovascular Disease | 55.9 |
| Chronic Lower Respiratory Diseases | 44.1 |
| Diabetes Mellitus | 24.3 |
| Influenza & Pneumonia | 14.4 |
| ● Intentional Self-Harm (Suicide) | 5.8 |
| Chronic Liver Disease and Cirrhosis | 7.8 |
| Infant Mortality | LNE |

Leading Causes of Death

| Leading Causes of Death | Total Deaths |
|---|--------------|
| 1. Heart Disease | 110 |
| 2. Malignant Neoplasms (Cancer) | 83 |
| 3. Cerebrovascular Diseases | 34 |
| 4. Chronic Lower Respiratory Diseases | 25 |
| 5. Alzheimer's Disease | 23 |
| 6. Accidents | 21 |
| 7. Diabetes | 12 |
| 8. Influenza and Pneumonia | 9 |
| 9. Nephritis, Nephrotic Syndrome, and Nephrosis | 6 |
| Percent of Deaths due to tobacco use | 19.5 |
| Median age at death | 83 |

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Deuel County

Demographic Information



Deuel County borders Minnesota and averages 7.0 persons per square mile.

2013 Population Information

| Subject | Number | Percent |
|--|--------|---------|
| Total population | 4,320 | 100.0 |
| White | 4,217 | 97.6 |
| Black or African American | 44 | 1.0 |
| American Indian & Alaska Native | 17 | 0.4 |
| Asian | 7 | 0.2 |
| Native Hawaiian & Other Pacific Islander | 0 | 0.0 |
| Two or More Races | 35 | 0.8 |
| Under 5 years | 259 | 6.0 |
| Under 18 years | 989 | 22.9 |
| 65 years and over | 889 | 20.6 |

Source: United States Census Bureau, 2013 Population Estimates

Health Status Indicators 2009-2013

Nativity

| | |
|---|------|
| ● Percent of Low Birth Weight Infants | 3.4 |
| ○ Percent of Mothers Receiving Care in 1st Trimester | 83.6 |
| Percent of Mothers Who Used Tobacco While Pregnant ¹ | 17.4 |
| ● Percent of Births Less Than 37 Wks. of Gestation | 3.8 |
| Average Age of Mother | 27.1 |
| Teenage Birth Rate ² | 8.7 |
| Percent White Births | 99.1 |
| Percent American Indian Births | LNE |
| ● Percent Unmarried | 25.2 |
| Percent WIC births | 31.6 |
| Percent Breastfeeding at discharge | 82.1 |
| ○ Percent Payment-Private Insurance | 73.4 |
| ● Percent Payment-Medicaid | 20.2 |
| Percent C-Section | 19.7 |

●Denotes a health status indicator which is significantly lower than the state average.
○Denotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

| | |
|-------------------------------------|-------|
| ● All Causes | 584.2 |
| Heart Disease | 150.8 |
| ● Malignant Neoplasms (Cancer) | 115.7 |
| Trachea, Bronchus, & Lung | 49.0 |
| Colon, Rectum, & Anus | LNE |
| Pancreas | LNE |
| Female Breast | 16.3 |
| Leukemia | LNE |
| Prostate | LNE |
| Accidents | 41.8 |
| Motor Vehicle Accidents | LNE |
| ● Alzheimer's Disease | 19.3 |
| Cerebrovascular Disease | 32.8 |
| Chronic Lower Respiratory Diseases | 53.0 |
| Diabetes Mellitus | 14.3 |
| Influenza & Pneumonia | 13.8 |
| Intentional Self-Harm (Suicide) | 15.6 |
| Chronic Liver Disease and Cirrhosis | LNE |
| Infant Mortality | LNE |

Leading Causes of Death

| Leading Causes of Death | Total Deaths |
|---------------------------------------|--------------|
| 1. Heart Disease | 55 |
| 2. Malignant Neoplasms (Cancer) | 39 |
| 3. Chronic Lower Respiratory Diseases | 18 |
| 4. Cerebrovascular Diseases | 12 |
| T5. Accidents | 11 |
| T5. Septicemia | 11 |
| 7. Alzheimer's Disease | 8 |
| 8. Influenza and Pneumonia | 6 |
| 9. Diabetes | 5 |
| Percent of Deaths due to tobacco use | 20.7 |
| Median age at death | 82 |

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Dewey County

Demographic Information



Dewey County is located in the north central region of the state and averages 2.3 persons per square mile.

2013 Population Information

| Subject | Number | Percent |
|--|--------|---------|
| Total population | 5,586 | 100.0 |
| White | 1,282 | 23.0 |
| Black or African American | 13 | 0.2 |
| American Indian & Alaska Native | 4,071 | 72.9 |
| Asian | 9 | 0.2 |
| Native Hawaiian & Other Pacific Islander | 0 | 0.0 |
| Two or More Races | 211 | 3.8 |
| Under 5 years | 628 | 11.2 |
| Under 18 years | 1,934 | 34.6 |
| 65 years and over | 558 | 10.0 |

Source: United States Census Bureau, 2013 Population Estimates

Health Status Indicators 2009-2013

Nativity

| | |
|---|------|
| Percent of Low Birth Weight Infants | 6.6 |
| ● Percent of Mothers Receiving Care in 1st Trimester | 50.2 |
| ○ Percent of Mothers Who Used Tobacco While Pregnant ¹ | 26.2 |
| Percent of Births Less Than 37 Wks. of Gestation | 8.7 |
| ● Average Age of Mother | 25.1 |
| ○ Teenage Birth Rate ² | 49.1 |
| Percent White Births | 9.5 |
| Percent American Indian Births | 85.4 |
| ○ Percent Unmarried | 70.1 |
| ○ Percent WIC births | 81.8 |
| ● Percent Breastfeeding at discharge | 49.9 |
| ● Percent Payment-Private Insurance | 15.0 |
| ○ Percent Payment-Medicaid | 76.6 |
| ○ Percent C-Section | 32.8 |

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

| | |
|---------------------------------------|---------|
| ○ All Causes | 1,250.7 |
| ○ Heart Disease | 264.5 |
| Malignant Neoplasms (Cancer) | 225.1 |
| Trachea, Bronchus, & Lung | 76.0 |
| Colon, Rectum, & Anus | 28.8 |
| Pancreas | LNE |
| Female Breast | 26.0 |
| Leukemia | 13.1 |
| Prostate | LNE |
| ○ Accidents | 126.1 |
| ○ Motor Vehicle Accidents | 63.9 |
| Alzheimer's Disease | 23.1 |
| Cerebrovascular Disease | 34.7 |
| Chronic Lower Respiratory Diseases | 57.4 |
| ○ Diabetes Mellitus | 93.2 |
| ○ Influenza & Pneumonia | 54.7 |
| ○ Intentional Self-Harm (Suicide) | 56.4 |
| ○ Chronic Liver Disease and Cirrhosis | 51.6 |
| Infant Mortality | 11.76 |

Leading Causes of Death

| Leading Causes of Death | Total Deaths |
|---|--------------|
| 1. Heart Disease | 58 |
| 2. Malignant Neoplasms (Cancer) | 49 |
| 3. Accidents | 30 |
| 4. Diabetes | 21 |
| 5. Intentional Self-Harm (Suicide) | 15 |
| T6. Chronic Lower Respiratory Diseases | 13 |
| T6. Chronic Liver Disease and Cirrhosis | 13 |
| 8. Influenza and Pneumonia | 12 |
| 9. Cerebrovascular Diseases | 7 |
| 10. Liver Disease, Unspecified | 6 |

| | |
|--------------------------------------|------|
| Percent of Deaths due to tobacco use | 21.4 |
| Median age at death | 65 |

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Douglas County

Demographic Information



Douglas County is located in south central South Dakota and averages 7.0 persons per square mile.

2013 Population Information

| Subject | Number | Percent |
|--|--------|---------|
| Total population | 3,023 | 100.0 |
| White | 2,903 | 96.0 |
| Black or African American | 15 | 0.5 |
| American Indian & Alaska Native | 66 | 2.2 |
| Asian | 10 | 0.3 |
| Native Hawaiian & Other Pacific Islander | 0 | 0.0 |
| Two or More Races | 29 | 1.0 |
| Under 5 years | 182 | 6.0 |
| Under 18 years | 685 | 22.7 |
| 65 years and over | 738 | 24.4 |

Source: United States Census Bureau, 2013 Population Estimates

Health Status Indicators 20089-2013

Nativity

| | |
|---|------|
| Percent of Low Birth Weight Infants | 6.6 |
| Percent of Mothers Receiving Care in 1st Trimester | 72.0 |
| ● Percent of Mothers Who Used Tobacco While Pregnant ¹ | 10.2 |
| Percent of Births Less Than 37 Wks. of Gestation | 4.8 |
| Average Age of Mother | 27.8 |
| Teenage Birth Rate ² | 8.8 |
| Percent White Births | 96.4 |
| Percent American Indian Births | LNE |
| ● Percent Unmarried | 15.0 |
| ● Percent WIC births | 27.1 |
| Percent Breastfeeding at discharge | 85.0 |
| Percent Payment-Private Insurance | 64.2 |
| ● Percent Payment-Medicaid | 22.4 |
| Percent C-Section | 28.1 |

●Denotes a health status indicator which is significantly lower than the state average.
○Denotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

| | |
|-------------------------------------|-------|
| All Causes | 668.7 |
| Heart Disease | 155.5 |
| Malignant Neoplasms (Cancer) | 171.1 |
| Trachea, Bronchus, & Lung | 34.2 |
| Colon, Rectum, & Anus | 23.4 |
| Pancreas | 15.6 |
| Female Breast | 25.4 |
| Leukemia | 24.2 |
| Prostate | LNE |
| Accidents | 34.1 |
| Motor Vehicle Accidents | LNE |
| Alzheimer's Disease | 46.5 |
| Cerebrovascular Disease | 43.6 |
| Chronic Lower Respiratory Diseases | 46.5 |
| Diabetes Mellitus | 23.0 |
| Influenza & Pneumonia | 10.6 |
| Intentional Self-Harm (Suicide) | LNE |
| Chronic Liver Disease and Cirrhosis | LNE |
| Infant Mortality | LNE |

Leading Causes of Death

| Leading Causes of Death | Total Deaths |
|---|--------------|
| 1. Heart Disease | 58 |
| 2. Malignant Neoplasms (Cancer) | 44 |
| 3. Alzheimer's Disease | 21 |
| 4. Cerebrovascular Diseases | 17 |
| 5. Chronic Lower Respiratory Diseases | 15 |
| 6. Accidents | 12 |
| 7. Diabetes | 9 |
| T8. Influenza and Pneumonia | 5 |
| T8. Pneumonitis due to Solids and Liquids | 5 |
| Percent of Deaths due to tobacco use | 12.8 |
| Median age at death | 85 |

●Denotes a health status indicator which is significantly lower than the state average.


○Denotes a health status indicator which is significantly higher than the state average.

³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics


Edmunds County

| Demographic Information | Health Status Indicators 2009-2013 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
|--|------------------------------------|---------|---------|------------------|-------|-------|-------|-------|------|---------------------------|----|-----|---------------------------------|----|-----|-------|---|-----|--|---|-----|-------------------|----|--|---------------|-----|-----|----------------|-----|------|-------------------|-----|------|--|-------------------------------------|-----|--|------|---|------|--|-----|-----------------------|------|-----------------------------------|-----|----------------------|------|--------------------------------|-----|---------------------|------|----------------------|------|------------------------------------|------|-------------------------------------|------|----------------------------|------|-------------------|------|------------|-------|---------------|-------|------------------------------|-------|---------------------------|------|-----------------------|------|----------|------|---------------|-----|----------|-----|----------|------|-----------|------|-------------------------|------|---------------------|------|-------------------------|------|------------------------------------|------|---------------------|------|-----------------------|------|---------------------------------|-----|-------------------------------------|-----|------------------|-----|-------------------------|--------------|---------------------------------|----|------------------|----|---------------------------------------|----|-------------|----|------------------------|----|--------------|---|------------------------------|---|---|---|----------------------------|---|
| <div style="text-align: center;">  </div> <p>Edmunds County is located in the north central region of the state and averages 3.6 persons per square mile.</p> <p style="text-align: center;">2013 Population Information</p> <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="text-align: left;">Subject</th> <th style="text-align: right;">Number</th> <th style="text-align: right;">Percent</th> </tr> </thead> <tbody> <tr> <td>Total population</td> <td style="text-align: right;">4,041</td> <td style="text-align: right;">100.0</td> </tr> <tr> <td> White</td> <td style="text-align: right;">3,959</td> <td style="text-align: right;">98.0</td> </tr> <tr> <td> Black or African American</td> <td style="text-align: right;">10</td> <td style="text-align: right;">0.2</td> </tr> <tr> <td> American Indian & Alaska Native</td> <td style="text-align: right;">23</td> <td style="text-align: right;">0.6</td> </tr> <tr> <td> Asian</td> <td style="text-align: right;">7</td> <td style="text-align: right;">0.2</td> </tr> <tr> <td> Native Hawaiian & Other Pacific Islander</td> <td style="text-align: right;">0</td> <td style="text-align: right;">0.0</td> </tr> <tr> <td> Two or More Races</td> <td style="text-align: right;">42</td> <td></td> </tr> <tr> <td>Under 5 years</td> <td style="text-align: right;">226</td> <td style="text-align: right;">5.6</td> </tr> <tr> <td>Under 18 years</td> <td style="text-align: right;">953</td> <td style="text-align: right;">23.6</td> </tr> <tr> <td>65 years and over</td> <td style="text-align: right;">843</td> <td style="text-align: right;">20.9</td> </tr> </tbody> </table> | Subject | Number | Percent | Total population | 4,041 | 100.0 | White | 3,959 | 98.0 | Black or African American | 10 | 0.2 | American Indian & Alaska Native | 23 | 0.6 | Asian | 7 | 0.2 | Native Hawaiian & Other Pacific Islander | 0 | 0.0 | Two or More Races | 42 | | Under 5 years | 226 | 5.6 | Under 18 years | 953 | 23.6 | 65 years and over | 843 | 20.9 | <p style="text-align: center;">Nativity</p> <table border="1" style="width: 100%; border-collapse: collapse;"> <tbody> <tr> <td>Percent of Low Birth Weight Infants</td> <td style="text-align: right;">3.7</td> </tr> <tr> <td>Percent of Mothers Receiving Care in 1st Trimester</td> <td style="text-align: right;">70.0</td> </tr> <tr> <td>Percent of Mothers Who Used Tobacco While Pregnant¹</td> <td style="text-align: right;">12.4</td> </tr> <tr> <td>Percent of Births Less Than 37 Wks. of Gestation</td> <td style="text-align: right;">5.5</td> </tr> <tr> <td>Average Age of Mother</td> <td style="text-align: right;">27.8</td> </tr> <tr> <td>● Teenage Birth Rate²</td> <td style="text-align: right;">6.3</td> </tr> <tr> <td>Percent White Births</td> <td style="text-align: right;">98.2</td> </tr> <tr> <td>Percent American Indian Births</td> <td style="text-align: right;">LNE</td> </tr> <tr> <td>● Percent Unmarried</td> <td style="text-align: right;">20.1</td> </tr> <tr> <td>● Percent WIC births</td> <td style="text-align: right;">22.9</td> </tr> <tr> <td>Percent Breastfeeding at discharge</td> <td style="text-align: right;">81.1</td> </tr> <tr> <td>○ Percent Payment-Private Insurance</td> <td style="text-align: right;">72.9</td> </tr> <tr> <td>● Percent Payment-Medicaid</td> <td style="text-align: right;">19.3</td> </tr> <tr> <td>Percent C-Section</td> <td style="text-align: right;">26.5</td> </tr> </tbody> </table> <p style="text-align: center;">Mortality³</p> <table border="1" style="width: 100%; border-collapse: collapse;"> <tbody> <tr> <td>All Causes</td> <td style="text-align: right;">661.9</td> </tr> <tr> <td>Heart Disease</td> <td style="text-align: right;">126.1</td> </tr> <tr> <td>Malignant Neoplasms (Cancer)</td> <td style="text-align: right;">170.7</td> </tr> <tr> <td> Trachea, Bronchus, & Lung</td> <td style="text-align: right;">33.5</td> </tr> <tr> <td> Colon, Rectum, & Anus</td> <td style="text-align: right;">18.4</td> </tr> <tr> <td> Pancreas</td> <td style="text-align: right;">20.8</td> </tr> <tr> <td> Female Breast</td> <td style="text-align: right;">LNE</td> </tr> <tr> <td> Leukemia</td> <td style="text-align: right;">8.0</td> </tr> <tr> <td> Prostate</td> <td style="text-align: right;">39.2</td> </tr> <tr> <td>Accidents</td> <td style="text-align: right;">34.8</td> </tr> <tr> <td> Motor Vehicle Accidents</td> <td style="text-align: right;">23.5</td> </tr> <tr> <td>Alzheimer's Disease</td> <td style="text-align: right;">26.7</td> </tr> <tr> <td>Cerebrovascular Disease</td> <td style="text-align: right;">24.0</td> </tr> <tr> <td>Chronic Lower Respiratory Diseases</td> <td style="text-align: right;">59.3</td> </tr> <tr> <td> ○ Diabetes Mellitus</td> <td style="text-align: right;">52.5</td> </tr> <tr> <td>Influenza & Pneumonia</td> <td style="text-align: right;">16.9</td> </tr> <tr> <td>Intentional Self-Harm (Suicide)</td> <td style="text-align: right;">LNE</td> </tr> <tr> <td>Chronic Liver Disease and Cirrhosis</td> <td style="text-align: right;">LNE</td> </tr> <tr> <td>Infant Mortality</td> <td style="text-align: right;">LNE</td> </tr> </tbody> </table> <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="text-align: left;">Leading Causes of Death</th> <th style="text-align: right;">Total Deaths</th> </tr> </thead> <tbody> <tr> <td>1. Malignant Neoplasms (Cancer)</td> <td style="text-align: right;">58</td> </tr> <tr> <td>2. Heart Disease</td> <td style="text-align: right;">52</td> </tr> <tr> <td>3. Chronic Lower Respiratory Diseases</td> <td style="text-align: right;">23</td> </tr> <tr> <td>4. Diabetes</td> <td style="text-align: right;">20</td> </tr> <tr> <td>5. Alzheimer's Disease</td> <td style="text-align: right;">12</td> </tr> <tr> <td>6. Accidents</td> <td style="text-align: right;">9</td> </tr> <tr> <td>T7. Cerebrovascular Diseases</td> <td style="text-align: right;">8</td> </tr> <tr> <td>T7. Essential (Primary) Hypertension and Hypertensive Renal Disease</td> <td style="text-align: right;">8</td> </tr> <tr> <td>9. Influenza and Pneumonia</td> <td style="text-align: right;">7</td> </tr> </tbody> </table> <p>Percent of Deaths due to tobacco use: 14.8 Median age at death: 83</p> <p>●Denotes a health status indicator which is significantly lower than the state average. ○Denotes a health status indicator which is significantly higher than the state average.</p> <p>³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births. See technical notes for more information. Source: South Dakota Department of Health, Office of Health Statistics</p> | Percent of Low Birth Weight Infants | 3.7 | Percent of Mothers Receiving Care in 1st Trimester | 70.0 | Percent of Mothers Who Used Tobacco While Pregnant ¹ | 12.4 | Percent of Births Less Than 37 Wks. of Gestation | 5.5 | Average Age of Mother | 27.8 | ● Teenage Birth Rate ² | 6.3 | Percent White Births | 98.2 | Percent American Indian Births | LNE | ● Percent Unmarried | 20.1 | ● Percent WIC births | 22.9 | Percent Breastfeeding at discharge | 81.1 | ○ Percent Payment-Private Insurance | 72.9 | ● Percent Payment-Medicaid | 19.3 | Percent C-Section | 26.5 | All Causes | 661.9 | Heart Disease | 126.1 | Malignant Neoplasms (Cancer) | 170.7 | Trachea, Bronchus, & Lung | 33.5 | Colon, Rectum, & Anus | 18.4 | Pancreas | 20.8 | Female Breast | LNE | Leukemia | 8.0 | Prostate | 39.2 | Accidents | 34.8 | Motor Vehicle Accidents | 23.5 | Alzheimer's Disease | 26.7 | Cerebrovascular Disease | 24.0 | Chronic Lower Respiratory Diseases | 59.3 | ○ Diabetes Mellitus | 52.5 | Influenza & Pneumonia | 16.9 | Intentional Self-Harm (Suicide) | LNE | Chronic Liver Disease and Cirrhosis | LNE | Infant Mortality | LNE | Leading Causes of Death | Total Deaths | 1. Malignant Neoplasms (Cancer) | 58 | 2. Heart Disease | 52 | 3. Chronic Lower Respiratory Diseases | 23 | 4. Diabetes | 20 | 5. Alzheimer's Disease | 12 | 6. Accidents | 9 | T7. Cerebrovascular Diseases | 8 | T7. Essential (Primary) Hypertension and Hypertensive Renal Disease | 8 | 9. Influenza and Pneumonia | 7 |
| Subject | Number | Percent | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Total population | 4,041 | 100.0 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| White | 3,959 | 98.0 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Black or African American | 10 | 0.2 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| American Indian & Alaska Native | 23 | 0.6 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Asian | 7 | 0.2 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Native Hawaiian & Other Pacific Islander | 0 | 0.0 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Two or More Races | 42 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Under 5 years | 226 | 5.6 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Under 18 years | 953 | 23.6 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 65 years and over | 843 | 20.9 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Percent of Low Birth Weight Infants | 3.7 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Percent of Mothers Receiving Care in 1st Trimester | 70.0 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Percent of Mothers Who Used Tobacco While Pregnant ¹ | 12.4 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Percent of Births Less Than 37 Wks. of Gestation | 5.5 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Average Age of Mother | 27.8 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| ● Teenage Birth Rate ² | 6.3 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Percent White Births | 98.2 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Percent American Indian Births | LNE | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| ● Percent Unmarried | 20.1 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| ● Percent WIC births | 22.9 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Percent Breastfeeding at discharge | 81.1 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| ○ Percent Payment-Private Insurance | 72.9 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| ● Percent Payment-Medicaid | 19.3 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Percent C-Section | 26.5 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| All Causes | 661.9 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Heart Disease | 126.1 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Malignant Neoplasms (Cancer) | 170.7 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Trachea, Bronchus, & Lung | 33.5 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Colon, Rectum, & Anus | 18.4 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Pancreas | 20.8 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Female Breast | LNE | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Leukemia | 8.0 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Prostate | 39.2 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Accidents | 34.8 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Motor Vehicle Accidents | 23.5 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Alzheimer's Disease | 26.7 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Cerebrovascular Disease | 24.0 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Chronic Lower Respiratory Diseases | 59.3 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| ○ Diabetes Mellitus | 52.5 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Influenza & Pneumonia | 16.9 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Intentional Self-Harm (Suicide) | LNE | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Chronic Liver Disease and Cirrhosis | LNE | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Infant Mortality | LNE | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Leading Causes of Death | Total Deaths | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 1. Malignant Neoplasms (Cancer) | 58 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 2. Heart Disease | 52 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 3. Chronic Lower Respiratory Diseases | 23 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 4. Diabetes | 20 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 5. Alzheimer's Disease | 12 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 6. Accidents | 9 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| T7. Cerebrovascular Diseases | 8 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| T7. Essential (Primary) Hypertension and Hypertensive Renal Disease | 8 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 9. Influenza and Pneumonia | 7 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |

Source: United States Census Bureau, 2013 Population Estimates

Fall River County

Demographic Information



Fall River County is located in the southwestern corner of the state and averages 4.1 persons per square mile.

2013 Population Information

| Subject | Number | Percent |
|--|--------|---------|
| Total population | 6,839 | 100.0 |
| White | 6,032 | 88.2 |
| Black or African American | 82 | 1.2 |
| American Indian & Alaska Native | 479 | 7.0 |
| Asian | 40 | 0.6 |
| Native Hawaiian & Other Pacific Islander | 5 | 0.1 |
| Two or More Races | 201 | 2.9 |
| Under 5 years | 234 | 3.4 |
| Under 18 years | 1,184 | 17.3 |
| 65 years and over | 1,774 | 25.9 |

Source: United States Census Bureau, 2013 Population Estimates

Health Status Indicators 2009-2013

Nativity

| | |
|---|------|
| Percent of Low Birth Weight Infants | 8.4 |
| Percent of Mothers Receiving Care in 1st Trimester | 62.4 |
| ○ Percent of Mothers Who Used Tobacco While Pregnant ¹ | 23.4 |
| Percent of Births Less Than 37 Wks. of Gestation | 8.8 |
| Average Age of Mother | 26.8 |
| Teenage Birth Rate ² | 17.8 |
| Percent White Births | 82.1 |
| Percent American Indian Births | 9.2 |
| Percent Unmarried | 40.9 |
| ○ Percent WIC births | 51.8 |
| Percent Breastfeeding at discharge | 77.7 |
| ● Percent Payment-Private Insurance | 45.3 |
| ○ Percent Payment-Medicaid | 45.6 |
| ● Percent C-Section | 17.9 |

●Denotes a health status indicator which is significantly lower than the state average.
○Denotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

| | |
|--------------------------------------|-------|
| ○ All Causes | 868.7 |
| ○ Heart Disease | 202.7 |
| Malignant Neoplasms (Cancer) | 186.6 |
| Trachea, Bronchus, & Lung | 56.0 |
| Colon, Rectum, & Anus | 13.8 |
| Pancreas | 7.3 |
| Female Breast | 31.9 |
| Leukemia | LNE |
| Prostate | 22.6 |
| ○ Accidents | 78.2 |
| Motor Vehicle Accidents | 24.6 |
| ● Alzheimer's Disease | 23.0 |
| ● Cerebrovascular Disease | 23.3 |
| ○ Chronic Lower Respiratory Diseases | 84.2 |
| ○ Diabetes Mellitus | 46.8 |
| Influenza & Pneumonia | 14.6 |
| ● Intentional Self-Harm (Suicide) | 7.7 |
| Chronic Liver Disease and Cirrhosis | 15.2 |
| Infant Mortality | LNE |

Leading Causes of Death

| | Total Deaths |
|---|--------------|
| 1. Heart Disease | 134 |
| 2. Malignant Neoplasms (Cancer) | 123 |
| 3. Chronic Lower Respiratory Diseases | 59 |
| 4. Accidents | 32 |
| 5. Diabetes | 30 |
| T6. Alzheimer's Disease | 17 |
| T6. Cerebrovascular Diseases | 17 |
| 8. Influenza and Pneumonia | 11 |
| T9. Unspecified Dementia | 10 |
| T9. Septicemia | 10 |
| T9. Pneumonitis due to Solids and Liquids | 10 |
| Liquids | |
| Percent of Deaths due to tobacco use | 21.4 |
| Median age at death | 79 |

●Denotes a health status indicator which is significantly lower than the state average.


○Denotes a health status indicator which is significantly higher than the state average.

³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics


Faulk County

| Demographic Information | Health Status Indicators 2009-2013 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
|---|------------------------------------|--|---------|------------------|-------|-------|-------|-------|------|---------------------------|----|-----|---------------------------------|---|-----|-------|---|-----|--|---|-----|-------------------|----|-----|---------------|-----|-----|----------------|-----|------|-------------------|-----|------|---|--|-------|--------------|---------------------------------|----|------------------|----|------------------------|----|---------------------------------------|----|-------------|---|-----------------------------|---|----------------------|---|
| <div style="text-align: center;">  </div> <p>Faulk County is located in north central South Dakota and averages 2.4 persons per square mile.</p> <h3 style="text-align: center;">2013 Population Information</h3> <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="text-align: left;">Subject</th> <th style="text-align: right;">Number</th> <th style="text-align: right;">Percent</th> </tr> </thead> <tbody> <tr> <td>Total population</td> <td style="text-align: right;">2,386</td> <td style="text-align: right;">100.0</td> </tr> <tr> <td>White</td> <td style="text-align: right;">2,345</td> <td style="text-align: right;">98.3</td> </tr> <tr> <td>Black or African American</td> <td style="text-align: right;">10</td> <td style="text-align: right;">0.4</td> </tr> <tr> <td>American Indian & Alaska Native</td> <td style="text-align: right;">3</td> <td style="text-align: right;">0.1</td> </tr> <tr> <td>Asian</td> <td style="text-align: right;">7</td> <td style="text-align: right;">0.3</td> </tr> <tr> <td>Native Hawaiian & Other Pacific Islander</td> <td style="text-align: right;">0</td> <td style="text-align: right;">0.0</td> </tr> <tr> <td>Two or More Races</td> <td style="text-align: right;">21</td> <td style="text-align: right;">0.9</td> </tr> <tr> <td>Under 5 years</td> <td style="text-align: right;">145</td> <td style="text-align: right;">6.1</td> </tr> <tr> <td>Under 18 years</td> <td style="text-align: right;">541</td> <td style="text-align: right;">22.7</td> </tr> <tr> <td>65 years and over</td> <td style="text-align: right;">557</td> <td style="text-align: right;">23.3</td> </tr> </tbody> </table> | Subject | Number | Percent | Total population | 2,386 | 100.0 | White | 2,345 | 98.3 | Black or African American | 10 | 0.4 | American Indian & Alaska Native | 3 | 0.1 | Asian | 7 | 0.3 | Native Hawaiian & Other Pacific Islander | 0 | 0.0 | Two or More Races | 21 | 0.9 | Under 5 years | 145 | 6.1 | Under 18 years | 541 | 22.7 | 65 years and over | 557 | 23.3 | <h3 style="text-align: center;">Natality</h3> <ul style="list-style-type: none"> ● Percent of Low Birth Weight Infants 3.0 ○ Percent of Mothers Receiving Care in 1st Trimester 57.7 ● Percent of Mothers Who Used Tobacco While Pregnant¹ 5.3 ● Percent of Births Less Than 37 Wks. of Gestation 3.8 ○ Average Age of Mother 29.5 Teenage Birth Rate² LNE Percent White Births 100.0 Percent American Indian Births LNE ● Percent Unmarried 10.6 ● Percent WIC births 15.3 Percent Breastfeeding at discharge 87.8 ○ Percent Payment-Private Insurance 79.2 ● Percent Payment-Medicaid 10.8 Percent C-Section 18.9 | <h3 style="text-align: center;">Mortality³</h3> <ul style="list-style-type: none"> ● All Causes 525.3 ● Heart Disease 97.0 Malignant Neoplasms (Cancer) 140.1 <ul style="list-style-type: none"> Trachea, Bronchus, & Lung 26.9 Colon, Rectum, & Anus 9.0 Pancreas LNE Female Breast LNE Leukemia LNE Prostate LNE ● Accidents 9.0 <ul style="list-style-type: none"> Motor Vehicle Accidents LNE Alzheimer's Disease 43.0 Cerebrovascular Disease 22.6 Chronic Lower Respiratory Diseases 40.7 Diabetes Mellitus 30.5 Influenza & Pneumonia 18.5 Intentional Self-Harm (Suicide) LNE Chronic Liver Disease and Cirrhosis LNE Infant Mortality LNE <h3 style="text-align: center;">Leading Causes of Death</h3> <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="text-align: left;">Cause</th> <th style="text-align: right;">Total Deaths</th> </tr> </thead> <tbody> <tr> <td>1. Malignant Neoplasms (Cancer)</td> <td style="text-align: right;">31</td> </tr> <tr> <td>2. Heart Disease</td> <td style="text-align: right;">24</td> </tr> <tr> <td>3. Alzheimer's Disease</td> <td style="text-align: right;">14</td> </tr> <tr> <td>4. Chronic Lower Respiratory Diseases</td> <td style="text-align: right;">10</td> </tr> <tr> <td>5. Diabetes</td> <td style="text-align: right;">8</td> </tr> <tr> <td>6. Cerebrovascular Diseases</td> <td style="text-align: right;">6</td> </tr> <tr> <td>7. Vascular Dementia</td> <td style="text-align: right;">5</td> </tr> </tbody> </table> <ul style="list-style-type: none"> Percent of Deaths due to tobacco use 9.8 Median age at death 85 | Cause | Total Deaths | 1. Malignant Neoplasms (Cancer) | 31 | 2. Heart Disease | 24 | 3. Alzheimer's Disease | 14 | 4. Chronic Lower Respiratory Diseases | 10 | 5. Diabetes | 8 | 6. Cerebrovascular Diseases | 6 | 7. Vascular Dementia | 5 |
| Subject | Number | Percent | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Total population | 2,386 | 100.0 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| White | 2,345 | 98.3 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Black or African American | 10 | 0.4 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| American Indian & Alaska Native | 3 | 0.1 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Asian | 7 | 0.3 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Native Hawaiian & Other Pacific Islander | 0 | 0.0 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Two or More Races | 21 | 0.9 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Under 5 years | 145 | 6.1 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Under 18 years | 541 | 22.7 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 65 years and over | 557 | 23.3 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Cause | Total Deaths | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 1. Malignant Neoplasms (Cancer) | 31 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 2. Heart Disease | 24 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 3. Alzheimer's Disease | 14 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 4. Chronic Lower Respiratory Diseases | 10 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 5. Diabetes | 8 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 6. Cerebrovascular Diseases | 6 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 7. Vascular Dementia | 5 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| <p>●Denotes a health status indicator which is significantly lower than the state average. ○Denotes a health status indicator which is significantly higher than the state average.</p> <p>¹Data for mothers who used tobacco are self-reported. ²Teenage Birth rate is live births per 1,000 females age 15-17.</p> | | <p>●Denotes a health status indicator which is significantly lower than the state average. ○Denotes a health status indicator which is significantly higher than the state average. ³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.</p> <p>See technical notes for more information. Source: South Dakota Department of Health, Office of Health Statistics</p> | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |

Source: United States Census Bureau, 2013 Population Estimates

Grant County

Demographic Information



Grant County borders Minnesota in northeastern South Dakota and averages 10.8 persons per square mile.

2013 Population Information

| Subject | Number | Percent |
|--|--------|---------|
| Total population | 7,281 | 100.0 |
| White | 7,073 | 97.1 |
| Black or African American | 33 | 0.5 |
| American Indian & Alaska Native | 63 | 0.9 |
| Asian | 34 | 0.5 |
| Native Hawaiian & Other Pacific Islander | 2 | 0.0 |
| Two or More Races | 76 | 1.0 |
| Under 5 years | 412 | 5.7 |
| Under 18 years | 1,641 | 22.5 |
| 65 years and over | 1,441 | 19.8 |

Source: United States Census Bureau, 2013 Population Estimates

Health Status Indicators 2009-2013

Nativity

| | |
|---|------|
| Percent of Low Birth Weight Infants | 4.6 |
| Percent of Mothers Receiving Care in 1st Trimester | 76.4 |
| Percent of Mothers Who Used Tobacco While Pregnant ¹ | 13.8 |
| ● Percent of Births Less Than 37 Wks. of Gestation | 5.1 |
| Average Age of Mother | 27.6 |
| Teenage Birth Rate ² | 14.5 |
| Percent White Births | 96.7 |
| Percent American Indian Births | 0.8 |
| ● Percent Unmarried | 29.9 |
| Percent WIC births | 36.9 |
| Percent Breastfeeding at discharge | 74.2 |
| Percent Payment-Private Insurance | 60.5 |
| ● Percent Payment-Medicaid | 22.1 |
| Percent C-Section | 23.6 |

●Denotes a health status indicator which is significantly lower than the state average.
○Denotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

| | |
|-------------------------------------|-------|
| All Causes | 680.2 |
| Heart Disease | 147.0 |
| Malignant Neoplasms (Cancer) | 141.5 |
| Trachea, Bronchus, & Lung | 32.6 |
| Colon, Rectum, & Anus | 15.6 |
| Pancreas | 9.2 |
| Female Breast | 11.2 |
| Leukemia | LNE |
| Prostate | 21.0 |
| Accidents | 36.7 |
| Motor Vehicle Accidents | 15.8 |
| Alzheimer's Disease | 36.2 |
| Cerebrovascular Disease | 51.6 |
| Chronic Lower Respiratory Diseases | 43.3 |
| Diabetes Mellitus | 32.6 |
| Influenza & Pneumonia | 14.7 |
| Intentional Self-Harm (Suicide) | LNE |
| Chronic Liver Disease and Cirrhosis | 7.7 |
| Infant Mortality | LNE |

Leading Causes of Death

| Leading Causes of Death | Total Deaths |
|--|--------------|
| 1. Heart Disease | 90 |
| 2. Malignant Neoplasms (Cancer) | 78 |
| 3. Cerebrovascular Diseases | 35 |
| T4. Chronic Lower Respiratory Diseases | 25 |
| T4. Alzheimer's Disease | 25 |
| 6. Diabetes | 19 |
| 7. Accidents | 16 |
| 8. Influenza and Pneumonia | 9 |
| 9. Disorders of Lipoprotein Metabolism and Other Lipidemias | 6 |
| T10. Essential (Primary) Hypertension and Hypertensive Renal Disease | 5 |
| T10. Nephritis, Nephrotic Syndrome, and Nephrosis | 5 |
| T10. Pneumonitis due to Solids and Liquids | 5 |
| T10. Aortic Aneurysm and Dissection | 5 |
| Percent of Deaths due to tobacco use | 13.8 |
| Median age at death | 82 |

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Gregory County

Demographic Information



Gregory County borders the west bank of the Missouri River and the state of Nebraska and averages 4.2 persons per square mile.

2013 Population Information

| Subject | Number | Percent |
|--|--------|---------|
| Total population | 4,242 | 100.0 |
| White | 3,786 | 89.3 |
| Black or African American | 15 | 0.4 |
| American Indian & Alaska Native | 317 | 7.5 |
| Asian | 19 | 0.4 |
| Native Hawaiian & Other Pacific Islander | 10 | 0.2 |
| Two or More Races | 95 | 2.2 |
| Under 5 years | 245 | 5.8 |
| Under 18 years | 958 | 22.6 |
| 65 years and over | 1,023 | 24.1 |

Source: United States Census Bureau, 2013 Population Estimates

Health Status Indicators 2009-2013

Natality

| | |
|---|------|
| Percent of Low Birth Weight Infants | 5.4 |
| Percent of Mothers Receiving Care in 1st Trimester | 60.3 |
| Percent of Mothers Who Used Tobacco While Pregnant ¹ | 18.8 |
| Percent of Births Less Than 37 Wks. of Gestation | 8.0 |
| Average Age of Mother | 27.5 |
| Teenage Birth Rate ² | 12.0 |
| Percent White Births | 79.9 |
| Percent American Indian Births | 15.2 |
| Percent Unmarried | 37.5 |
| ○ Percent WIC births | 50.2 |
| Percent Breastfeeding at discharge | 71.4 |
| Percent Payment-Private Insurance | 50.0 |
| ○ Percent Payment-Medicaid | 44.2 |
| Percent C-Section | 33.5 |

●Denotes a health status indicator which is significantly lower than the state average.
○Denotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

| | |
|-------------------------------------|-------|
| All Causes | 771.3 |
| ○ Heart Disease | 227.3 |
| Malignant Neoplasms (Cancer) | 169.9 |
| Trachea, Bronchus, & Lung | 57.6 |
| Colon, Rectum, & Anus | 15.0 |
| Pancreas | LNE |
| Female Breast | LNE |
| Leukemia | LNE |
| Prostate | 14.4 |
| Accidents | 64.6 |
| Motor Vehicle Accidents | 36.2 |
| Alzheimer's Disease | 21.2 |
| Cerebrovascular Disease | 35.1 |
| Chronic Lower Respiratory Diseases | 55.7 |
| Diabetes Mellitus | 41.1 |
| ● Influenza & Pneumonia | 5.8 |
| Intentional Self-Harm (Suicide) | 15.8 |
| Chronic Liver Disease and Cirrhosis | 16.5 |
| Infant Mortality | LNE |

Leading Causes of Death

| Leading Causes of Death | Total Deaths |
|---|--------------|
| 1. Heart Disease | 103 |
| 2. Malignant Neoplasms (Cancer) | 67 |
| 3. Chronic Lower Respiratory Diseases | 26 |
| T4. Accidents | 18 |
| T4. Diabetes | 18 |
| 6. Cerebrovascular Diseases | 14 |
| 7. Alzheimer's Disease | 12 |
| 8. Disorders of Lipoprotein Metabolism and Other Lipidemias | 10 |

| | |
|--------------------------------------|------|
| Percent of Deaths due to tobacco use | 18.6 |
| Median age at death | 83 |

●Denotes a health status indicator which is significantly lower than the state average.
○Denotes a health status indicator which is significantly higher than the state average.


³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Haakon County

Demographic Information



Haakon County is located in the west central region of the state and averages 1.1 persons per square mile.

2013 Population Information

| Subject | Number | Percent |
|--|--------|---------|
| Total population | 1,894 | 100.0 |
| White | 1,768 | 93.3 |
| Black or African American | 15 | 0.8 |
| American Indian & Alaska Native | 41 | 2.2 |
| Asian | 9 | 0.5 |
| Native Hawaiian & Other Pacific Islander | 1 | 0.1 |
| Two or More Races | 60 | 3.2 |
| Under 5 years | 117 | 6.2 |
| Under 18 years | 417 | 22.0 |
| 65 years and over | 441 | 23.3 |

Source: United States Census Bureau, 2013 Population Estimates

Health Status Indicators 2009-2013

Nativity

| | |
|---|------|
| ● Percent of Low Birth Weight Infants | 2.7 |
| Percent of Mothers Receiving Care in 1st Trimester | 68.2 |
| ● Percent of Mothers Who Used Tobacco While Pregnant ¹ | 10.1 |
| Percent of Births Less Than 37 Wks. of Gestation | 4.5 |
| Average Age of Mother | 27.9 |
| Teenage Birth Rate ² | 16.9 |
| Percent White Births | 91.8 |
| Percent American Indian Births | 2.7 |
| ● Percent Unmarried | 22.7 |
| ● Percent WIC births | 26.9 |
| Percent Breastfeeding at discharge | 86.2 |
| ○ Percent Payment-Private Insurance | 77.3 |
| ● Percent Payment-Medicaid | 17.3 |
| Percent C-Section | 27.3 |

●Denotes a health status indicator which is significantly lower than the state average.
○Denotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

| | |
|-------------------------------------|-------|
| All Causes | 708.3 |
| Heart Disease | 144.8 |
| Malignant Neoplasms (Cancer) | 178.2 |
| Trachea, Bronchus, & Lung | 23.6 |
| Colon, Rectum, & Anus | 34.2 |
| Pancreas | LNE |
| Female Breast | LNE |
| Leukemia | LNE |
| Prostate | 70.8 |
| Accidents | 27.8 |
| Motor Vehicle Accidents | 20.0 |
| Alzheimer's Disease | LNE |
| Cerebrovascular Disease | 57.8 |
| Chronic Lower Respiratory Diseases | 33.6 |
| Diabetes Mellitus | LNE |
| ○ Influenza & Pneumonia | 53.2 |
| Intentional Self-Harm (Suicide) | LNE |
| Chronic Liver Disease and Cirrhosis | LNE |
| Infant Mortality | LNE |

Leading Causes of Death

| Leading Causes of Death | Total Deaths |
|---|--------------|
| 1. Malignant Neoplasms (Cancer) | 28 |
| 2. Heart Disease | 27 |
| T3. Cerebrovascular Diseases | 11 |
| T3. Influenza and Pneumonia | 11 |
| T5. Chronic Lower Respiratory Diseases | 6 |
| T5. Pneumonitis due to Solids and Liquids | 6 |
| 7. Accidents | 5 |
| Percent of Deaths due to tobacco use | 11.2 |
| Median age at death | 85 |

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Hamlin County

Demographic Information



Hamlin County is located in the northeastern region of the state and averages 11.6 persons per square mile.

2013 Population Information

| Subject | Number | Percent |
|--|--------|---------|
| Total population | 5,961 | 100.0 |
| White | 5,847 | 98.1 |
| Black or African American | 31 | 0.5 |
| American Indian & Alaska Native | 29 | 0.5 |
| Asian | 13 | 0.2 |
| Native Hawaiian & Other Pacific Islander | 1 | 0.0 |
| Two or More Races | 40 | 0.7 |
| Under 5 years | 537 | 9.0 |
| Under 18 years | 1,833 | 30.7 |
| 65 years and over | 1,026 | 17.2 |

Source: United States Census Bureau, 2013 Population Estimates

Health Status Indicators 2009-2013

Nativity

| | |
|---|------|
| ● Percent of Low Birth Weight Infants | 3.7 |
| Percent of Mothers Receiving Care in 1st Trimester | 69.0 |
| Percent of Mothers Who Used Tobacco While Pregnant ¹ | 15.3 |
| ● Percent of Births Less Than 37 Wks. of Gestation | 4.5 |
| Average Age of Mother | 27.4 |
| Teenage Birth Rate ² | 15.7 |
| Percent White Births | 99.3 |
| Percent American Indian Births | LNE |
| ● Percent Unmarried | 13.3 |
| Percent WIC births | 40.6 |
| ○ Percent Breastfeeding at discharge | 87.3 |
| ○ Percent Payment-Private Insurance | 72.2 |
| ● Percent Payment-Medicaid | 23.2 |
| ● Percent C-Section | 16.7 |

Mortality³

| | |
|-------------------------------------|-------|
| All Causes | 789.3 |
| Heart Disease | 155.8 |
| Malignant Neoplasms (Cancer) | 150.6 |
| Trachea, Bronchus, & Lung | 26.2 |
| Colon, Rectum, & Anus | 19.6 |
| Pancreas | 12.4 |
| Female Breast | 11.9 |
| Leukemia | 8.6 |
| Prostate | 43.6 |
| Accidents | 40.9 |
| Motor Vehicle Accidents | 11.5 |
| ○ Alzheimer's Disease | 115.8 |
| Cerebrovascular Disease | 56.5 |
| Chronic Lower Respiratory Diseases | 47.5 |
| Diabetes Mellitus | 29.3 |
| Influenza & Pneumonia | 15.7 |
| Intentional Self-Harm (Suicide) | LNE |
| Chronic Liver Disease and Cirrhosis | LNE |
| Infant Mortality | LNE |

Leading Causes of Death

| Leading Causes of Death | Total Deaths |
|---------------------------------------|--------------|
| 1. Heart Disease | 72 |
| 2. Malignant Neoplasms (Cancer) | 60 |
| 3. Alzheimer's Disease | 58 |
| 4. Cerebrovascular Diseases | 25 |
| 5. Chronic Lower Respiratory Diseases | 22 |
| 6. Accidents | 15 |
| 7. Diabetes | 13 |
| 8. Vascular Dementia | 11 |
| 9. Parkinson's Disease | 8 |
| 10. Influenza and Pneumonia | 7 |
| Percent of Deaths due to tobacco use | 13.2 |
| Median age at death | 84 |

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Hand County

Demographic Information



Hand County is located in central South Dakota and averages 2.4 persons per square mile.

2013 Population Information

| Subject | Number | Percent |
|--|--------|---------|
| Total population | 3,391 | 100.0 |
| White | 3,323 | 98.0 |
| Black or African American | 4 | 0.1 |
| American Indian & Alaska Native | 21 | 0.6 |
| Asian | 12 | 0.4 |
| Native Hawaiian & Other Pacific Islander | 1 | 0.0 |
| Two or More Races | 30 | 0.9 |
| Under 5 years | 172 | 5.1 |
| Under 18 years | 693 | 20.4 |
| 65 years and over | 846 | 24.9 |

Source: United States Census Bureau, 2013 Population Estimates

Health Status Indicators 2009-2013

Nativity

| | |
|---|------|
| Percent of Low Birth Weight Infants | 6.9 |
| Percent of Mothers Receiving Care in 1st Trimester | 64.2 |
| ● Percent of Mothers Who Used Tobacco While Pregnant ¹ | 9.8 |
| Percent of Births Less Than 37 Wks. of Gestation | 9.1 |
| Average Age of Mother | 27.6 |
| Teenage Birth Rate ² | LNE |
| Percent White Births | 98.3 |
| Percent American Indian Births | 1.7 |
| ● Percent Unmarried | 20.6 |
| ● Percent WIC births | 28.0 |
| Percent Breastfeeding at discharge | 78.0 |
| ○ Percent Payment-Private Insurance | 77.7 |
| ● Percent Payment-Medicaid | 18.9 |
| ○ Percent C-Section | 37.7 |

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

| | |
|-------------------------------------|-------|
| ● All Causes | 544.3 |
| ● Heart Disease | 107.2 |
| Malignant Neoplasms (Cancer) | 128.1 |
| Trachea, Bronchus, & Lung | 24.7 |
| Colon, Rectum, & Anus | 10.1 |
| Pancreas | 13.8 |
| Female Breast | LNE |
| Leukemia | 10.7 |
| Prostate | LNE |
| ● Accidents | 23.0 |
| Motor Vehicle Accidents | LNE |
| ● Alzheimer's Disease | 13.5 |
| Cerebrovascular Disease | 45.5 |
| Chronic Lower Respiratory Diseases | 46.2 |
| Diabetes Mellitus | 22.5 |
| Influenza & Pneumonia | 8.0 |
| Intentional Self-Harm (Suicide) | LNE |
| Chronic Liver Disease and Cirrhosis | LNE |
| Infant Mortality | LNE |

Leading Causes of Death

| Leading Causes of Death | Total Deaths |
|---|--------------|
| 1. Heart Disease | 47 |
| 2. Malignant Neoplasms (Cancer) | 42 |
| 3. Cerebrovascular Diseases | 20 |
| 4. Chronic Lower Respiratory Diseases | 16 |
| 5. Atherosclerosis | 15 |
| 6. Diabetes | 10 |
| T7. Accidents | 6 |
| T7. Essential (Primary) Hypertension and Hypertensive Renal Disease | 6 |
| 9. Alzheimer's Disease | 5 |
| Percent of Deaths due to tobacco use | 21.0 |
| Median age at death | 84 |

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Hanson County

Demographic Information



Hanson County is located in southeastern South Dakota and averages 7.7 people per square mile.

2013 Population Information

| Subject | Number | Percent |
|--|--------|---------|
| Total population | 3,405 | 100.0 |
| White | 3,355 | 98.5 |
| Black or African American | 6 | 0.2 |
| American Indian & Alaska Native | 12 | 0.4 |
| Asian | 13 | 0.4 |
| Native Hawaiian & Other Pacific Islander | 2 | 0.1 |
| Two or More Races | 17 | 0.5 |
| Under 5 years | 324 | 9.5 |
| Under 18 years | 1,099 | 32.3 |
| 65 years and over | 469 | 13.8 |

Source: United States Census Bureau, 2013 Population Estimates

Health Status Indicators 2009-2013

Nativity

| | |
|---|------|
| Percent of Low Birth Weight Infants | 8.0 |
| Percent of Mothers Receiving Care in 1st Trimester | 62.5 |
| ● Percent of Mothers Who Used Tobacco While Pregnant ¹ | 8.0 |
| Percent of Births Less Than 37 Wks. of Gestation | 7.2 |
| ○ Average Age of Mother | 29.9 |
| Teenage Birth Rate ² | LNE |
| Percent White Births | 99.6 |
| Percent American Indian Births | LNE |
| ● Percent Unmarried | 12.4 |
| ● Percent WIC births | 18.5 |
| Percent Breastfeeding at discharge | 81.0 |
| ○ Percent Payment-Private Insurance | 79.4 |
| ● Percent Payment-Medicaid | 10.5 |
| Percent C-Section | 30.0 |

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

| | |
|--------------------------------------|-------|
| All Causes | 587.0 |
| Heart Disease | 159.6 |
| Malignant Neoplasms (Cancer) | 146.7 |
| Trachea, Bronchus, & Lung | 58.2 |
| Colon, Rectum, & Anus | LNE |
| Pancreas | LNE |
| Female Breast | LNE |
| Leukemia | LNE |
| Prostate | LNE |
| Accidents | 54.8 |
| Motor Vehicle Accidents | 16.2 |
| Alzheimer's Disease | LNE |
| Cerebrovascular Disease | 43.9 |
| ● Chronic Lower Respiratory Diseases | 18.4 |
| Diabetes Mellitus | LNE |
| Influenza & Pneumonia | LNE |
| Intentional Self-Harm (Suicide) | LNE |
| Chronic Liver Disease and Cirrhosis | LNE |
| Infant Mortality | 24.00 |

Leading Causes of Death

| Leading Causes of Death | Total Deaths |
|---|--------------|
| 1. Malignant Neoplasms (Cancer) | 26 |
| 2. Heart Disease | 24 |
| 3. Accidents | 8 |
| 4. Cerebrovascular Diseases | 6 |
| 5. Certain Conditions Originating in the Perinatal Period | 5 |
| Percent of Deaths due to tobacco use | 12.4 |
| Median age at death | 74 |

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Harding County

Demographic Information



Harding County is located in the northwestern corner of the state and averages 0.5 persons per square mile.

2013 Population Information

| Subject | Number | Percent |
|--|--------|---------|
| Total population | 1,262 | 100.0 |
| White | 1,206 | 95.6 |
| Black or African American | 7 | 0.6 |
| American Indian & Alaska Native | 23 | 1.8 |
| Asian | 2 | 0.2 |
| Native Hawaiian & Other Pacific Islander | 0 | 0.0 |
| Two or More Races | 24 | 1.9 |
| Under 5 years | 89 | 7.1 |
| Under 18 years | 271 | 21.5 |
| 65 years and over | 197 | 15.6 |

Source: United States Census Bureau, 2013 Population Estimates

Health Status Indicators 2009-2013

Nativity

| | |
|---|------|
| Percent of Low Birth Weight Infants | LNE |
| Percent of Mothers Receiving Care in 1st Trimester | 73.4 |
| ● Percent of Mothers Who Used Tobacco While Pregnant ¹ | 6.2 |
| Percent of Births Less Than 37 Wks. of Gestation | 4.6 |
| ○ Average Age of Mother | 29.0 |
| Teenage Birth Rate ² | LNE |
| Percent White Births | 98.5 |
| Percent American Indian Births | LNE |
| ● Percent Unmarried | 12.3 |
| ● Percent WIC births | 7.7 |
| Percent Breastfeeding at discharge | 89.2 |
| ○ Percent Payment-Private Insurance | 78.1 |
| ● Percent Payment-Medicaid | 12.5 |
| Percent C-Section | 30.8 |

Mortality³

| | |
|-------------------------------------|-------|
| All Causes | 578.9 |
| Heart Disease | 144.3 |
| ● Malignant Neoplasms (Cancer) | 72.4 |
| Trachea, Bronchus, & Lung | LNE |
| Colon, Rectum, & Anus | LNE |
| Pancreas | LNE |
| Female Breast | LNE |
| Leukemia | LNE |
| Prostate | LNE |
| Accidents | 102.3 |
| Motor Vehicle Accidents | LNE |
| Alzheimer's Disease | LNE |
| Cerebrovascular Disease | 54.1 |
| Chronic Lower Respiratory Diseases | 55.7 |
| Diabetes Mellitus | LNE |
| Influenza & Pneumonia | LNE |
| Intentional Self-Harm (Suicide) | LNE |
| Chronic Liver Disease and Cirrhosis | LNE |
| Infant Mortality | LNE |

Leading Causes of Death

| Leading Causes of Death | Total Deaths |
|--------------------------------------|--------------|
| 1. Heart Disease | 11 |
| T2. Malignant Neoplasms (Cancer) | 6 |
| T2. Accidents | 6 |
| Percent of Deaths due to tobacco use | 16.3 |
| Median age at death | 77 |

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Hughes County

Demographic Information



Hughes County is located in the center of the state and averages 23.0 persons per square mile.

2013 Population Information

| Subject | Number | Percent |
|--|--------|---------|
| Total population | 17,508 | 100.0 |
| White | 14,900 | 85.1 |
| Black or African American | 226 | 1.3 |
| American Indian & Alaska Native | 1,831 | 10.5 |
| Asian | 112 | 0.6 |
| Native Hawaiian & Other Pacific Islander | 3 | 0.0 |
| Two or More Races | 436 | 2.5 |
| Under 5 years | 1,228 | 7.0 |
| Under 18 years | 4,133 | 23.6 |
| 65 years and over | 2,541 | 14.5 |

Source: United States Census Bureau, 2013 Population Estimates

Health Status Indicators 2009-2013

Nativity

| | |
|---|------|
| Percent of Low Birth Weight Infants | 5.1 |
| Percent of Mothers Receiving Care in 1st Trimester | 66.4 |
| Percent of Mothers Who Used Tobacco While Pregnant ¹ | 18.8 |
| Percent of Births Less Than 37 Wks. of Gestation | 7.4 |
| Average Age of Mother | 27.4 |
| ● Teenage Birth Rate ² | 8.9 |
| Percent White Births | 74.4 |
| Percent American Indian Births | 19.3 |
| Percent Unmarried | 41.0 |
| Percent WIC births | 36.6 |
| Percent Breastfeeding at discharge | 73.9 |
| Percent Payment-Private Insurance | 57.7 |
| Percent Payment-Medicaid | 36.5 |
| ○ Percent C-Section | 35.8 |

●Denotes a health status indicator which is significantly lower than the state average.
○Denotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

| | |
|-------------------------------------|-------|
| All Causes | 672.7 |
| Heart Disease | 150.0 |
| Malignant Neoplasms (Cancer) | 161.0 |
| ● Trachea, Bronchus, & Lung | 29.9 |
| Colon, Rectum, & Anus | 22.1 |
| Pancreas | 14.8 |
| Female Breast | 15.7 |
| Leukemia | 6.0 |
| Prostate | 14.0 |
| Accidents | 33.6 |
| Motor Vehicle Accidents | 13.7 |
| Alzheimer's Disease | 37.5 |
| Cerebrovascular Disease | 53.2 |
| Chronic Lower Respiratory Diseases | 47.6 |
| Diabetes Mellitus | 22.5 |
| Influenza & Pneumonia | 13.4 |
| Intentional Self-Harm (Suicide) | 18.4 |
| Chronic Liver Disease and Cirrhosis | 10.9 |
| Infant Mortality | 7.52 |

Leading Causes of Death

| Leading Causes of Death | Total Deaths |
|--|--------------|
| 1. Heart Disease | 165 |
| 2. Malignant Neoplasms (Cancer) | 163 |
| 3. Cerebrovascular Diseases | 58 |
| 4. Chronic Lower Respiratory Diseases | 48 |
| 5. Alzheimer's Disease | 41 |
| 6. Accidents | 34 |
| 7. Diabetes | 24 |
| T8. Influenza and Pneumonia | 15 |
| T8. Intentional Self-Harm (Suicide) | 15 |
| T10. Chronic Liver Disease and Cirrhosis | 11 |
| T10. Essential (Primary) Hypertension and Hypertensive Renal Disease | 11 |
| Percent of Deaths due to tobacco use | 18.4 |
| Median age at death | 81 |

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Hutchinson County

Demographic Information



Hutchinson County is located in the southeastern region of the state and averages 9.0 persons per square mile.

2013 Population Information

| Subject | Number | Percent |
|--|--------|---------|
| Total population | 7,145 | 100.0 |
| White | 6,926 | 96.9 |
| Black or African American | 52 | 0.7 |
| American Indian & Alaska Native | 87 | 1.2 |
| Asian | 16 | 0.2 |
| Native Hawaiian & Other Pacific Islander | 1 | 0.0 |
| Two or More Races | 63 | 0.9 |
| Under 5 years | 451 | 6.3 |
| Under 18 years | 1,672 | 23.4 |
| 65 years and over | 1,721 | 24.1 |

Source: United States Census Bureau, 2013 Population Estimates

Health Status Indicators 2009-2013

Nativity

| | |
|---|------|
| ● Percent of Low Birth Weight Infants | 2.8 |
| ○ Percent of Mothers Receiving Care in 1st Trimester | 62.0 |
| ● Percent of Mothers Who Used Tobacco While Pregnant ¹ | 9.0 |
| ○ Percent of Births Less Than 37 Wks. of Gestation | 5.9 |
| ○ Average Age of Mother | 28.4 |
| ● Teenage Birth Rate ² | 4.6 |
| Percent White Births | 98.8 |
| Percent American Indian Births | LNE |
| ● Percent Unmarried | 18.3 |
| ● Percent WIC births | 20.6 |
| Percent Breastfeeding at discharge | 83.0 |
| ○ Percent Payment-Private Insurance | 77.8 |
| ● Percent Payment-Medicaid | 17.5 |
| Percent C-Section | 23.2 |

●Denotes a health status indicator which is significantly lower than the state average.
○Denotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

| | |
|-------------------------------------|-------|
| All Causes | 647.4 |
| Heart Disease | 142.3 |
| Malignant Neoplasms (Cancer) | 136.3 |
| Trachea, Bronchus, & Lung | 33.0 |
| Colon, Rectum, & Anus | 11.4 |
| Pancreas | 7.3 |
| Female Breast | 22.1 |
| Leukemia | 6.0 |
| Prostate | 22.4 |
| Accidents | 44.4 |
| Motor Vehicle Accidents | 17.0 |
| Alzheimer's Disease | 29.0 |
| Cerebrovascular Disease | 42.1 |
| Chronic Lower Respiratory Diseases | 36.7 |
| Diabetes Mellitus | 19.2 |
| Influenza & Pneumonia | 21.4 |
| Intentional Self-Harm (Suicide) | 12.5 |
| Chronic Liver Disease and Cirrhosis | LNE |
| Infant Mortality | LNE |

Leading Causes of Death

| Leading Causes of Death | Total Deaths |
|--|--------------|
| 1. Heart Disease | 139 |
| 2. Malignant Neoplasms (Cancer) | 93 |
| 3. Cerebrovascular Diseases | 42 |
| 4. Alzheimer's Disease | 37 |
| 5. Chronic Lower Respiratory Diseases | 26 |
| 6. Accidents | 24 |
| 7. Influenza and Pneumonia | 20 |
| 8. Diabetes | 14 |
| T9. Nephritis, Nephrotic Syndrome, and Nephrosis | 12 |
| T9. Pneumonitis due to Solids and Liquids | 12 |

| | |
|--------------------------------------|------|
| Percent of Deaths due to tobacco use | 10.2 |
| Median age at death | 86 |

●Denotes a health status indicator which is significantly lower than the state average.
○Denotes a health status indicator which is significantly higher than the state average.


³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Hyde County

Demographic Information



Hyde County is located in the central region of the state and averages 1.7 people per square mile.

2013 Population Information

| Subject | Number | Percent |
|--|--------|---------|
| Total population | 1,391 | 100.0 |
| White | 1,224 | 88.0 |
| Black or African American | 5 | 0.4 |
| American Indian & Alaska Native | 132 | 9.5 |
| Asian | 5 | 0.4 |
| Native Hawaiian & Other Pacific Islander | 1 | 0.1 |
| Two or More Races | 24 | 1.7 |
| Under 5 years | 74 | 5.3 |
| Under 18 years | 293 | 21.1 |
| 65 years and over | 332 | 23.9 |

Source: United States Census Bureau, 2013 Population Estimates

Health Status Indicators 2009-2013

Nativity

| | |
|---|------|
| Percent of Low Birth Weight Infants | LNE |
| Percent of Mothers Receiving Care in 1st Trimester | 73.1 |
| Percent of Mothers Who Used Tobacco While Pregnant ¹ | 14.9 |
| Percent of Births Less Than 37 Wks. of Gestation | 13.4 |
| Average Age of Mother | 27.0 |
| Teenage Birth Rate ² | LNE |
| Percent White Births | 86.6 |
| Percent American Indian Births | 7.5 |
| Percent Unmarried | 29.9 |
| Percent WIC births | 40.9 |
| Percent Breastfeeding at discharge | 69.2 |
| Percent Payment-Private Insurance | 67.2 |
| Percent Payment-Medicaid | 29.9 |
| Percent C-Section | 28.4 |

●Denotes a health status indicator which is significantly lower than the state average.
○Denotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

| | |
|-------------------------------------|-------|
| All Causes | 696.4 |
| ○ Heart Disease | 301.4 |
| Malignant Neoplasms (Cancer) | 107.7 |
| Trachea, Bronchus, & Lung | 24.7 |
| Colon, Rectum, & Anus | LNE |
| Pancreas | LNE |
| Female Breast | LNE |
| Leukemia | LNE |
| Prostate | LNE |
| Accidents | LNE |
| Motor Vehicle Accidents | LNE |
| Alzheimer's Disease | LNE |
| Cerebrovascular Disease | 57.3 |
| Chronic Lower Respiratory Diseases | 26.8 |
| Diabetes Mellitus | LNE |
| Influenza & Pneumonia | LNE |
| Intentional Self-Harm (Suicide) | LNE |
| Chronic Liver Disease and Cirrhosis | LNE |
| Infant Mortality | LNE |

Leading Causes of Death

| Leading Causes of Death | Total Deaths |
|--------------------------------------|--------------|
| 1. Heart Disease | 46 |
| 2. Malignant Neoplasms (Cancer) | 13 |
| 3. Cerebrovascular Diseases | 10 |
| Percent of Deaths due to tobacco use | 9.6 |
| Median age at death | 86 |

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Jackson County

Demographic Information



Jackson County is located in western South Dakota and averages 1.6 persons per square mile.

2013 Population Information

| Subject | Number | Percent |
|--|--------|---------|
| Total population | 3,216 | 100.0 |
| White | 1,387 | 43.1 |
| Black or African American | 21 | 0.7 |
| American Indian & Alaska Native | 1,655 | 51.5 |
| Asian | 3 | 0.1 |
| Native Hawaiian & Other Pacific Islander | 0 | 0.0 |
| Two or More Races | 150 | 4.7 |
| Under 5 years | 323 | 10.0 |
| Under 18 years | 1,072 | 33.3 |
| 65 years and over | 428 | 13.3 |

Source: United States Census Bureau, 2013 Population Estimates

Health Status Indicators 2009-2013

Nativity

| | |
|---|------|
| Percent of Low Birth Weight Infants | 8.9 |
| ● Percent of Mothers Receiving Care in 1st Trimester | 44.9 |
| Percent of Mothers Who Used Tobacco While Pregnant ¹ | 20.2 |
| ○ Percent of Births Less Than 37 Wks. of Gestation | 13.8 |
| ● Average Age of Mother | 25.2 |
| ○ Teenage Birth Rate ² | 42.1 |
| Percent White Births | 19.4 |
| Percent American Indian Births | 76.9 |
| ○ Percent Unmarried | 75.6 |
| ○ Percent WIC births | 74.4 |
| ● Percent Breastfeeding at discharge | 53.1 |
| ● Percent Payment-Private Insurance | 14.4 |
| ○ Percent Payment-Medicaid | 62.5 |
| Percent C-Section | 21.6 |

●Denotes a health status indicator which is significantly lower than the state average.
○Denotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

| | |
|---------------------------------------|---------|
| ○ All Causes | 1,044.4 |
| Heart Disease | 171.8 |
| Malignant Neoplasms (Cancer) | 156.5 |
| Trachea, Bronchus, & Lung | 72.3 |
| Colon, Rectum, & Anus | LNE |
| Pancreas | LNE |
| Female Breast | LNE |
| Leukemia | LNE |
| Prostate | LNE |
| ○ Accidents | 148.6 |
| ○ Motor Vehicle Accidents | 74.2 |
| Alzheimer's Disease | 27.1 |
| Cerebrovascular Disease | LNE |
| Chronic Lower Respiratory Diseases | 77.2 |
| Diabetes Mellitus | 45.3 |
| Influenza & Pneumonia | 27.3 |
| Intentional Self-Harm (Suicide) | 40.2 |
| ○ Chronic Liver Disease and Cirrhosis | 80.2 |
| Infant Mortality | 16.62 |

Leading Causes of Death

| Leading Causes of Death | Total Deaths |
|--|--------------|
| 1. Heart Disease | 27 |
| 2. Malignant Neoplasms (Cancer) | 25 |
| 3. Accidents | 22 |
| 4. Chronic Lower Respiratory Diseases | 13 |
| 5. Chronic Liver Disease and Cirrhosis | 11 |
| 6. Diabetes | 7 |
| 7. Intentional Self-Harm (Suicide) | 6 |
| T8. Alzheimer's Disease | 5 |
| T8. Influenza and Pneumonia | 5 |
| Percent of Deaths due to tobacco use | 14.4 |
| Median age at death | 71 |

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Jerauld County

Demographic Information



Jerauld County is located in the central region of the state and averages 3.9 persons per square mile.

2013 Population Information

| Subject | Number | Percent |
|--|--------|---------|
| Total population | 2,066 | 100.0 |
| White | 2,027 | 98.1 |
| Black or African American | 4 | 0.2 |
| American Indian & Alaska Native | 8 | 0.4 |
| Asian | 4 | 0.2 |
| Native Hawaiian & Other Pacific Islander | 3 | 0.1 |
| Two or More Races | 20 | 1.0 |
| Under 5 years | 141 | 6.8 |
| Under 18 years | 474 | 22.9 |
| 65 years and over | 502 | 24.3 |

Source: United States Census Bureau, 2013 Population Estimates

Health Status Indicators 2009-2013

Nativity

| | |
|---|------|
| Percent of Low Birth Weight Infants | 4.7 |
| Percent of Mothers Receiving Care in 1st Trimester | 66.7 |
| Percent of Mothers Who Used Tobacco While Pregnant ¹ | 11.7 |
| Percent of Births Less Than 37 Wks. of Gestation | 7.0 |
| Average Age of Mother | 28.1 |
| Teenage Birth Rate ² | LNE |
| Percent White Births | 99.2 |
| Percent American Indian Births | LNE |
| • Percent Unmarried | 26.4 |
| Percent WIC births | 30.5 |
| Percent Breastfeeding at discharge | 83.7 |
| Percent Payment-Private Insurance | 64.3 |
| • Percent Payment-Medicaid | 25.6 |
| Percent C-Section | 22.5 |

•Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

| | |
|-------------------------------------|-------|
| All Causes | 642.6 |
| Heart Disease | 121.1 |
| Malignant Neoplasms (Cancer) | 169.5 |
| • Trachea, Bronchus, & Lung | 20.4 |
| Colon, Rectum, & Anus | 24.0 |
| Pancreas | LNE |
| Female Breast | 31.2 |
| Leukemia | 14.1 |
| Prostate | LNE |
| Accidents | 56.7 |
| • Motor Vehicle Accidents | 33.7 |
| ○ Alzheimer's Disease | 91.2 |
| Cerebrovascular Disease | 60.7 |
| Chronic Lower Respiratory Diseases | 36.1 |
| Diabetes Mellitus | LNE |
| Influenza & Pneumonia | 13.0 |
| Intentional Self-Harm (Suicide) | LNE |
| Chronic Liver Disease and Cirrhosis | LNE |
| Infant Mortality | LNE |

Leading Causes of Death

| Leading Causes of Death | Total Deaths |
|---------------------------------------|--------------|
| 1. Malignant Neoplasms (Cancer) | 31 |
| 2. Heart Disease | 30 |
| 3. Alzheimer's Disease | 27 |
| 4. Cerebrovascular Diseases | 14 |
| 5. Chronic Lower Respiratory Diseases | 9 |
| 6. Accidents | 7 |
| Percent of Deaths due to tobacco use | 11.2 |
| Median age at death | 85 |

•Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Jones County

Demographic Information



Jones County is located in western South Dakota and averages 1.0 persons per square mile.

2013 Population Information

| Subject | Number | Percent |
|--|--------|---------|
| Total population | 1,001 | 100.0 |
| White | 919 | 91.8 |
| Black or African American | 7 | 0.7 |
| American Indian & Alaska Native | 35 | 3.5 |
| Asian | 6 | 0.6 |
| Native Hawaiian & Other Pacific Islander | 5 | 0.5 |
| Two or More Races | 29 | 2.9 |
| Under 5 years | 64 | 6.4 |
| Under 18 years | 213 | 21.3 |
| 65 years and over | 214 | 21.4 |

Source: United States Census Bureau, 2013 Population Estimates

Health Status Indicators 2009-2013

Nativity

| | |
|---|------|
| Percent of Low Birth Weight Infants | 9.1 |
| Percent of Mothers Receiving Care in 1st Trimester | 81.8 |
| Percent of Mothers Who Used Tobacco While Pregnant ¹ | 16.4 |
| Percent of Births Less Than 37 Wks. of Gestation | 16.4 |
| Average Age of Mother | 26.1 |
| Teenage Birth Rate ² | 25.4 |
| Percent White Births | 94.5 |
| Percent American Indian Births | LNE |
| • Percent Unmarried | 20.0 |
| Percent WIC births | 47.3 |
| Percent Breastfeeding at discharge | 87.3 |
| Percent Payment-Private Insurance | 68.5 |
| Percent Payment-Medicaid | 24.1 |
| Percent C-Section | 32.7 |

•Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

| | |
|-------------------------------------|-------|
| All Causes | 566.9 |
| • Heart Disease | 73.9 |
| Malignant Neoplasms (Cancer) | 110.9 |
| Trachea, Bronchus, & Lung | LNE |
| Colon, Rectum, & Anus | LNE |
| Pancreas | LNE |
| Female Breast | LNE |
| Leukemia | LNE |
| Prostate | LNE |
| Accidents | LNE |
| Motor Vehicle Accidents | LNE |
| Alzheimer's Disease | LNE |
| Cerebrovascular Disease | LNE |
| Chronic Lower Respiratory Diseases | 44.9 |
| Diabetes Mellitus | LNE |
| Influenza & Pneumonia | LNE |
| Intentional Self-Harm (Suicide) | LNE |
| Chronic Liver Disease and Cirrhosis | LNE |
| Infant Mortality | LNE |

Leading Causes of Death

| Leading Causes of Death | Total Deaths |
|---------------------------------|--------------|
| 1. Malignant Neoplasms (Cancer) | 10 |
| 2. Heart Disease | 7 |

| | |
|--------------------------------------|------|
| Percent of Deaths due to tobacco use | 19.1 |
| Median age at death | 80 |

•Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Kingsbury County

Demographic Information



Kingsbury County is located in east central South Dakota and averages 6.2 persons per square mile.

2013 Population Information

| Subject | Number | Percent |
|--|--------|---------|
| Total population | 5,065 | 100.0 |
| White | 4,927 | 97.3 |
| Black or African American | 21 | 0.4 |
| American Indian & Alaska Native | 34 | 0.7 |
| Asian | 22 | 0.4 |
| Native Hawaiian & Other Pacific Islander | 0 | 0.0 |
| Two or More Races | 61 | 1.2 |
| Under 5 years | 336 | 6.6 |
| Under 18 years | 1,121 | 22.1 |
| 65 years and over | 1,096 | 21.6 |

Source: United States Census Bureau, 2013 Population Estimates

Health Status Indicators 2009-2013

Nativity

| | |
|---|------|
| Percent of Low Birth Weight Infants | 5.3 |
| Percent of Mothers Receiving Care in 1st Trimester | 70.3 |
| Percent of Mothers Who Used Tobacco While Pregnant ¹ | 15.1 |
| Percent of Births Less Than 37 Wks. of Gestation | 6.7 |
| ○ Average Age of Mother | 28.5 |
| Teenage Birth Rate ² | LNE |
| Percent White Births | 98.0 |
| Percent American Indian Births | 1.0 |
| ● Percent Unmarried | 19.0 |
| ● Percent WIC births | 23.7 |
| Percent Breastfeeding at discharge | 79.9 |
| ○ Percent Payment-Private Insurance | 79.9 |
| ● Percent Payment-Medicaid | 17.4 |
| Percent C-Section | 22.7 |

●Denotes a health status indicator which is significantly lower than the state average.
 ○Denotes a health status indicator which is significantly higher than the state average.
¹Data for mothers who used tobacco are self-reported.
²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

| | |
|--------------------------------------|-------|
| All Causes | 737.8 |
| Heart Disease | 168.7 |
| Malignant Neoplasms (Cancer) | 164.9 |
| Trachea, Bronchus, & Lung | 44.8 |
| Colon, Rectum, & Anus | 17.6 |
| Pancreas | 15.0 |
| Female Breast | 15.4 |
| Leukemia | 19.9 |
| Prostate | LNE |
| Accidents | 58.6 |
| Motor Vehicle Accidents | 38.4 |
| Alzheimer's Disease | 37.5 |
| Cerebrovascular Disease | 43.1 |
| ● Chronic Lower Respiratory Diseases | 25.9 |
| Diabetes Mellitus | 19.8 |
| Influenza & Pneumonia | 23.5 |
| Intentional Self-Harm (Suicide) | 24.1 |
| Chronic Liver Disease and Cirrhosis | LNE |
| Infant Mortality | 16.67 |


Leading Causes of Death

| Leading Causes of Death | Total Deaths |
|---|--------------|
| 1. Heart Disease | 95 |
| 2. Malignant Neoplasms (Cancer) | 76 |
| 3. Cerebrovascular Diseases | 25 |
| 4. Alzheimer's Disease | 22 |
| 5. Accidents | 17 |
| 6. Chronic Lower Respiratory Diseases | 14 |
| 7. Influenza and Pneumonia | 13 |
| T8. Diabetes | 8 |
| T8. Unspecified Dementia | 8 |
| T10. Intentional Self-Harm (Suicide) | 6 |
| T10. Nephritis, Nephrotic Syndrome, and Nephrosis | 6 |
| T10. Septicemia | 6 |
| Percent of Deaths due to tobacco use | 12.7 |
| Median age at death | 83 |

●Denotes a health status indicator which is significantly lower than the state average.
 ○Denotes a health status indicator which is significantly higher than the state average.
³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.
 See technical notes for more information.
 Source: South Dakota Department of Health, Office of Health Statistics

Lake County

Demographic Information



Lake County is located in the east central region of the state and averages 19.9 persons per square mile.

2013 Population Information

| Subject | Number | Percent |
|--|--------|---------|
| Total population | 12,055 | 100.0 |
| White | 11,537 | 95.7 |
| Black or African American | 128 | 1.1 |
| American Indian & Alaska Native | 113 | 0.9 |
| Asian | 104 | 0.9 |
| Native Hawaiian & Other Pacific Islander | 4 | 0.0 |
| Two or More Races | 169 | 1.4 |
| Under 5 years | 659 | 5.5 |
| Under 18 years | 2,417 | 20.0 |
| 65 years and over | 2,302 | 19.1 |

Source: United States Census Bureau, 2013 Population Estimates

Health Status Indicators 2009-2013

Nativity

| | |
|---|------|
| Percent of Low Birth Weight Infants | 5.2 |
| Percent of Mothers Receiving Care in 1st Trimester | 73.3 |
| ● Percent of Mothers Who Used Tobacco While Pregnant ¹ | 12.9 |
| Percent of Births Less Than 37 Wks. of Gestation | 7.1 |
| ○ Average Age of Mother | 28.2 |
| ● Teenage Birth Rate ² | 9.0 |
| Percent White Births | 96.8 |
| Percent American Indian Births | 1.5 |
| ● Percent Unmarried | 23.2 |
| ● Percent WIC births | 26.4 |
| Percent Breastfeeding at discharge | 78.7 |
| ○ Percent Payment-Private Insurance | 74.2 |
| ● Percent Payment-Medicaid | 20.5 |
| Percent C-Section | 26.9 |

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

| | |
|-------------------------------------|-------|
| ● All Causes | 630.0 |
| Heart Disease | 141.8 |
| Malignant Neoplasms (Cancer) | 151.7 |
| Trachea, Bronchus, & Lung | 35.3 |
| Colon, Rectum, & Anus | 11.1 |
| Pancreas | 7.9 |
| Female Breast | 25.6 |
| Leukemia | 4.0 |
| Prostate | 10.6 |
| ● Accidents | 28.1 |
| Motor Vehicle Accidents | 11.6 |
| ○ Alzheimer's Disease | 56.0 |
| Cerebrovascular Disease | 47.5 |
| Chronic Lower Respiratory Diseases | 45.2 |
| Diabetes Mellitus | 16.6 |
| Influenza & Pneumonia | 8.9 |
| Intentional Self-Harm (Suicide) | 13.1 |
| Chronic Liver Disease and Cirrhosis | 10.2 |
| Infant Mortality | 7.59 |

Leading Causes of Death

| Leading Causes of Death | Total Deaths |
|--|--------------|
| 1. Malignant Neoplasms (Cancer) | 129 |
| 2. Heart Disease | 126 |
| 3. Alzheimer's Disease | 53 |
| 4. Cerebrovascular Diseases | 41 |
| 5. Chronic Lower Respiratory Diseases | 39 |
| 6. Accidents | 21 |
| 7. Diabetes | 14 |
| 8. Essential (Primary) Hypertension and Hypertensive Renal Disease | 9 |
| T9. Influenza and Pneumonia | 8 |
| T9. Aortic Aneurysm and Dissection | 8 |
| Percent of Deaths due to tobacco use | 21.6 |
| Median age at death | 82 |

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Lawrence County

Demographic Information



Lawrence County is located along the Wyoming border and averages 30.1 persons per square mile.

2013 Population Information

| Subject | Number | Percent |
|--|--------|---------|
| Total population | 24,910 | 100.0 |
| White | 23,328 | 93.6 |
| Black or African American | 321 | 1.3 |
| American Indian & Alaska Native | 569 | 2.3 |
| Asian | 189 | 0.8 |
| Native Hawaiian & Other Pacific Islander | 17 | 0.1 |
| Two or More Races | 486 | 2.0 |
| Under 5 years | 1,174 | 4.7 |
| Under 18 years | 4,658 | 18.7 |
| 65 years and over | 4,369 | 17.5 |

Source: United States Census Bureau, 2013 Population Estimates

Health Status Indicators 2009-2013

Nativity

| | |
|---|------|
| ○ Percent of Low Birth Weight Infants | 8.6 |
| ○ Percent of Mothers Receiving Care in 1st Trimester | 80.4 |
| ○ Percent of Mothers Who Used Tobacco While Pregnant ¹ | 19.9 |
| Percent of Births Less Than 37 Wks. of Gestation | 8.5 |
| Average Age of Mother | 27.4 |
| Teenage Birth Rate ² | 14.8 |
| Percent White Births | 93.7 |
| Percent American Indian Births | 1.9 |
| Percent Unmarried | 37.5 |
| Percent WIC births | 38.5 |
| ○ Percent Breastfeeding at discharge | 80.9 |
| Percent Payment-Private Insurance | 55.7 |
| Percent Payment-Medicaid | 36.7 |
| ● Percent C-Section | 21.1 |

Mortality³

| | |
|-------------------------------------|-------|
| ● All Causes | 636.5 |
| Heart Disease | 141.5 |
| Malignant Neoplasms (Cancer) | 165.0 |
| Trachea, Bronchus, & Lung | 38.3 |
| Colon, Rectum, & Anus | 11.7 |
| Pancreas | 16.4 |
| Female Breast | 21.8 |
| Leukemia | 5.4 |
| Prostate | 29.2 |
| ● Accidents | 29.1 |
| ● Motor Vehicle Accidents | 8.0 |
| Alzheimer's Disease | 27.9 |
| Cerebrovascular Disease | 36.8 |
| Chronic Lower Respiratory Diseases | 53.7 |
| Diabetes Mellitus | 24.4 |
| ● Influenza & Pneumonia | 8.7 |
| Intentional Self-Harm (Suicide) | 17.5 |
| Chronic Liver Disease and Cirrhosis | 6.8 |
| Infant Mortality | 6.71 |

Leading Causes of Death

| Leading Causes of Death | Total Deaths |
|---------------------------------------|--------------|
| 1. Malignant Neoplasms (Cancer) | 274 |
| 2. Heart Disease | 248 |
| 3. Chronic Lower Respiratory Diseases | 94 |
| 4. Cerebrovascular Diseases | 66 |
| 5. Alzheimer's Disease | 54 |
| 6. Accidents | 44 |
| 7. Diabetes | 42 |
| 8. Intentional Self-Harm (Suicide) | 21 |
| 9. Unspecified Dementia | 18 |
| 10. Influenza and Pneumonia | 17 |

| | |
|--------------------------------------|------|
| Percent of Deaths due to tobacco use | 20.5 |
| Median age at death | 81 |

● Denotes a health status indicator which is significantly lower than the state average.

○ Denotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

● Denotes a health status indicator which is significantly lower than the state average.

○ Denotes a health status indicator which is significantly higher than the state average.


³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Lincoln County

Demographic Information



Lincoln County is located in southeastern South Dakota and averages 77.7 persons per square mile.

2013 Population Information

| Subject | Number | Percent |
|--|--------|---------|
| Total population | 49,858 | 100.0 |
| White | 47,598 | 95.5 |
| Black or African American | 603 | 1.2 |
| American Indian & Alaska Native | 319 | 0.6 |
| Asian | 602 | 1.2 |
| Native Hawaiian & Other Pacific Islander | 15 | 0.0 |
| Two or More Races | 721 | 1.4 |
| Under 5 years | 4,306 | 8.6 |
| Under 18 years | 14,677 | 29.4 |
| 65 years and over | 4,907 | 9.8 |

Source: United States Census Bureau, 2013 Population Estimates

Health Status Indicators 2009-2013

Nativity

| | |
|---|------|
| Percent of Low Birth Weight Infants | 5.7 |
| ○ Percent of Mothers Receiving Care in 1st Trimester | 80.9 |
| ● Percent of Mothers Who Used Tobacco While Pregnant ¹ | 7.0 |
| Percent of Births Less Than 37 Wks. of Gestation | 7.5 |
| ○ Average Age of Mother | 29.0 |
| ● Teenage Birth Rate ² | 6.4 |
| Percent White Births | 97.6 |
| Percent American Indian Births | 0.4 |
| ● Percent Unmarried | 16.0 |
| ● Percent WIC births | 10.8 |
| ○ Percent Breastfeeding at discharge | 78.5 |
| ○ Percent Payment-Private Insurance | 86.6 |
| ● Percent Payment-Medicaid | 11.4 |
| Percent C-Section | 26.9 |

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

| | |
|---------------------------------------|-------|
| ● All Causes | 507.8 |
| ● Heart Disease | 110.7 |
| ● Malignant Neoplasms (Cancer) | 122.6 |
| Trachea, Bronchus, & Lung | 33.9 |
| Colon, Rectum, & Anus | 13.0 |
| Pancreas | 7.9 |
| Female Breast | 21.4 |
| Leukemia | 5.8 |
| Prostate | 19.6 |
| ● Accidents | 25.8 |
| ● Motor Vehicle Accidents | 6.7 |
| Alzheimer's Disease | 34.1 |
| Cerebrovascular Disease | 32.2 |
| ● Chronic Lower Respiratory Diseases | 31.5 |
| ● Diabetes Mellitus | 13.8 |
| ● Influenza & Pneumonia | 8.3 |
| ● Intentional Self-Harm (Suicide) | 9.6 |
| ● Chronic Liver Disease and Cirrhosis | 2.4 |
| Infant Mortality | 5.47 |

Leading Causes of Death

| Leading Causes of Death | Total Deaths |
|---------------------------------------|--------------|
| 1. Malignant Neoplasms (Cancer) | 222 |
| 2. Heart Disease | 206 |
| 3. Alzheimer's Disease | 63 |
| 4. Cerebrovascular Diseases | 56 |
| 5. Chronic Lower Respiratory Diseases | 54 |
| 6. Accidents | 53 |
| 7. Diabetes | 24 |
| 8. Intentional Self-Harm (Suicide) | 21 |
| 9. Unspecified Dementia | 20 |
| 10. Parkinson's Disease | 17 |
| Percent of Deaths due to tobacco use | 15.3 |
| Median age at death | 81 |

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Lyman County

Demographic Information



Lyman County is located in central South Dakota and averages 2.3 persons per square mile.

2013 Population Information

| Subject | Number | Percent |
|--|--------|---------|
| Total population | 3,892 | 100.0 |
| White | 2,261 | 58.1 |
| Black or African American | 19 | 0.5 |
| American Indian & Alaska Native | 1,486 | 38.2 |
| Asian | 14 | 0.4 |
| Native Hawaiian & Other Pacific Islander | 1 | 0.0 |
| Two or More Races | 111 | 2.9 |
| Under 5 years | 308 | 7.9 |
| Under 18 years | 1,114 | 28.6 |
| 65 years and over | 585 | 15.0 |

Source: United States Census Bureau, 2013 Population Estimates

Health Status Indicators 2009-2013

Nativity

| | |
|---|------|
| Percent of Low Birth Weight Infants | 4.8 |
| ● Percent of Mothers Receiving Care in 1st Trimester | 49.7 |
| ○ Percent of Mothers Who Used Tobacco While Pregnant ¹ | 26.8 |
| Percent of Births Less Than 37 Wks. of Gestation | 6.7 |
| ● Average Age of Mother | 26.3 |
| ○ Teenage Birth Rate ² | 32.1 |
| Percent White Births | 36.7 |
| Percent American Indian Births | 60.3 |
| ○ Percent Unmarried | 61.5 |
| ○ Percent WIC births | 64.3 |
| ● Percent Breastfeeding at discharge | 62.5 |
| ● Percent Payment-Private Insurance | 35.4 |
| ○ Percent Payment-Medicaid | 59.8 |
| ○ Percent C-Section | 33.9 |

●Denotes a health status indicator which is significantly lower than the state average.
○Denotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

| | |
|-------------------------------------|-------|
| ○ All Causes | 858.8 |
| Heart Disease | 210.3 |
| Malignant Neoplasms (Cancer) | 135.7 |
| Trachea, Bronchus, & Lung | 59.1 |
| Colon, Rectum, & Anus | 22.0 |
| Pancreas | LNE |
| Female Breast | 33.2 |
| Leukemia | LNE |
| Prostate | LNE |
| ○ Accidents | 148.9 |
| ○ Motor Vehicle Accidents | 87.6 |
| Alzheimer's Disease | 59.1 |
| Cerebrovascular Disease | 30.1 |
| Chronic Lower Respiratory Diseases | 55.3 |
| Diabetes Mellitus | 28.7 |
| Influenza & Pneumonia | 20.1 |
| Intentional Self-Harm (Suicide) | LNE |
| Chronic Liver Disease and Cirrhosis | 13.8 |
| Infant Mortality | LNE |

Leading Causes of Death

| Leading Causes of Death | Total Deaths |
|--|--------------|
| 1. Heart Disease | 42 |
| 2. Malignant Neoplasms (Cancer) | 31 |
| 3. Accidents | 27 |
| T4. Chronic Lower Respiratory Diseases | 11 |
| T4. Alzheimer's Disease | 11 |
| 6. Diabetes | 7 |
| 7. Cerebrovascular Diseases | 6 |
| Percent of Deaths due to tobacco use | 25.3 |
| Median age at death | 74 |

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

McCook County

Demographic Information



McCook County is located in eastern South Dakota and averages 9.8 persons per square mile.

2013 Population Information

| Subject | Number | Percent |
|--|--------|---------|
| Total population | 5,654 | 100.0 |
| White | 5,528 | 97.8 |
| Black or African American | 29 | 0.5 |
| American Indian & Alaska Native | 32 | 0.6 |
| Asian | 18 | 0.3 |
| Native Hawaiian & Other Pacific Islander | 2 | 0.0 |
| Two or More Races | 45 | 0.8 |
| Under 5 years | 388 | 6.9 |
| Under 18 years | 1,462 | 25.9 |
| 65 years and over | 1,081 | 19.1 |

Source: United States Census Bureau, 2013 Population Estimates

Health Status Indicators 2009-2013

Nativity

| | |
|---|------|
| Percent of Low Birth Weight Infants | 6.9 |
| Percent of Mothers Receiving Care in 1st Trimester | 69.0 |
| ● Percent of Mothers Who Used Tobacco While Pregnant ¹ | 11.1 |
| Percent of Births Less Than 37 Wks. of Gestation | 8.7 |
| ○ Average Age of Mother | 28.4 |
| Teenage Birth Rate ² | 9.8 |
| Percent White Births | 98.4 |
| Percent American Indian Births | 0.8 |
| ● Percent Unmarried | 21.6 |
| ● Percent WIC births | 30.6 |
| Percent Breastfeeding at discharge | 76.3 |
| ○ Percent Payment-Private Insurance | 75.7 |
| ● Percent Payment-Medicaid | 21.1 |
| Percent C-Section | 24.3 |

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

| | |
|-------------------------------------|-------|
| ○ All Causes | 851.7 |
| ○ Heart Disease | 227.6 |
| Malignant Neoplasms (Cancer) | 203.3 |
| Trachea, Bronchus, & Lung | 58.7 |
| Colon, Rectum, & Anus | 19.1 |
| Pancreas | 10.0 |
| Female Breast | LNE |
| Leukemia | 10.8 |
| Prostate | 29.9 |
| Accidents | 48.9 |
| Motor Vehicle Accidents | 27.5 |
| Alzheimer's Disease | 58.3 |
| Cerebrovascular Disease | 56.5 |
| Chronic Lower Respiratory Diseases | 37.8 |
| Diabetes Mellitus | 22.7 |
| Influenza & Pneumonia | 14.4 |
| Intentional Self-Harm (Suicide) | 10.6 |
| Chronic Liver Disease and Cirrhosis | LNE |
| Infant Mortality | 7.92 |

Leading Causes of Death

| Leading Causes of Death | Total Deaths |
|--|--------------|
| 1. Heart Disease | 104 |
| 2. Malignant Neoplasms (Cancer) | 87 |
| 3. Alzheimer's Disease | 31 |
| 4. Cerebrovascular Diseases | 28 |
| 5. Chronic Lower Respiratory Diseases | 18 |
| 6. Accidents | 12 |
| 7. Diabetes | 10 |
| T8. Influenza and Pneumonia | 8 |
| T8. Septicemia | 8 |
| 10. Nephritis, Nephrotic Syndrome, and Nephrosis | 6 |

| | |
|--------------------------------------|------|
| Percent of Deaths due to tobacco use | 14.3 |
| Median age at death | 81 |

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

McPherson County

Demographic Information



McPherson County is located in the north central region of the state and averages 2.2 persons per square mile.

2013 Population Information

| Subject | Number | Percent |
|--|--------|---------|
| Total population | 2,457 | 100.0 |
| White | 2,411 | 98.1 |
| Black or African American | 9 | 0.4 |
| American Indian & Alaska Native | 6 | 0.2 |
| Asian | 11 | 0.4 |
| Native Hawaiian & Other Pacific Islander | 2 | 0.1 |
| Two or More Races | 18 | 0.7 |
| Under 5 years | 133 | 5.4 |
| Under 18 years | 531 | 21.6 |
| 65 years and over | 709 | 28.9 |

Source: United States Census Bureau, 2013 Population Estimates

Health Status Indicators 2009-2013

Nativity

| | |
|---|------|
| Percent of Low Birth Weight Infants | 7.3 |
| Percent of Mothers Receiving Care in 1st Trimester | 62.3 |
| Percent of Mothers Who Used Tobacco While Pregnant ¹ | 14.7 |
| Percent of Births Less Than 37 Wks. of Gestation | 7.3 |
| ○ Average Age of Mother | 29.9 |
| Teenage Birth Rate ² | LNE |
| Percent White Births | 98.1 |
| Percent American Indian Births | LNE |
| ● Percent Unmarried | 15.6 |
| ● Percent WIC births | 19.6 |
| Percent Breastfeeding at discharge | 84.3 |
| ○ Percent Payment-Private Insurance | 75.9 |
| ● Percent Payment-Medicaid | 17.6 |
| Percent C-Section | 19.3 |

●Denotes a health status indicator which is significantly lower than the state average.
○Denotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

| | |
|-------------------------------------|-------|
| ● All Causes | 578.9 |
| Heart Disease | 177.2 |
| ● Malignant Neoplasms (Cancer) | 92.2 |
| Trachea, Bronchus, & Lung | 34.0 |
| Colon, Rectum, & Anus | 14.0 |
| Pancreas | LNE |
| Female Breast | LNE |
| Leukemia | LNE |
| Prostate | 25.4 |
| Accidents | 49.1 |
| Motor Vehicle Accidents | 20.3 |
| ● Alzheimer's Disease | 18.8 |
| Cerebrovascular Disease | 63.0 |
| Chronic Lower Respiratory Diseases | 33.7 |
| Diabetes Mellitus | 20.9 |
| Influenza & Pneumonia | 10.7 |
| Intentional Self-Harm (Suicide) | LNE |
| Chronic Liver Disease and Cirrhosis | LNE |
| Infant Mortality | LNE |

Leading Causes of Death

| Leading Causes of Death | Total Deaths |
|--|--------------|
| 1. Heart Disease | 52 |
| 2. Malignant Neoplasms (Cancer) | 25 |
| 3. Cerebrovascular Diseases | 20 |
| 4. Chronic Lower Respiratory Diseases | 12 |
| 5. Essential (Primary) Hypertension and Hypertensive Renal Disease | 11 |
| 6. Accidents | 9 |
| 7. Alzheimer's Disease | 8 |
| 8. Diabetes | 7 |

| | |
|--------------------------------------|------|
| Percent of Deaths due to tobacco use | 10.6 |
| Median age at death | 84 |

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Marshall County

Demographic Information



Marshall County is located in the northeastern part of the state and averages 5.6 persons per square mile.

2013 Population Information

| Subject | Number | Percent |
|--|--------|---------|
| Total population | 4,763 | 100.0 |
| White | 4,144 | 87.0 |
| Black or African American | 69 | 1.4 |
| American Indian & Alaska Native | 467 | 9.8 |
| Asian | 14 | 0.3 |
| Native Hawaiian & Other Pacific Islander | 0 | 0.0 |
| Two or More Races | 69 | 1.4 |
| Under 5 years | 284 | 6.0 |
| Under 18 years | 1,046 | 22.0 |
| 65 years and over | 927 | 19.5 |

Source: United States Census Bureau, 2013 Population Estimates

Health Status Indicators 2009-2013

Nativity

| | |
|---|------|
| Percent of Low Birth Weight Infants | 5.2 |
| Percent of Mothers Receiving Care in 1st Trimester | 61.4 |
| Percent of Mothers Who Used Tobacco While Pregnant ¹ | 17.5 |
| Percent of Births Less Than 37 Wks. of Gestation | 5.6 |
| Average Age of Mother | 27.5 |
| Teenage Birth Rate ² | 12.7 |
| Percent White Births | 81.3 |
| Percent American Indian Births | 15.8 |
| ● Percent Unmarried | 30.3 |
| ● Percent WIC births | 30.8 |
| Percent Breastfeeding at discharge | 72.8 |
| Percent Payment-Private Insurance | 59.8 |
| Percent Payment-Medicaid | 29.0 |
| Percent C-Section | 28.9 |

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

| | |
|-------------------------------------|-------|
| ○ All Causes | 823.2 |
| Heart Disease | 164.9 |
| Malignant Neoplasms (Cancer) | 197.5 |
| Trachea, Bronchus, & Lung | 52.2 |
| Colon, Rectum, & Anus | 20.9 |
| Pancreas | 25.3 |
| Female Breast | 25.6 |
| Leukemia | 7.9 |
| Prostate | 25.0 |
| Accidents | 55.6 |
| Motor Vehicle Accidents | 22.1 |
| Alzheimer's Disease | 39.6 |
| ○ Cerebrovascular Disease | 68.6 |
| Chronic Lower Respiratory Diseases | 63.9 |
| Diabetes Mellitus | 28.3 |
| Influenza & Pneumonia | 21.4 |
| Intentional Self-Harm (suicide) | 12.6 |
| Chronic Liver Disease and Cirrhosis | 8.1 |
| Infant Mortality | 20.91 |

Leading Causes of Death

| Leading Causes of Death | Total Deaths |
|--|--------------|
| 1. Malignant Neoplasms (Cancer) | 70 |
| 2. Heart Disease | 64 |
| 3. Cerebrovascular Diseases | 29 |
| 4. Chronic Lower Respiratory Diseases | 27 |
| 5. Alzheimer's Disease | 18 |
| 6. Accidents | 12 |
| 7. Influenza and Pneumonia | 9 |
| 8. Diabetes | 8 |
| 9. Other Interstitial Pulmonary Diseases with Fibrosis | 5 |
| Percent of Deaths due to tobacco use | 18.0 |
| Median age at death | 82 |

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Meade County

Demographic Information



Meade County is located in west central South Dakota and averages 7.3 persons per square mile.

2013 Population Information

| Subject | Number | Percent |
|--|--------|---------|
| Total population | 27,202 | 100.0 |
| White | 24,842 | 91.3 |
| Black or African American | 551 | 2.0 |
| American Indian & Alaska Native | 790 | 2.9 |
| Asian | 240 | 0.9 |
| Native Hawaiian & Other Pacific Islander | 27 | 0.1 |
| Two or More Races | 752 | 2.8 |
| Under 5 years | 1,759 | 6.5 |
| Under 18 years | 6,541 | 24.0 |
| 65 years and over | 3,508 | 12.9 |

Source: United States Census Bureau, 2013 Population Estimates

Health Status Indicators 2009-2013

Nativity

| | |
|---|------|
| Percent of Low Birth Weight Infants | 6.8 |
| ○ Percent of Mothers Receiving Care in 1st Trimester | 78.1 |
| Percent of Mothers Who Used Tobacco While Pregnant ¹ | 15.0 |
| Percent of Births Less Than 37 Wks. of Gestation | 7.7 |
| Average Age of Mother | 27.3 |
| ● Teenage Birth Rate ² | 11.0 |
| Percent White Births | 93.1 |
| Percent American Indian Births | 2.6 |
| ● Percent Unmarried | 26.3 |
| ● Percent WIC births | 33.9 |
| ○ Percent Breastfeeding at discharge | 84.4 |
| Percent Payment-Private Insurance | 51.7 |
| ● Percent Payment-Medicaid | 28.0 |
| ● Percent C-Section | 21.0 |

● Denotes a health status indicator which is significantly lower than the state average.
 ○ Denotes a health status indicator which is significantly higher than the state average.
¹Data for mothers who used tobacco are self-reported.
²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

| | |
|-------------------------------------|-------|
| All Causes | 686.6 |
| Heart Disease | 153.8 |
| Malignant Neoplasms (Cancer) | 149.9 |
| Trachea, Bronchus, & Lung | 44.8 |
| Colon, Rectum, & Anus | 12.1 |
| Pancreas | 13.5 |
| Female Breast | 9.2 |
| Leukemia | 6.5 |
| Prostate | 15.6 |
| Accidents | 34.6 |
| Motor Vehicle Accidents | 12.3 |
| ● Alzheimer's Disease | 20.4 |
| Cerebrovascular Disease | 46.4 |
| Chronic Lower Respiratory Diseases | 55.1 |
| Diabetes Mellitus | 20.7 |
| Influenza & Pneumonia | 12.0 |
| ○ Intentional Self-Harm (suicide) | 30.8 |
| Chronic Liver Disease and Cirrhosis | 13.9 |
| Infant Mortality | 6.22 |

Leading Causes of Death


| Leading Causes of Death | Total Deaths |
|--|--------------|
| 1. Malignant Neoplasms (Cancer) | 200 |
| 2. Heart Disease | 195 |
| 3. Chronic Lower Respiratory Diseases | 69 |
| 4. Cerebrovascular Diseases | 58 |
| 5. Accidents | 42 |
| 6. Intentional Self-Harm (Suicide) | 36 |
| 7. Diabetes | 26 |
| 8. Alzheimer's Disease | 24 |
| 9. Chronic Liver Disease and Cirrhosis | 19 |
| 10. Influenza and Pneumonia | 15 |
| Percent of Deaths due to tobacco use | 20.7 |
| Median age at death | 76 |

● Denotes a health status indicator which is significantly lower than the state average.
 ○ Denotes a health status indicator which is significantly higher than the state average.
³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.
 Source: South Dakota Department of Health, Office of Health Statistics

Mellette County

Demographic Information



Mellette County is located in the south central region of the state and averages 1.6 persons per square mile.

2013 Population Information

| Subject | Number | Percent |
|--|--------|---------|
| Total population | 2,081 | 100.0 |
| White | 858 | 41.2 |
| Black or African American | 8 | 0.4 |
| American Indian & Alaska Native | 1,099 | 52.8 |
| Asian | 5 | 0.2 |
| Native Hawaiian & Other Pacific Islander | 0 | 0.0 |
| Two or More Races | 111 | 5.3 |
| Under 5 years | 184 | 8.8 |
| Under 18 years | 629 | 30.2 |
| 65 years and over | 317 | 15.2 |

Source: United States Census Bureau, 2013 Population Estimates

Health Status Indicators 2009-2013

Nativity

| | |
|---|------|
| Percent of Low Birth Weight Infants | 5.5 |
| ● Percent of Mothers Receiving Care in 1st Trimester | 47.1 |
| Percent of Mothers Who Used Tobacco While Pregnant ¹ | 22.5 |
| Percent of Births Less Than 37 Wks. of Gestation | 5.5 |
| ● Average Age of Mother | 24.3 |
| ○ Teenage Birth Rate ² | 49.5 |
| Percent White Births | 18.6 |
| Percent American Indian Births | 68.3 |
| ○ Percent Unmarried | 71.3 |
| ○ Percent WIC births | 66.3 |
| Percent Breastfeeding at discharge | 72.6 |
| ● Percent Payment-Private Insurance | 12.8 |
| ○ Percent Payment-Medicaid | 61.0 |
| Percent C-Section | 34.1 |

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

| | |
|---------------------------------------|-------|
| ○ All Causes | 996.2 |
| Heart Disease | 137.6 |
| Malignant Neoplasms (Cancer) | 226.1 |
| Trachea, Bronchus, & Lung | 65.8 |
| Colon, Rectum, & Anus | 39.0 |
| Pancreas | LNE |
| Female Breast | LNE |
| Leukemia | LNE |
| Prostate | LNE |
| Accidents | 82.6 |
| Motor Vehicle Accidents | 63.2 |
| Alzheimer's Disease | LNE |
| Cerebrovascular Disease | 80.0 |
| Chronic Lower Respiratory Diseases | 70.0 |
| Diabetes Mellitus | 63.4 |
| Influenza & Pneumonia | LNE |
| Intentional Self-Harm (suicide) | 31.3 |
| ○ Chronic Liver Disease and Cirrhosis | 70.8 |
| Infant Mortality | LNE |

Leading Causes of Death

| Leading Causes of Death | Total Deaths |
|---|--------------|
| 1. Malignant Neoplasms (Cancer) | 27 |
| 2. Heart Disease | 17 |
| 3. Cerebrovascular Diseases | 10 |
| T4. Chronic Lower Respiratory Diseases | 8 |
| T4. Accidents | 8 |
| T6. Diabetes | 7 |
| T6. Chronic Liver Disease and Cirrhosis | 7 |
| Percent of Deaths due to tobacco use | 20.0 |
| Median age at death | 74 |

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Miner County

Demographic Information



Miner County is located in the east central region of the state and averages 4.2 persons per square mile.

2013 Population Information

| Subject | Number | Percent |
|--|--------|---------|
| Total population | 2,333 | 100.0 |
| White | 2,276 | 97.6 |
| Black or African American | 12 | 0.5 |
| American Indian & Alaska Native | 10 | 0.4 |
| Asian | 10 | 0.4 |
| Native Hawaiian & Other Pacific Islander | 0 | 0.0 |
| Two or More Races | 25 | 1.1 |
| Under 5 years | 131 | 5.6 |
| Under 18 years | 558 | 23.9 |
| 65 years and over | 502 | 21.5 |

Source: United States Census Bureau, 2013 Population Estimates

Health Status Indicators 2009-2013

Nativity

| | |
|---|------|
| Percent of Low Birth Weight Infants | LNE |
| Percent of Mothers Receiving Care in 1st Trimester | 70.9 |
| Percent of Mothers Who Used Tobacco While Pregnant ¹ | 11.8 |
| ● Percent of Births Less Than 37 Wks. of Gestation | 2.5 |
| ○ Average Age of Mother | 28.9 |
| Teenage Birth Rate ² | LNE |
| Percent White Births | 96.6 |
| Percent American Indian Births | LNE |
| ● Percent Unmarried | 14.3 |
| ● Percent WIC births | 21.8 |
| Percent Breastfeeding at discharge | 79.7 |
| ○ Percent Payment-Private Insurance | 78.0 |
| ● Percent Payment-Medicaid | 17.8 |
| Percent C-Section | 29.4 |

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

| | |
|-------------------------------------|-------|
| All Causes | 702.1 |
| Heart Disease | 163.9 |
| Malignant Neoplasms (Cancer) | 176.1 |
| Trachea, Bronchus, & Lung | 53.7 |
| Colon, Rectum, & Anus | 20.1 |
| Pancreas | 16.0 |
| Female Breast | LNE |
| Leukemia | LNE |
| Prostate | 32.1 |
| Accidents | 53.3 |
| Motor Vehicle Accidents | 36.3 |
| ○ Alzheimer's Disease | 77.8 |
| Cerebrovascular Disease | 43.5 |
| Chronic Lower Respiratory Diseases | 45.7 |
| Diabetes Mellitus | LNE |
| Influenza & Pneumonia | 16.7 |
| Intentional Self-Harm (Suicide) | LNE |
| Chronic Liver Disease and Cirrhosis | LNE |
| Infant Mortality | LNE |

| Leading Causes of Death | Total Deaths |
|---------------------------------------|--------------|
| 1. Heart Disease | 43 |
| 2. Malignant Neoplasms (Cancer) | 41 |
| 3. Alzheimer's Disease | 24 |
| 4. Cerebrovascular Diseases | 14 |
| 5. Chronic Lower Respiratory Diseases | 10 |
| 6. Accidents | 6 |
| Percent of Deaths due to tobacco use | 18.9 |
| Median age at death | 85 |

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Minnehaha County

Demographic Information



Minnehaha County is located in southeastern South Dakota and averages 210.0 persons per square mile.

2013 Population Information

| Subject | Number | Percent |
|--|---------|---------|
| Total population | 179,640 | 100.0 |
| White | 159,249 | 88.6 |
| Black or African American | 8,195 | 4.6 |
| American Indian & Alaska Native | 4,857 | 2.7 |
| Asian | 3,073 | 1.7 |
| Native Hawaiian & Other Pacific Islander | 202 | 0.1 |
| Two or More Races | 4,064 | 2.3 |
| Under 5 years | 13,751 | 7.7 |
| Under 18 years | 44,835 | 25.0 |
| 65 years and over | 21,305 | 11.9 |

Source: United States Census Bureau, 2013 Population Estimates

Health Status Indicators 2009-2013

Nativity

| | |
|---|------|
| ○ Percent of Low Birth Weight Infants | 7.0 |
| Percent of Mothers Receiving Care in 1st Trimester | 71.5 |
| ● Percent of Mothers Who Used Tobacco While Pregnant ¹ | 14.5 |
| Percent of Births Less Than 37 Wks. of Gestation | 8.2 |
| ○ Average Age of Mother | 27.7 |
| Teenage Birth Rate ² | 15.4 |
| Percent White Births | 83.1 |
| Percent American Indian Births | 4.9 |
| ● Percent Unmarried | 35.7 |
| ● Percent WIC births | 30.6 |
| Percent Breastfeeding at discharge | 74.1 |
| ○ Percent Payment-Private Insurance | 63.7 |
| ● Percent Payment-Medicaid | 32.7 |
| Percent C-Section | 26.6 |

●Denotes a health status indicator which is significantly lower than the state average.
○Denotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

| | |
|-------------------------------------|-------|
| All Causes | 704.2 |
| Heart Disease | 145.6 |
| ○ Malignant Neoplasms (Cancer) | 178.8 |
| ○ Trachea, Bronchus, & Lung | 50.6 |
| Colon, Rectum, & Anus | 16.0 |
| Pancreas | 10.6 |
| Female Breast | 22.2 |
| Leukemia | 8.4 |
| Prostate | 21.4 |
| ● Accidents | 36.1 |
| ● Motor Vehicle Accidents | 9.9 |
| Alzheimer's Disease | 40.9 |
| Cerebrovascular Disease | 39.3 |
| Chronic Lower Respiratory Diseases | 48.5 |
| ● Diabetes Mellitus | 17.5 |
| Influenza & Pneumonia | 12.5 |
| Intentional Self-Harm (Suicide) | 15.3 |
| Chronic Liver Disease and Cirrhosis | 9.6 |
| ● Infant Mortality | 4.80 |

Leading Causes of Death

| Leading Causes of Death | Total Deaths |
|---|--------------|
| 1. Malignant Neoplasms (Cancer) | 1,560 |
| 2. Heart Disease | 1,322 |
| 3. Chronic Lower Respiratory Diseases | 424 |
| 4. Alzheimer's Disease | 381 |
| 5. Cerebrovascular Diseases | 348 |
| 6. Accidents | 325 |
| 7. Diabetes | 155 |
| 8. Intentional Self-Harm (Suicide) | 134 |
| 9. Influenza and Pneumonia | 116 |
| 10. Chronic Liver Disease and Cirrhosis | 87 |
| Percent of Deaths due to tobacco use | 18.2 |
| Median age at death | 79 |

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Moody County

Demographic Information



Moody County is located on the Minnesota border and averages 12.5 persons per square mile.

2013 Population Information

| Subject | Number | Percent |
|--|--------|---------|
| Total population | 6,404 | 100.0 |
| White | 5,221 | 81.5 |
| Black or African American | 65 | 1.0 |
| American Indian & Alaska Native | 867 | 13.5 |
| Asian | 75 | 1.2 |
| Native Hawaiian & Other Pacific Islander | 0 | 0.0 |
| Two or More Races | 176 | 2.7 |
| Under 5 years | 458 | 7.2 |
| Under 18 years | 1,622 | 25.3 |
| 65 years and over | 1,055 | 16.5 |

Source: United States Census Bureau, 2013 Population Estimates

Health Status Indicators 2009-2013

Nativity

| | |
|---|------|
| Percent of Low Birth Weight Infants | 6.2 |
| Percent of Mothers Receiving Care in 1st Trimester | 62.9 |
| Percent of Mothers Who Used Tobacco While Pregnant ¹ | 21.1 |
| Percent of Births Less Than 37 Wks. of Gestation | 10.4 |
| Average Age of Mother | 27.5 |
| Teenage Birth Rate ² | 14.7 |
| Percent White Births | 77.8 |
| Percent American Indian Births | 17.0 |
| Percent Unmarried | 36.6 |
| Percent WIC births | 39.8 |
| Percent Breastfeeding at discharge | 73.6 |
| Percent Payment-Private Insurance | 62.0 |
| Percent Payment-Medicaid | 32.3 |
| Percent C-Section | 23.3 |

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

| | |
|--------------------------------------|-------|
| All Causes | 633.4 |
| Heart Disease | 163.7 |
| Malignant Neoplasms (Cancer) | 173.1 |
| Trachea, Bronchus, & Lung | 32.2 |
| Colon, Rectum, & Anus | 24.3 |
| Pancreas | 12.1 |
| Female Breast | 17.2 |
| Leukemia | LNE |
| Prostate | 26.3 |
| Accidents | 42.9 |
| Motor Vehicle Accidents | 15.8 |
| ○ Alzheimer's Disease | 66.1 |
| Cerebrovascular Disease | 36.8 |
| ● Chronic Lower Respiratory Diseases | 24.9 |
| Diabetes Mellitus | 34.1 |
| Influenza & Pneumonia | LNE |
| Intentional Self-Harm (Suicide) | 15.8 |
| Chronic Liver Disease and Cirrhosis | LNE |
| Infant Mortality | LNE |

Leading Causes of Death

| Leading Causes of Death | Total Deaths |
|---------------------------------------|--------------|
| 1. Heart Disease | 71 |
| 2. Malignant Neoplasms (Cancer) | 70 |
| 3. Alzheimer's Disease | 32 |
| 4. Accidents | 16 |
| T5. Cerebrovascular Diseases | 14 |
| T5. Diabetes | 14 |
| 7. Chronic Lower Respiratory Diseases | 10 |
| 8. Intentional Self-Harm (Suicide) | 6 |
| Percent of Deaths due to tobacco use | 21.8 |
| Median age at death | 81 |

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Pennington County

Demographic Information



Pennington County is located on the Wyoming border and averages 36.4 persons per square mile.

2013 Population Information

| Subject | Number | Percent |
|--|---------|---------|
| Total population | 105,761 | 100.0 |
| White | 88,275 | 83.5 |
| Black or African American | 1,798 | 1.7 |
| American Indian & Alaska Native | 10,471 | 9.9 |
| Asian | 1,279 | 1.2 |
| Native Hawaiian & Other Pacific Islander | 117 | 0.1 |
| Two or More Races | 3,821 | 3.6 |
| Under 5 years | 7,424 | 7.0 |
| Under 18 years | 25,435 | 24.0 |
| 65 years and over | 15,787 | 14.9 |

Source: United States Census Bureau, 2013 Population Estimates

Health Status Indicators 2009-2013

Nativity

| | |
|---|------|
| Percent of Low Birth Weight Infants | 6.8 |
| Percent of Mothers Receiving Care in 1st Trimester | 72.3 |
| ○ Percent of Mothers Who Used Tobacco While Pregnant ¹ | 18.5 |
| Percent of Births Less Than 37 Wks. of Gestation | 8.5 |
| ● Average Age of Mother | 26.8 |
| Teenage Birth Rate ² | 16.1 |
| Percent White Births | 71.8 |
| Percent American Indian Births | 18.9 |
| ○ Percent Unmarried | 43.1 |
| ○ Percent WIC births | 45.1 |
| ○ Percent Breastfeeding at discharge | 80.2 |
| ● Percent Payment-Private Insurance | 38.6 |
| ○ Percent Payment-Medicaid | 41.8 |
| ● Percent C-Section | 20.2 |

● Denotes a health status indicator which is significantly lower than the state average.

○ Denotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

| | |
|-------------------------------------|-------|
| ● All Causes | 640.4 |
| Heart Disease | 143.0 |
| Malignant Neoplasms (Cancer) | 157.5 |
| Trachea, Bronchus, & Lung | 45.4 |
| ● Colon, Rectum, & Anus | 11.2 |
| Pancreas | 11.8 |
| Female Breast | 21.5 |
| Leukemia | 8.1 |
| Prostate | 17.1 |
| Accidents | 44.2 |
| ● Motor Vehicle Accidents | 9.0 |
| ● Alzheimer's Disease | 28.6 |
| ● Cerebrovascular Disease | 29.5 |
| Chronic Lower Respiratory Diseases | 39.6 |
| ● Diabetes Mellitus | 11.1 |
| Influenza & Pneumonia | 15.4 |
| Intentional Self-Harm (Suicide) | 17.1 |
| Chronic Liver Disease and Cirrhosis | 10.6 |
| Infant Mortality | 7.49 |

Leading Causes of Death

| Leading Causes of Death | Total Deaths |
|---------------------------------------|--------------|
| 1. Malignant Neoplasms (Cancer) | 931 |
| 2. Heart Disease | 866 |
| 3. Accidents | 239 |
| 4. Chronic Lower Respiratory Diseases | 237 |
| 5. Alzheimer's Disease | 179 |
| 6. Cerebrovascular Diseases | 177 |
| 7. Unspecified Dementia | 121 |
| 8. Influenza and Pneumonia | 94 |
| 9. Intentional Self-Harm (Suicide) | 89 |
| 10. Diabetes | 66 |
| Percent of Deaths due to tobacco use | 16.9 |
| Median age at death | 77 |

● Denotes a health status indicator which is significantly lower than the state average.

○ Denotes a health status indicator which is significantly higher than the state average.


³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Perkins County

Demographic Information



Perkins County is located in northwestern South Dakota and averages 1.0 persons per square mile.

2013 Population Information

| Subject | Number | Percent |
|--|--------|---------|
| Total population | 3,037 | 100.0 |
| White | 2,943 | 96.9 |
| Black or African American | 10 | 0.3 |
| American Indian & Alaska Native | 47 | 1.5 |
| Asian | 8 | 0.3 |
| Native Hawaiian & Other Pacific Islander | 0 | 0.0 |
| Two or More Races | 29 | 1.0 |
| Under 5 years | 158 | 5.2 |
| Under 18 years | 648 | 21.3 |
| 65 years and over | 713 | 23.5 |

Source: United States Census Bureau, 2013 Population Estimates

Health Status Indicators 2009-2013

Natality

| | |
|---|------|
| Percent of Low Birth Weight Infants | 6.3 |
| Percent of Mothers Receiving Care in 1st Trimester | 61.1 |
| Percent of Mothers Who Used Tobacco While Pregnant ¹ | 13.2 |
| Percent of Births Less Than 37 Wks. of Gestation | 6.3 |
| ○ Average Age of Mother | 28.9 |
| Teenage Birth Rate ² | LNE |
| Percent White Births | 97.5 |
| Percent American Indian Births | 2.5 |
| ● Percent Unmarried | 19.5 |
| Percent WIC births | 38.2 |
| Percent Breastfeeding at discharge | 80.3 |
| Percent Payment-Private Insurance | 60.5 |
| ● Percent Payment-Medicaid | 22.4 |
| Percent C-Section | 32.1 |

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

| | |
|--------------------------------------|-------|
| All Causes | 759.4 |
| Heart Disease | 124.5 |
| Malignant Neoplasms (Cancer) | 189.2 |
| Trachea, Bronchus, & Lung | 32.4 |
| Colon, Rectum, & Anus | 23.7 |
| Pancreas | LNE |
| Female Breast | 51.5 |
| Leukemia | LNE |
| Prostate | 31.0 |
| Accidents | 88.2 |
| Motor Vehicle Accidents | 30.6 |
| Alzheimer's Disease | 30.4 |
| Cerebrovascular Disease | 32.1 |
| ● Chronic Lower Respiratory Diseases | 25.5 |
| Diabetes Mellitus | 24.9 |
| Influenza & Pneumonia | 13.6 |
| Intentional Self-Harm (Suicide) | 24.4 |
| Chronic Liver Disease and Cirrhosis | LNE |
| Infant Mortality | LNE |

Leading Causes of Death

| Leading Causes of Death | Total Deaths |
|--|--------------|
| 1. Malignant Neoplasms (Cancer) | 52 |
| 2. Heart Disease | 43 |
| 3. Accidents | 15 |
| 4. Cerebrovascular Diseases | 13 |
| 5. Alzheimer's Disease | 11 |
| T6. Chronic Lower Respiratory Diseases | 9 |
| T6. Diabetes | 9 |
| 8. Unspecified Dementia | 6 |
| 9. Septicemia | 5 |

| | |
|--------------------------------------|------|
| Percent of Deaths due to tobacco use | 15.0 |
| Median age at death | 83 |

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Potter County

Demographic Information



Potter County is located in north central South Dakota and averages 2.7 persons per square mile.

2013 Population Information

| Subject | Number | Percent |
|--|--------|---------|
| Total population | 2,394 | 100.0 |
| White | 2,315 | 96.7 |
| Black or African American | 9 | 0.4 |
| American Indian & Alaska Native | 37 | 1.5 |
| Asian | 9 | 0.4 |
| Native Hawaiian & Other Pacific Islander | 0 | 0.0 |
| Two or More Races | 24 | 1.0 |
| Under 5 years | 133 | 5.6 |
| Under 18 years | 502 | 21.0 |
| 65 years and over | 631 | 26.4 |

Source: United States Census Bureau, 2013 Population Estimates

Health Status Indicators 2009-2013

Nativity

| | |
|---|------|
| Percent of Low Birth Weight Infants | 4.3 |
| Percent of Mothers Receiving Care in 1st Trimester | 65.8 |
| Percent of Mothers Who Used Tobacco While Pregnant ¹ | 14.5 |
| Percent of Births Less Than 37 Wks. of Gestation | 6.8 |
| Average Age of Mother | 27.7 |
| Teenage Birth Rate ² | LNE |
| Percent White Births | 94.0 |
| Percent American Indian Births | 4.3 |
| ● Percent Unmarried | 24.8 |
| ● Percent WIC births | 23.3 |
| Percent Breastfeeding at discharge | 77.6 |
| ○ Percent Payment-Private Insurance | 76.9 |
| ● Percent Payment-Medicaid | 20.5 |
| Percent C-Section | 25.6 |

●Denotes a health status indicator which is significantly lower than the state average.
○Denotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

| | |
|-------------------------------------|-------|
| All Causes | 640.3 |
| Heart Disease | 134.6 |
| Malignant Neoplasms (Cancer) | 153.6 |
| Trachea, Bronchus, & Lung | 58.3 |
| Colon, Rectum, & Anus | 17.9 |
| Pancreas | LNE |
| Female Breast | LNE |
| Leukemia | LNE |
| Prostate | LNE |
| Accidents | 37.9 |
| Motor Vehicle Accidents | LNE |
| Alzheimer's Disease | 27.7 |
| Cerebrovascular Disease | 21.3 |
| Chronic Lower Respiratory Diseases | 37.7 |
| Diabetes Mellitus | 24.4 |
| Influenza & Pneumonia | 19.5 |
| Intentional Self-Harm (Suicide) | LNE |
| Chronic Liver Disease and Cirrhosis | LNE |
| Infant Mortality | LNE |

Leading Causes of Death

| Leading Causes of Death | Total Deaths |
|---------------------------------------|--------------|
| 1. Heart Disease | 42 |
| 2. Malignant Neoplasms (Cancer) | 40 |
| 3. Chronic Lower Respiratory Diseases | 12 |
| 4. Alzheimer's Disease | 10 |
| 5. Influenza and Pneumonia | 8 |
| T6. Cerebrovascular Diseases | 7 |
| T6. Unspecified Dementia | 7 |
| 8. Diabetes | 6 |
| T9. Accidents | 5 |
| T9. Parkinson's Disease | 5 |
| Percent of Deaths due to tobacco use | 16.9 |
| Median age at death | 85 |

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Roberts County

Demographic Information



Roberts County is located in the extreme northeastern corner of the state and averages 9.2 persons per square mile.

2013 Population Information

| Subject | Number | Percent |
|--|--------|---------|
| Total population | 10,251 | 100.0 |
| White | 6,201 | 60.5 |
| Black or African American | 26 | 0.3 |
| American Indian & Alaska Native | 3,668 | 35.8 |
| Asian | 34 | 0.3 |
| Native Hawaiian & Other Pacific Islander | 1 | 0.0 |
| Two or More Races | 321 | 3.1 |
| Under 5 years | 862 | 8.4 |
| Under 18 years | 2,883 | 28.1 |
| 65 years and over | 1,859 | 18.1 |

Source: United States Census Bureau, 2013 Population Estimates

Health Status Indicators 2009-2013

Nativity

| | |
|---|------|
| Percent of Low Birth Weight Infants | 5.7 |
| ● Percent of Mothers Receiving Care in 1st Trimester | 53.1 |
| ○ Percent of Mothers Who Used Tobacco While Pregnant ¹ | 27.7 |
| Percent of Births Less Than 37 Wks. of Gestation | 8.8 |
| ● Average Age of Mother | 25.3 |
| ○ Teenage Birth Rate ² | 33.8 |
| Percent White Births | 34.6 |
| Percent American Indian Births | 57.3 |
| ○ Percent Unmarried | 63.1 |
| ○ Percent WIC births | 58.9 |
| ● Percent Breastfeeding at discharge | 68.0 |
| ● Percent Payment-Private Insurance | 32.2 |
| ○ Percent Payment-Medicaid | 60.3 |
| Percent C-Section | 23.1 |

●Denotes a health status indicator which is significantly lower than the state average.
○Denotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

| | |
|-------------------------------------|-------|
| ○ All Causes | 831.3 |
| ○ Heart Disease | 192.1 |
| Malignant Neoplasms (Cancer) | 191.0 |
| Trachea, Bronchus, & Lung | 37.1 |
| Colon, Rectum, & Anus | 15.9 |
| Pancreas | 8.5 |
| Female Breast | 24.8 |
| Leukemia | 10.0 |
| Prostate | 19.6 |
| ○ Accidents | 76.4 |
| ○ Motor Vehicle Accidents | 41.1 |
| Alzheimer's Disease | 28.5 |
| Cerebrovascular Disease | 49.8 |
| Chronic Lower Respiratory Diseases | 32.5 |
| Diabetes Mellitus | 30.9 |
| Influenza & Pneumonia | 11.8 |
| Intentional Self-Harm (Suicide) | 25.7 |
| Chronic Liver Disease and Cirrhosis | 27.8 |
| Infant Mortality | 6.73 |

Leading Causes of Death

| Leading Causes of Death | Total Deaths |
|--|--------------|
| 1. Heart Disease | 144 |
| 2. Malignant Neoplasms (Cancer) | 129 |
| 3. Accidents | 40 |
| 4. Cerebrovascular Diseases | 37 |
| 5. Chronic Lower Respiratory Diseases | 24 |
| 6. Alzheimer's Disease | 23 |
| 7. Diabetes | 19 |
| 8. Chronic Liver Disease and Cirrhosis | 13 |
| 9. Intentional Self-Harm (Suicide) | 11 |
| 10. Septicemia | 9 |

| | |
|--------------------------------------|------|
| Percent of Deaths due to tobacco use | 17.0 |
| Median age at death | 79 |

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Sanborn County

Demographic Information



Sanborn County is located in east central South Dakota and averages 4.1 persons per square mile.

2013 Population Information

| Subject | Number | Percent |
|--|--------|---------|
| Total population | 2,324 | 100.0 |
| White | 2,267 | 97.5 |
| Black or African American | 6 | 0.3 |
| American Indian & Alaska Native | 8 | 0.3 |
| Asian | 6 | 0.3 |
| Native Hawaiian & Other Pacific Islander | 2 | 0.1 |
| Two or More Races | 35 | 1.5 |
| Under 5 years | 123 | 5.3 |
| Under 18 years | 489 | 21.0 |
| 65 years and over | 449 | 19.3 |

Source: United States Census Bureau, 2013 Population Estimates

Health Status Indicators 2009-2013

Nativity

| | |
|---|------|
| Percent of Low Birth Weight Infants | 5.5 |
| Percent of Mothers Receiving Care in 1st Trimester | 66.7 |
| Percent of Mothers Who Used Tobacco While Pregnant ¹ | 15.9 |
| Percent of Births Less Than 37 Wks. of Gestation | 5.5 |
| ○ Average Age of Mother | 28.5 |
| Teenage Birth Rate ² | LNE |
| Percent White Births | 99.3 |
| Percent American Indian Births | LNE |
| ● Percent Unmarried | 25.5 |
| Percent WIC births | 29.4 |
| Percent Breastfeeding at discharge | 80.7 |
| ○ Percent Payment-Private Insurance | 74.5 |
| ● Percent Payment-Medicaid | 20.7 |
| Percent C-Section | 25.5 |

●Denotes a health status indicator which is significantly lower than the state average.
○Denotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

| | |
|-------------------------------------|-------|
| All Causes | 708.2 |
| Heart Disease | 137.0 |
| Malignant Neoplasms (Cancer) | 159.3 |
| Trachea, Bronchus, & Lung | 42.1 |
| Colon, Rectum, & Anus | 23.9 |
| Pancreas | LNE |
| Female Breast | LNE |
| Leukemia | LNE |
| Prostate | LNE |
| Accidents | 61.6 |
| Motor Vehicle Accidents | 31.9 |
| ○ Alzheimer's Disease | 94.1 |
| Cerebrovascular Disease | 35.8 |
| Chronic Lower Respiratory Diseases | 28.2 |
| Diabetes Mellitus | 19.6 |
| Influenza & Pneumonia | 16.5 |
| Intentional Self-Harm (Suicide) | LNE |
| Chronic Liver Disease and Cirrhosis | LNE |
| Infant Mortality | LNE |

Leading Causes of Death

| Leading Causes of Death | Total Deaths |
|---------------------------------------|--------------|
| T1. Heart Disease | 28 |
| T1. Malignant Neoplasms (Cancer) | 28 |
| 3. Alzheimer's Disease | 23 |
| 4. Accidents | 10 |
| 5. Cerebrovascular Diseases | 8 |
| 6. Chronic Lower Respiratory Diseases | 6 |
| Percent of Deaths due to tobacco use | 15.0 |
| Median age at death | 83 |

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Shannon County

Demographic Information



Shannon County is located in the southwestern part of the state, along the Nebraska border and averages 6.5 persons per square mile.

2013 Population Information

| Subject | Number | Percent |
|--|--------|---------|
| Total population | 14,118 | 100.0 |
| White | 747 | 5.3 |
| Black or African American | 28 | 0.2 |
| American Indian & Alaska Native | 13,114 | 92.9 |
| Asian | 19 | 0.1 |
| Native Hawaiian & Other Pacific Islander | 3 | 0.0 |
| Two or More Races | 207 | 1.5 |
| Under 5 years | 1,631 | 11.6 |
| Under 18 years | 5,392 | 38.2 |
| 65 years and over | 946 | 6.7 |

Source: United States Census Bureau, 2013 Population Estimates

Health Status Indicators 2009-2013

Natality

| | |
|---|------|
| Percent of Low Birth Weight Infants | 6.6 |
| ● Percent of Mothers Receiving Care in 1st Trimester | 52.0 |
| ○ Percent of Mothers Who Used Tobacco While Pregnant ¹ | 21.7 |
| ○ Percent of Births Less Than 37 Wks. of Gestation | 11.2 |
| ● Average Age of Mother | 24.2 |
| ○ Teenage Birth Rate ² | 57.5 |
| Percent White Births | 2.1 |
| Percent American Indian Births | 97.2 |
| ○ Percent Unmarried | 87.3 |
| ○ Percent WIC births | 75.0 |
| ● Percent Breastfeeding at discharge | 52.0 |
| ● Percent Payment-Private Insurance | 3.7 |
| ○ Percent Payment-Medicaid | 65.9 |
| ● Percent C-Section | 22.7 |

●Denotes a health status indicator which is significantly lower than the state average.
○Denotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

| | |
|---------------------------------------|--------|
| ○ All Causes | 1338.2 |
| Heart Disease | 174.4 |
| ○ Malignant Neoplasms (Cancer) | 219.3 |
| Trachea, Bronchus, & Lung | 53.8 |
| Colon, Rectum, & Anus | 18.1 |
| Pancreas | 17.1 |
| Female Breast | LNE |
| Leukemia | 12.4 |
| Prostate | 29.0 |
| ○ Accidents | 143.2 |
| ○ Motor Vehicle Accidents | 65.8 |
| ● Alzheimer's Disease | 13.8 |
| Cerebrovascular Disease | 53.5 |
| Chronic Lower Respiratory Diseases | 69.2 |
| ○ Diabetes Mellitus | 93.4 |
| ○ Influenza & Pneumonia | 54.3 |
| ○ Intentional Self-Harm (Suicide) | 39.6 |
| ○ Chronic Liver Disease and Cirrhosis | 99.2 |
| ○ Infant Mortality | 13.64 |

Leading Causes of Death

| Leading Causes of Death | Total Deaths |
|--|--------------|
| 1. Accidents | 89 |
| 2. Malignant Neoplasms (Cancer) | 80 |
| 3. Heart Disease | 67 |
| 4. Chronic Liver Disease and Cirrhosis | 52 |
| 5. Diabetes | 39 |
| 6. Intentional Self-Harm (Suicide) | 31 |
| 7. Chronic Lower Respiratory Diseases | 23 |
| 8. Cerebrovascular Diseases | 20 |
| 9. Influenza and Pneumonia | 18 |
| 10. Septicemia | 17 |
| Percent of Deaths due to tobacco use | 11.3 |
| Median age at death | 54 |

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Spink County

Demographic Information



Spink County is located in the center of eastern South Dakota and averages 4.3 persons per square mile.

2013 Population Information

| Subject | Number | Percent |
|--|--------|---------|
| Total population | 6,610 | 100.0 |
| White | 6,392 | 96.7 |
| Black or African American | 46 | 0.7 |
| American Indian & Alaska Native | 101 | 1.5 |
| Asian | 8 | 0.1 |
| Native Hawaiian & Other Pacific Islander | 0 | 0.0 |
| Two or More Races | 63 | 1.0 |
| Under 5 years | 385 | 5.8 |
| Under 18 years | 1,607 | 24.3 |
| 65 years and over | 1,294 | 19.6 |

Source: United States Census Bureau, 2013 Population Estimates

Health Status Indicators 2009-2013

Nativity

| | |
|---|------|
| Percent of Low Birth Weight Infants | 7.6 |
| Percent of Mothers Receiving Care in 1st Trimester | 69.5 |
| Percent of Mothers Who Used Tobacco While Pregnant ¹ | 15.8 |
| Percent of Births Less Than 37 Wks. of Gestation | 8.7 |
| Average Age of Mother | 27.6 |
| ● Teenage Birth Rate ² | 5.3 |
| Percent White Births | 96.9 |
| Percent American Indian Births | LNE |
| ● Percent Unmarried | 27.8 |
| ● Percent WIC births | 32.1 |
| Percent Breastfeeding at discharge | 79.6 |
| ○ Percent Payment-Private Insurance | 70.7 |
| ● Percent Payment-Medicaid | 24.8 |
| Percent C-Section | 26.2 |

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

| | |
|-------------------------------------|-------|
| All Causes | 738.1 |
| Heart Disease | 146.9 |
| Malignant Neoplasms (Cancer) | 156.3 |
| Trachea, Bronchus, & Lung | 46.1 |
| Colon, Rectum, & Anus | 18.5 |
| Pancreas | 11.8 |
| Female Breast | 20.2 |
| Leukemia | LNE |
| Prostate | 15.7 |
| ○ Accidents | 75.2 |
| Motor Vehicle Accidents | 24.5 |
| ○ Alzheimer's Disease | 65.2 |
| Cerebrovascular Disease | 30.6 |
| Chronic Lower Respiratory Diseases | 62.5 |
| Diabetes Mellitus | 24.5 |
| Influenza & Pneumonia | 24.3 |
| Intentional Self-Harm (Suicide) | 9.7 |
| Chronic Liver Disease and Cirrhosis | 13.8 |
| Infant Mortality | LNE |

Leading Causes of Death

| Leading Causes of Death | Total Deaths |
|--|--------------|
| 1. Heart Disease | 86 |
| 2. Malignant Neoplasms (Cancer) | 85 |
| 3. Alzheimer's Disease | 45 |
| 4. Chronic Lower Respiratory Diseases | 34 |
| 5. Accidents | 28 |
| 6. Cerebrovascular Diseases | 18 |
| 7. Influenza and Pneumonia | 16 |
| 8. Diabetes | 10 |
| T9. Chronic Liver Disease and Cirrhosis | 6 |
| T9. Disorders of Lipoprotein Metabolism and other Lipidemias | 6 |

| | |
|--------------------------------------|------|
| Percent of Deaths due to tobacco use | 14.4 |
| Median age at death | 83 |

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Stanley County

Demographic Information



Stanley County is located in the center of the state and averages 2.1 persons per square mile.

2013 Population Information

| Subject | Number | Percent |
|--|--------|---------|
| Total population | 2,981 | 100.0 |
| White | 2,631 | 88.3 |
| Black or African American | 23 | 0.8 |
| American Indian & Alaska Native | 251 | 8.4 |
| Asian | 5 | 0.2 |
| Native Hawaiian & Other Pacific Islander | 0 | 0.0 |
| Two or More Races | 71 | 2.4 |
| Under 5 years | 176 | 5.9 |
| Under 18 years | 710 | 23.8 |
| 65 years and over | 502 | 16.8 |

Source: United States Census Bureau, 2013 Population Estimates

Health Status Indicators 2009-2013

Natality

| | |
|---|------|
| Percent of Low Birth Weight Infants | 4.6 |
| Percent of Mothers Receiving Care in 1st Trimester | 65.4 |
| Percent of Mothers Who Used Tobacco While Pregnant ¹ | 20.6 |
| Percent of Births Less Than 37 Wks. of Gestation | 6.7 |
| Average Age of Mother | 27.8 |
| Teenage Birth Rate ² | LNE |
| Percent White Births | 83.5 |
| Percent American Indian Births | 12.4 |
| Percent Unmarried | 41.2 |
| Percent WIC births | 46.4 |
| Percent Breastfeeding at discharge | 74.2 |
| Percent Payment-Private Insurance | 59.8 |
| Percent Payment-Medicaid | 36.1 |
| ○ Percent C-Section | 43.8 |

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

| | |
|-------------------------------------|-------|
| ● All Causes | 511.7 |
| ● Heart Disease | 89.1 |
| Malignant Neoplasms (Cancer) | 164.6 |
| Trachea, Bronchus, & Lung | 49.7 |
| Colon, Rectum, & Anus | 17.7 |
| Pancreas | LNE |
| Female Breast | 36.3 |
| Leukemia | LNE |
| Prostate | LNE |
| Accidents | 18.5 |
| Motor Vehicle Accidents | 18.5 |
| Alzheimer's Disease | LNE |
| Cerebrovascular Disease | 31.6 |
| Chronic Lower Respiratory Diseases | 58.8 |
| Diabetes Mellitus | 15.7 |
| Influenza & Pneumonia | LNE |
| Intentional Self-Harm (Suicide) | 21.7 |
| Chronic Liver Disease and Cirrhosis | LNE |
| Infant Mortality | LNE |

Leading Causes of Death

| Leading Causes of Death | Total Deaths |
|---------------------------------------|--------------|
| 1. Malignant Neoplasms (Cancer) | 29 |
| 2. Heart Disease | 14 |
| 3. Chronic Lower Respiratory Diseases | 11 |
| 4. Cerebrovascular Diseases | 5 |

Percent of Deaths due to tobacco use 33.3

Median age at death 70

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Sully County

Demographic Information



Sully County is located in the central region of the state and averages 1.4 persons per square mile.

2013 Population Information

| Subject | Number | Percent |
|--|--------|---------|
| Total population | 1,437 | 100.0 |
| White | 1,368 | 95.2 |
| Black or African American | 18 | 1.3 |
| American Indian & Alaska Native | 24 | 1.7 |
| Asian | 0 | 0.0 |
| Native Hawaiian & Other Pacific Islander | 0 | 0.0 |
| Two or More Races | 27 | 1.9 |
| Under 5 years | 71 | 4.9 |
| Under 18 years | 320 | 22.3 |
| 65 years and over | 265 | 18.4 |

Source: United States Census Bureau, 2013 Population Estimates

Health Status Indicators 2009-2013

Nativity

| | |
|---|------|
| Percent of Low Birth Weight Infants | 10.6 |
| Percent of Mothers Receiving Care in 1st Trimester | 60.7 |
| Percent of Mothers Who Used Tobacco While Pregnant ¹ | 10.7 |
| Percent of Births Less Than 37 Wks. of Gestation | 10.6 |
| Average Age of Mother | 27.9 |
| Teenage Birth Rate ² | LNE |
| Percent White Births | 87.1 |
| Percent American Indian Births | 10.6 |
| • Percent Unmarried | 25.9 |
| Percent WIC births | 29.4 |
| Percent Breastfeeding at discharge | 76.2 |
| Percent Payment-Private Insurance | 65.9 |
| Percent Payment-Medicaid | 29.4 |
| Percent C-Section | 28.2 |

•Denotes a health status indicator which is significantly lower than the state average.
 ○Denotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

| | |
|-------------------------------------|-------|
| • All Causes | 539.9 |
| Heart Disease | 115.7 |
| Malignant Neoplasms (Cancer) | 186.7 |
| Trachea, Bronchus, & Lung | 48.9 |
| Colon, Rectum, & Anus | LNE |
| Pancreas | 31.1 |
| Female Breast | LNE |
| Leukemia | LNE |
| Prostate | LNE |
| Accidents | 29.2 |
| Motor Vehicle Accidents | LNE |
| Alzheimer's Disease | LNE |
| Cerebrovascular Disease | LNE |
| Chronic Lower Respiratory Diseases | LNE |
| Diabetes Mellitus | 27.4 |
| Influenza & Pneumonia | LNE |
| Intentional Self-Harm (Suicide) | LNE |
| Chronic Liver Disease and Cirrhosis | LNE |
| Infant Mortality | LNE |

| Leading Causes of Death | Total Deaths |
|---------------------------------|--------------|
| 1. Malignant Neoplasms (Cancer) | 19 |
| 2. Heart Disease | 12 |

| | |
|--------------------------------------|------|
| Percent of Deaths due to tobacco use | 30.2 |
| Median age at death | 73 |

•Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Todd County

Demographic Information



Todd County is located in south central South Dakota, along the Nebraska border and averages 6.9 persons per square mile.

2013 Population Information

| Subject | Number | Percent |
|--|--------|---------|
| Total population | 9,982 | 100.0 |
| White | 1,079 | 10.8 |
| Black or African American | 24 | 0.2 |
| American Indian & Alaska Native | 8,631 | 86.5 |
| Asian | 27 | 0.3 |
| Native Hawaiian & Other Pacific Islander | 0 | 0.0 |
| Two or More Races | 221 | 2.2 |
| Under 5 years | 1,301 | 13.0 |
| Under 18 years | 4,023 | 40.3 |
| 65 years and over | 678 | 6.8 |

Source: United States Census Bureau, 2013 Population Estimates

Health Status Indicators 2009-2013

Nativity

| | |
|---|------|
| Percent of Low Birth Weight Infants | 7.4 |
| ● Percent of Mothers Receiving Care in 1st Trimester | 38.9 |
| ○ Percent of Mothers Who Used Tobacco While Pregnant ¹ | 25.5 |
| Percent of Births Less Than 37 Wks. of Gestation | 9.7 |
| ● Average Age of Mother | 24.6 |
| ○ Teenage Birth Rate ² | 63.0 |
| Percent White Births | 4.4 |
| Percent American Indian Births | 92.7 |
| ○ Percent Unmarried | 83.0 |
| ○ Percent WIC births | 74.4 |
| ● Percent Breastfeeding at discharge | 59.5 |
| ● Percent Payment-Private Insurance | 4.3 |
| ○ Percent Payment-Medicaid | 44.7 |
| ○ Percent C-Section | 33.9 |

●Denotes a health status indicator which is significantly lower than the state average.
○Denotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

| | |
|---------------------------------------|--------|
| ○ All Causes | 1304.2 |
| Heart Disease | 201.8 |
| ○ Malignant Neoplasms (Cancer) | 230.0 |
| Trachea, Bronchus, & Lung | 63.7 |
| Colon, Rectum, & Anus | 37.2 |
| Pancreas | 10.8 |
| Female Breast | LNE |
| Leukemia | 14.9 |
| Prostate | 31.1 |
| ○ Accidents | 158.9 |
| ○ Motor Vehicle Accidents | 90.6 |
| Alzheimer's Disease | 18.1 |
| ○ Cerebrovascular Disease | 74.4 |
| ○ Chronic Lower Respiratory Diseases | 81.3 |
| ○ Diabetes Mellitus | 90.5 |
| Influenza & Pneumonia | 26.2 |
| ○ Intentional Self-Harm (Suicide) | 55.7 |
| ○ Chronic Liver Disease and Cirrhosis | 59.8 |
| ○ Infant Mortality | 14.63 |

Leading Causes of Death

| Leading Causes of Death | Total Deaths |
|--|--------------|
| T1. Heart Disease | 64 |
| T1. Accidents | 64 |
| 3. Malignant Neoplasms(Cancer) | 62 |
| 4. Diabetes | 30 |
| 5. Intentional Self-Harm (Suicide) | 25 |
| 6. Chronic Lower Respiratory Diseases | 22 |
| 7. Chronic Liver Disease and Cirrhosis | 21 |
| 8. Cerebrovascular Diseases | 19 |
| 9. Assault (Homicide) | 11 |
| 10. Septicemia | 9 |

| | |
|--------------------------------------|------|
| Percent of Deaths due to tobacco use | 15.9 |
| Median age at death | 59 |

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Tripp County

Demographic Information



Tripp County is located along the Nebraska border in south central South Dakota and averages 3.5 persons per square mile.

2013 Population Information

| Subject | Number | Percent |
|--|--------|---------|
| Total population | 5,498 | 100.0 |
| White | 4,570 | 83.1 |
| Black or African American | 19 | 0.3 |
| American Indian & Alaska Native | 760 | 13.8 |
| Asian | 24 | 0.4 |
| Native Hawaiian & Other Pacific Islander | 1 | 0.0 |
| Two or More Races | 124 | 2.3 |
| Under 5 years | 335 | 6.1 |
| Under 18 years | 1,231 | 22.4 |
| 65 years and over | 1,191 | 21.7 |

Source: United States Census Bureau, 2013 Population Estimates

Health Status Indicators 2009-2013

Nativity

| | |
|---|------|
| Percent of Low Birth Weight Infants | 7.9 |
| Percent of Mothers Receiving Care in 1st Trimester | 64.5 |
| Percent of Mothers Who Used Tobacco While Pregnant ¹ | 21.4 |
| Percent of Births Less Than 37 Wks. of Gestation | 9.5 |
| Average Age of Mother | 26.9 |
| Teenage Birth Rate ² | 13.7 |
| Percent White Births | 63.1 |
| Percent American Indian Births | 32.0 |
| ○ Percent Unmarried | 47.6 |
| ○ Percent WIC births | 52.3 |
| Percent Breastfeeding at discharge | 67.3 |
| ● Percent Payment-Private Insurance | 42.3 |
| ○ Percent Payment-Medicaid | 50.9 |
| ○ Percent C-Section | 36.3 |

● Denotes a health status indicator which is significantly lower than the state average.
○ Denotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

| | |
|-------------------------------------|-------|
| All Causes | 733.5 |
| Heart Disease | 161.8 |
| Malignant Neoplasms (Cancer) | 163.6 |
| Trachea, Bronchus, & Lung | 28.3 |
| Colon, Rectum, & Anus | 19.4 |
| Pancreas | 6.9 |
| Female Breast | 17.8 |
| Leukemia | 8.7 |
| Prostate | 12.5 |
| Accidents | 50.2 |
| Motor Vehicle Accidents | 24.3 |
| Alzheimer's Disease | 47.6 |
| Cerebrovascular Disease | 36.4 |
| Chronic Lower Respiratory Diseases | 35.9 |
| Diabetes Mellitus | 21.3 |
| Influenza & Pneumonia | 20.1 |
| Intentional Self-Harm (Suicide) | LNE |
| Chronic Liver Disease and Cirrhosis | 27.4 |
| Infant Mortality | LNE |

| Leading Causes of Death | Total Deaths |
|---|--------------|
| 1. Heart Disease | 92 |
| 2. Malignant Neoplasms (Cancer) | 76 |
| 3. Alzheimer's Disease | 33 |
| T4. Cerebrovascular Diseases | 20 |
| T4. Accidents | 20 |
| 6. Chronic Lower Respiratory Diseases | 18 |
| 7. Influenza and Pneumonia | 13 |
| 8. Diabetes | 11 |
| 9. Chronic Liver Disease and Cirrhosis | 7 |
| 10. Essential (Primary) Hypertension and Hypertensive Renal Disease | 5 |

| | |
|--------------------------------------|------|
| Percent of Deaths due to tobacco use | 15.9 |
| Median age at death | 83 |

● Denotes a health status indicator which is significantly lower than the state average.

○ Denotes a health status indicator which is significantly higher than the state average.


³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Turner County

Demographic Information



Turner County is located in southeastern South Dakota and averages 13.5 persons per square mile.

2013 Population Information

| Subject | Number | Percent |
|--|--------|---------|
| Total population | 8,361 | 100.0 |
| White | 8,122 | 97.1 |
| Black or African American | 38 | 0.5 |
| American Indian & Alaska Native | 87 | 1.0 |
| Asian | 15 | 0.2 |
| Native Hawaiian & Other Pacific Islander | 4 | 0.0 |
| Two or More Races | 95 | 1.1 |
| Under 5 years | 506 | 6.1 |
| Under 18 years | 1,966 | 23.5 |
| 65 years and over | 1,660 | 19.9 |

Source: United States Census Bureau, 2013 Population Estimates

Health Status Indicators 2009-2013

Nativity

| | |
|---|------|
| ● Percent of Low Birth Weight Infants | 4.0 |
| Percent of Mothers Receiving Care in 1st Trimester | 75.6 |
| Percent of Mothers Who Used Tobacco While Pregnant ¹ | 13.4 |
| Percent of Births Less Than 37 Wks. of Gestation | 7.1 |
| ○ Average Age of Mother | 28.2 |
| Teenage Birth Rate ² | LNE |
| Percent White Births | 99.3 |
| Percent American Indian Births | LNE |
| ● Percent Unmarried | 25.9 |
| ● Percent WIC births | 27.5 |
| Percent Breastfeeding at discharge | 77.9 |
| ○ Percent Payment-Private Insurance | 72.4 |
| ● Percent Payment-Medicaid | 24.1 |
| Percent C-Section | 28.6 |

●Denotes a health status indicator which is significantly lower than the state average.
○Denotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

| | |
|-------------------------------------|-------|
| All Causes | 677.8 |
| Heart Disease | 158.7 |
| Malignant Neoplasms (Cancer) | 153.1 |
| Trachea, Bronchus, & Lung | 35.8 |
| Colon, Rectum, & Anus | 12.3 |
| Pancreas | 7.4 |
| Female Breast | 23.0 |
| Leukemia | 9.6 |
| Prostate | 21.2 |
| Accidents | 42.6 |
| Motor Vehicle Accidents | 19.4 |
| ○ Alzheimer's Disease | 64.1 |
| ● Cerebrovascular Disease | 20.7 |
| Chronic Lower Respiratory Diseases | 59.0 |
| Diabetes Mellitus | 17.1 |
| Influenza & Pneumonia | 11.8 |
| Intentional Self-Harm (Suicide) | 17.9 |
| Chronic Liver Disease and Cirrhosis | LNE |
| Infant Mortality | LNE |

Leading Causes of Death

| Leading Causes of Death | Total Deaths |
|---------------------------------------|--------------|
| 1. Heart Disease | 119 |
| 2. Malignant Neoplasms (Cancer) | 97 |
| 3. Alzheimer's Disease | 55 |
| 4. Chronic Lower Respiratory Diseases | 41 |
| 5. Accidents | 20 |
| 6. Cerebrovascular Diseases | 17 |
| 7. Diabetes | 12 |
| 8. Influenza and Pneumonia | 11 |
| 9. Intentional Self-Harm (Suicide) | 8 |
| 10. Unspecified Dementia | 7 |

| | |
|--------------------------------------|------|
| Percent of Deaths due to tobacco use | 15.8 |
| Median age at death | 83 |

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Union County

Demographic Information



Union County is located in the southeastern corner of the state and averages 31.3 persons per square mile.

2013 Population Information

| Subject | Number | Percent |
|--|--------|---------|
| Total population | 14,829 | 100.0 |
| White | 14,146 | 95.4 |
| Black or African American | 186 | 1.3 |
| American Indian & Alaska Native | 104 | 0.7 |
| Asian | 175 | 1.2 |
| Native Hawaiian & Other Pacific Islander | 9 | 0.1 |
| Two or More Races | 209 | 1.4 |
| Under 5 years | 932 | 6.3 |
| Under 18 years | 3,788 | 25.5 |
| 65 years and over | 2,233 | 15.1 |

Source: United States Census Bureau, 2013 Population Estimates

Health Status Indicators 2009-2013

Nativity

| | |
|---|------|
| Percent of Low Birth Weight Infants | 7.9 |
| ○ Percent of Mothers Receiving Care in 1st Trimester | 85.4 |
| ● Percent of Mothers Who Used Tobacco While Pregnant ¹ | 12.3 |
| Percent of Births Less Than 37 Wks. of Gestation | 10.3 |
| ○ Average Age of Mother | 28.7 |
| ● Teenage Birth Rate ² | 5.2 |
| Percent White Births | 95.9 |
| Percent American Indian Births | 0.5 |
| ● Percent Unmarried | 23.9 |
| ● Percent WIC births | 20.7 |
| Percent Breastfeeding at discharge | 80.0 |
| ○ Percent Payment-Private Insurance | 74.8 |
| ● Percent Payment-Medicaid | 21.7 |
| ○ Percent C-Section | 34.1 |

●Denotes a health status indicator which is significantly lower than the state average.
 ○Denotes a health status indicator which is significantly higher than the state average.
¹Data for mothers who used tobacco are self-reported.
²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

| | |
|-------------------------------------|-------|
| ● All Causes | 623.5 |
| Heart Disease | 157.0 |
| Malignant Neoplasms (Cancer) | 164.2 |
| Trachea, Bronchus, & Lung | 45.8 |
| Colon, Rectum, & Anus | 20.4 |
| Pancreas | 10.1 |
| Female Breast | 26.8 |
| Leukemia | 7.5 |
| Prostate | 23.4 |
| ● Accidents | 30.1 |
| ● Motor Vehicle Accidents | 7.7 |
| Alzheimer's Disease | 33.4 |
| ● Cerebrovascular Disease | 23.9 |
| Chronic Lower Respiratory Diseases | 39.9 |
| Diabetes Mellitus | 15.3 |
| Influenza & Pneumonia | 14.8 |
| ● Intentional Self-Harm (Suicide) | 7.9 |
| Chronic Liver Disease and Cirrhosis | 6.3 |
| Infant Mortality | LNE |

Leading Causes of Death


| Leading Causes of Death | Total Deaths |
|--|--------------|
| T1. Heart Disease | 140 |
| T1. Malignant Neoplasms (Cancer) | 140 |
| 3. Chronic Lower Respiratory Diseases | 35 |
| 4. Alzheimer's Disease | 30 |
| 5. Accidents | 24 |
| 6. Cerebrovascular Diseases | 22 |
| 7. Unspecified Dementia | 17 |
| 8. Diabetes | 13 |
| 9. Influenza and Pneumonia | 12 |
| T10. Essential (Primary) Hypertension and Hypertensive Renal Disease | 8 |
| T10. Other Interstitial Pulmonary Diseases with Fibrosis | 8 |
| Percent of Deaths due to tobacco use | 16.7 |
| Median age at death | 81 |

●Denotes a health status indicator which is significantly lower than the state average.
 ○Denotes a health status indicator which is significantly higher than the state average.
³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.
 Source: South Dakota Department of Health, Office of Health Statistics

Walworth County

Demographic Information



Walworth County is located in north central South Dakota, near the North Dakota border and averages 7.7 persons per square mile.

2013 Population Information

| Subject | Number | Percent |
|--|--------|---------|
| Total population | 5,524 | 100.0 |
| White | 4,541 | 82.2 |
| Black or African American | 24 | 0.4 |
| American Indian & Alaska Native | 772 | 14.0 |
| Asian | 16 | 0.3 |
| Native Hawaiian & Other Pacific Islander | 0 | 0.0 |
| Two or More Races | 171 | 3.1 |
| Under 5 years | 331 | 6.0 |
| Under 18 years | 1,236 | 22.4 |
| 65 years and over | 1,314 | 23.8 |

Source: United States Census Bureau, 2013 Population Estimates

Health Status Indicators 2009-2013

Natality

| | |
|---|------|
| ● Percent of Low Birth Weight Infants | 3.9 |
| Percent of Mothers Receiving Care in 1st Trimester | 61.8 |
| Percent of Mothers Who Used Tobacco While Pregnant ¹ | 21.4 |
| Percent of Births Less Than 37 Wks. of Gestation | 7.3 |
| ● Average Age of Mother | 25.9 |
| Teenage Birth Rate ² | 18.9 |
| Percent White Births | 61.3 |
| Percent American Indian Births | 34.1 |
| ○ Percent Unmarried | 48.5 |
| ○ Percent WIC births | 48.9 |
| Percent Breastfeeding at discharge | 70.6 |
| ● Percent Payment-Private Insurance | 40.5 |
| ○ Percent Payment-Medicaid | 51.4 |
| Percent C-Section | 24.9 |

●Denotes a health status indicator which is significantly lower than the state average.
○Denotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

| | |
|-------------------------------------|-------|
| All Causes | 725.9 |
| Heart Disease | 133.0 |
| Malignant Neoplasms (Cancer) | 148.2 |
| Trachea, Bronchus, & Lung | 43.9 |
| Colon, Rectum, & Anus | 17.5 |
| Pancreas | 5.6 |
| Female Breast | 19.3 |
| Leukemia | 6.8 |
| Prostate | 12.9 |
| Accidents | 43.8 |
| Motor Vehicle Accidents | 14.6 |
| Alzheimer's Disease | 60.3 |
| Cerebrovascular Disease | 43.4 |
| Chronic Lower Respiratory Diseases | 44.3 |
| Diabetes Mellitus | 29.0 |
| Influenza & Pneumonia | 19.8 |
| Intentional Self-Harm (Suicide) | LNE |
| Chronic Liver Disease and Cirrhosis | 20.9 |
| Infant Mortality | LNE |

Leading Causes of Death

| Leading Causes of Death | Total Deaths |
|--|--------------|
| T1. Heart Disease | 76 |
| T1. Malignant Neoplasms (Cancer) | 76 |
| 3. Alzheimer's Disease | 42 |
| 4. Cerebrovascular Diseases | 25 |
| 5. Chronic Lower Respiratory Diseases | 24 |
| 6. Accidents | 16 |
| 7. Diabetes | 15 |
| 8. Influenza and Pneumonia | 11 |
| T9. Chronic Liver Disease and Cirrhosis | 7 |
| T9. Neoplasms of Uncertain or Unknown Behavior | 7 |

| | |
|--------------------------------------|------|
| Percent of Deaths due to tobacco use | 17.0 |
| Median age at death | 82 |

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Yankton County

Demographic Information



Yankton County is located in southeastern South Dakota on the Nebraska border and averages 43.1 persons per square mile.

2013 Population Information

| Subject | Number | Percent |
|--|--------|---------|
| Total population | 22,696 | 100.0 |
| White | 21,095 | 92.9 |
| Black or African American | 473 | 2.1 |
| American Indian & Alaska Native | 651 | 2.9 |
| Asian | 141 | 0.6 |
| Native Hawaiian & Other Pacific Islander | 9 | 0.0 |
| Two or More Races | 327 | 1.4 |
| Under 5 years | 1,313 | 5.8 |
| Under 18 years | 4,866 | 21.4 |
| 65 years and over | 3,873 | 17.1 |

Source: United States Census Bureau, 2013 Population Estimates

Health Status Indicators 2009-2013

Nativity

| | |
|---|------|
| Percent of Low Birth Weight Infants | 6.6 |
| ○ Percent of Mothers Receiving Care in 1st Trimester | 82.8 |
| ○ Percent of Mothers Who Used Tobacco While Pregnant ¹ | 21.5 |
| Percent of Births Less Than 37 Wks. of Gestation | 8.0 |
| Average Age of Mother | 26.9 |
| Teenage Birth Rate ² | 13.0 |
| Percent White Births | 91.1 |
| Percent American Indian Births | 4.4 |
| Percent Unmarried | 40.2 |
| Percent WIC births | 39.2 |
| ● Percent Breastfeeding at discharge | 68.0 |
| ○ Percent Payment-Private Insurance | 64.0 |
| Percent Payment-Medicaid | 33.5 |
| Percent C-Section | 26.5 |

Mortality³

| | |
|-------------------------------------|-------|
| ● All Causes | 632.1 |
| Heart Disease | 148.9 |
| ● Malignant Neoplasms (Cancer) | 137.9 |
| Trachea, Bronchus, & Lung | 43.6 |
| Colon, Rectum, & Anus | 12.1 |
| Pancreas | 8.1 |
| Female Breast | 12.7 |
| Leukemia | 9.6 |
| Prostate | 14.7 |
| Accidents | 38.6 |
| Motor Vehicle Accidents | 15.0 |
| Alzheimer's Disease | 31.9 |
| Cerebrovascular Disease | 30.1 |
| Chronic Lower Respiratory Diseases | 51.0 |
| Diabetes Mellitus | 27.9 |
| Influenza & Pneumonia | 11.9 |
| Intentional Self-Harm (Suicide) | 17.6 |
| Chronic Liver Disease and Cirrhosis | 6.5 |
| Infant Mortality | 5.41 |

Leading Causes of Death

| Leading Causes of Death | Total Deaths |
|---------------------------------------|--------------|
| 1. Heart Disease | 257 |
| 2. Malignant Neoplasms (Cancer) | 205 |
| 3. Chronic Lower Respiratory Diseases | 78 |
| 4. Alzheimer's Disease | 64 |
| 5. Cerebrovascular Diseases | 54 |
| 6. Accidents | 52 |
| 7. Diabetes | 47 |
| 8. Influenza and Pneumonia | 22 |
| 9. Intentional Self-Harm (Suicide) | 20 |
| 10. Unspecified Dementia | 17 |

| | |
|--------------------------------------|------|
| Percent of Deaths due to tobacco use | 21.3 |
| Median age at death | 82 |

● Denotes a health status indicator which is significantly lower than the state average.

○ Denotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

● Denotes a health status indicator which is significantly lower than the state average.

○ Denotes a health status indicator which is significantly higher than the state average.


³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

Ziebach County

Demographic Information



Ziebach County is located in north central South Dakota and averages 1.4 persons per square mile.

2013 Population Information

| Subject | Number | Percent |
|--|--------|---------|
| Total population | 2,834 | 100.0 |
| White | 689 | 24.3 |
| Black or African American | 11 | 0.4 |
| American Indian & Alaska Native | 2,030 | 71.6 |
| Asian | 8 | 0.3 |
| Native Hawaiian & Other Pacific Islander | 1 | 0.0 |
| Two or More Races | 95 | 3.4 |
| Under 5 years | 206 | 7.3 |
| Under 18 years | 1,040 | 36.7 |
| 65 years and over | 198 | 7.0 |

Source: United States Census Bureau, 2013 Population Estimates

Health Status Indicators 2009-2013

Natality

| | |
|---|------|
| Percent of Low Birth Weight Infants | 7.9 |
| ● Percent of Mothers Receiving Care in 1st Trimester | 45.3 |
| ○ Percent of Mothers Who Used Tobacco While Pregnant ¹ | 24.0 |
| Percent of Births Less Than 37 Wks. of Gestation | 9.8 |
| ● Average Age of Mother | 25.8 |
| ○ Teenage Birth Rate ² | 40.8 |
| Percent White Births | 11.0 |
| Percent American Indian Births | 85.5 |
| ○ Percent Unmarried | 69.2 |
| ○ Percent WIC births | 81.3 |
| ● Percent Breastfeeding at discharge | 52.4 |
| ● Percent Payment-Private Insurance | 13.1 |
| ○ Percent Payment-Medicaid | 80.2 |
| ○ Percent C-Section | 33.9 |

●Denotes a health status indicator which is significantly lower than the state average.
○Denotes a health status indicator which is significantly higher than the state average.

¹Data for mothers who used tobacco are self-reported.

²Teenage Birth rate is live births per 1,000 females age 15-17.

Mortality³

| | |
|-------------------------------------|-------|
| All Causes | 877.3 |
| Heart Disease | 249.5 |
| ● Malignant Neoplasms (Cancer) | 89.1 |
| Trachea, Bronchus, & Lung | LNE |
| Colon, Rectum, & Anus | LNE |
| Pancreas | LNE |
| Female Breast | LNE |
| Leukemia | LNE |
| Prostate | LNE |
| ○ Accidents | 107.3 |
| Motor Vehicle Accidents | 47.3 |
| Alzheimer's Disease | LNE |
| Cerebrovascular Disease | 45.6 |
| Chronic Lower Respiratory Diseases | 58.6 |
| Diabetes Mellitus | 56.1 |
| Influenza & Pneumonia | 20.7 |
| Intentional Self-Harm (Suicide) | LNE |
| Chronic Liver Disease and Cirrhosis | 40.1 |
| Infant Mortality | 22.03 |

Leading Causes of Death

| Leading Causes of Death | Total Deaths |
|---------------------------------|--------------|
| 1. Heart Disease | 19 |
| 2. Accidents | 12 |
| 3. Malignant Neoplasms (Cancer) | 7 |

| | |
|--------------------------------------|-----|
| Percent of Deaths due to tobacco use | 8.0 |
| Median age at death | 71 |

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

³All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics