

The following table shows the percent of those who binge drink for various health behaviors and conditions. For example, 43 percent of respondents who use smokeless tobacco binge drink, while 18 percent of respondents who do not use smokeless tobacco binge drink.

**Binge Drinking for Selected Health Behaviors and Conditions, 2011-2014**

	<b>Binge Drinker</b>
Underweight (BMI < 18.5)	12%
Recommended Weight (BMI = 18.5-24.9)	21%
Overweight (BMI = 25.0-29.9)	21%
Obese - Class I (BMI = 30.0-34.9)	20%
Obese - Classes II & III (BMI = 35.0+)	18%
Watching Salt Intake	13%
Not Watching Salt Intake	20%
Participates in Leisure Time Exercise	21%
Does Not Participate in Leisure Time Exercise	18%
Current Smoker	35%
Former Smoker	17%
Never Smoked	16%
Uses Smokeless Tobacco	43%
Does Not Use Smokeless Tobacco	18%
Heard of "South Dakota Quitline"	19%
Not Heard of "South Dakota Quitline"	12%
Ever Diagnosed with Diabetes	9%
Never Diagnosed with Diabetes	21%
Has High Blood Pressure	15%
Does Not Have High Blood Pressure	22%
Has Health Insurance (18-64)	23%
Does Not Have Health Insurance (18-64)	27%
Employer Based Coverage (18-64)	23%
Private Health Plan (18-64)	20%
Medicare (18-64)	10%
Medicaid or Medical Assistance (18-64)	16%
The military, CHAMPUS, TriCare, or the VA (18-64)	23%
The Indian Health Service (18-64)	31%
Breast Exam Within Past Two Years	13%
No Breast Exam Within Past Two Years	11%
Mammogram Within Past Two Years (40+)	8%
No Mammogram Within Past Two Years (40+)	10%
Pap Smear Within Past Three Years	16%
No Pap Smear Within Past Three Years	12%
Blood Stool Test Within Past Two Years (50+)	8%
No Blood Stool Test Within Past Two Years (50+)	10%
Ever Had a Sigmoidoscopy or Colonoscopy (50+)	8%
Never Had a Sigmoidoscopy or Colonoscopy (50+)	13%
PSA Test Within Past Two Years (40+)	11%
No PSA Test Within Past Two Years (40+)	22%
Had a Flu Shot in Past 12 Months (65+)	3%
Did Not Have a Flu Shot in Past 12 Months (65+)	4%
Ever Had a Pneumonia Shot (65+)	2%
Never Had a Pneumonia Shot (65+)	5%
Ever Had a Shingles Shot (50+)	5%
Never Had a Shingles Shot (50+)	13%
Previously Had Heart Attack	8%
Never Had Heart Attack	20%

### Binge Drinking for Selected Health Behaviors and Conditions, 2011-2014

	<b>Binge Drinker</b>
Have Angina or Coronary Heart Disease	9%
Does Not Have Angina or Coronary Heart Disease	20%
Previously Had a Stroke	7%
Never Had a Stroke	20%
Ever Diagnosed with Cancer (Excluding Skin Cancer)	9%
Never Diagnosed with Cancer	21%
Ever Diagnosed with Skin Cancer	8%
Never Diagnosed with Skin Cancer	21%
Uses Sun Block Most of the Time	14%
Does Not Use Sun Block Most of the Time	21%
Current Asthma	16%
Former Asthma	23%
Never Had Asthma	20%
Ever Diagnosed with Arthritis	11%
Never Diagnosed with Arthritis	23%
Ever Diagnosed with COPD	12%
Never Diagnosed with COPD	20%
Ever Diagnosed with Depression	18%
Never Diagnosed with Depression	20%
Ever Diagnosed with Kidney Disease	8%
Never Diagnosed with Kidney Disease	20%
Has Severe Vision Impairment	11%
Does Not Have Severe Vision Impairment	19%
Recent Increased Confusion/Memory Loss	21%
No Recent Increased Confusion/Memory Loss	18%
< 6 hours of sleep	18%
6+ hours of sleep	18%
Been to the Dentist in the Past Year	19%
Hasn't Been to the Dentist in the Past Year	20%
Always/Almost Always Wears Seat Belt	17%
Sometimes/Seldom/Never Wears Seat Belt	31%
Drank Alcohol in Past 30 Days	35%
Did Not Drink Alcohol in Past 30 Days	0%
Heavy Drinker	85%
Not a Heavy Drinker	16%
Good/Very Good/Excellent Health Status	21%
Fair/Poor Health Status	12%
Physical Health Not Good for 0-29 Days of the Past 30	20%
Physical Health Not Good for 30 Days of the Past 30	10%
Mental Health Not Good for 0-19 Days of the Past 30	20%
Mental Health Not Good for 20-30 Days of the Past 30	23%
Usual Activities Unattainable for 0-9 Days of the Past 30	20%
Usual Activities Unattainable for 10-30 Days of the Past 30	14%
Has Physical, Mental, or Emotional Problems	12%
Does Not Have Physical, Mental, or Emotional Problems	22%
Has a Disability with Special Equipment Needed	8%
Does Not Have a Disability with Special Equipment Needed	21%
Injured in a Fall in Past 12 Months (45+)	10%
Not Injured in a Fall in Past 12 Months (45+)	12%
Ever Been Tested for HIV	25%
Never Been Tested for HIV	18%

Source: South Dakota Behavioral Risk Factor Surveillance System, 2011-2014