

The following table shows the percent of those who have COPD for various health behaviors and conditions. For example, 11 percent of respondents who are underweight have COPD, while five percent of respondents who are within the recommended weight have COPD.

COPD Prevalence for Selected Health Behaviors and Conditions, 2011-2014

	Ever Diagnosed with COPD
Underweight (BMI < 18.5)	11%
Recommended Weight (BMI = 18.5-24.9)	5%
Overweight (BMI = 25.0-29.9)	4%
Obese - Class I (BMI = 30.0-34.9)	6%
Obese - Classes II & III (BMI = 35.0+)	8%
Watching Salt Intake	7%
Not Watching Salt Intake	5%
Participates in Leisure Time Exercise	4%
Does Not Participate in Leisure Time Exercise	9%
Current Smoker	10%
Former Smoker	8%
Never Smoked	2%
Uses Smokeless Tobacco	5%
Does Not Use Smokeless Tobacco	5%
Heard of "South Dakota Quitline"	5%
Not Heard of "South Dakota Quitline"	5%
Ever Diagnosed with Diabetes	12%
Never Diagnosed with Diabetes	4%
Has High Blood Pressure	9%
Does Not Have High Blood Pressure	3%
Has Health Insurance (18-64)	3%
Does Not Have Health Insurance (18-64)	7%
Employer Based Coverage (18-64)	2%
Private Health Plan (18-64)	2%
Medicare (18-64)	16%
Medicaid or Medical Assistance (18-64)	9%
The military, CHAMPUS, TriCare, or the VA (18-64)	8%
The Indian Health Service (18-64)	3%
Breast Exam Within Past Two Years	5%
No Breast Exam Within Past Two Years	8%
Mammogram Within Past Two Years (40+)	6%
No Mammogram Within Past Two Years (40+)	10%
Pap Smear Within Past Three Years	4%
No Pap Smear Within Past Three Years	7%
Blood Stool Test Within Past Two Years (50+)	10%
No Blood Stool Test Within Past Two Years (50+)	8%
Ever Had a Sigmoidoscopy or Colonoscopy (50+)	8%
Never Had a Sigmoidoscopy or Colonoscopy (50+)	9%
PSA Test Within Past Two Years (40+)	9%
No PSA Test Within Past Two Years (40+)	6%
Had a Flu Shot in Past 12 Months (65+)	12%
Did Not Have a Flu Shot in Past 12 Months (65+)	8%
Ever Had a Pneumonia Shot (65+)	14%
Never Had a Pneumonia Shot (65+)	5%
Ever Had a Shingles Shot (50+)	11%
Never Had a Shingles Shot (50+)	8%
Previously Had Heart Attack	18%
Never Had Heart Attack	4%
Have Angina or Coronary Heart Disease	18%
Does Not Have Angina or Coronary Heart Disease	4%

COPD Prevalence for Selected Health Behaviors and Conditions, 2011-2014

	Ever Diagnosed with COPD
Previously Had a Stroke	19%
Never Had a Stroke	5%
Ever Diagnosed with Cancer (Excluding Skin Cancer)	12%
Never Diagnosed with Cancer	5%
Ever Diagnosed with Skin Cancer	11%
Never Diagnosed with Skin Cancer	5%
Uses Sun Block Most of the Time	4%
Does Not Use Sun Block Most of the Time	5%
Current Asthma	24%
Former Asthma	8%
Never Had Asthma	3%
Ever Diagnosed with Arthritis	11%
Never Diagnosed with Arthritis	3%
Ever Diagnosed with Depression	12%
Never Diagnosed with Depression	4%
Ever Diagnosed with Kidney Disease	16%
Never Diagnosed with Kidney Disease	5%
Has Severe Vision Impairment	20%
Does Not Have Severe Vision Impairment	4%
Recent Increased Confusion/Memory Loss	18%
No Recent Increased Confusion/Memory Loss	5%
< 6 hours of sleep	11%
6+ hours of sleep	4%
Been to the Dentist in the Past Year	4%
Hasn't Been to the Dentist in the Past Year	8%
Always/Almost Always Wears Seat Belt	5%
Sometimes/Seldom/Never Wears Seat Belt	7%
Drank Alcohol in Past 30 Days	4%
Did Not Drink Alcohol in Past 30 Days	7%
Binge Drinker	3%
Not a Binge Drinker	6%
Heavy Drinker	6%
Not a Heavy Drinker	5%
Good/Very Good/Excellent Health Status	3%
Fair/Poor Health Status	20%
Physical Health Not Good for 0-29 Days of the Past 30	4%
Physical Health Not Good for 30 Days of the Past 30	22%
Mental Health Not Good for 0-19 Days of the Past 30	4%
Mental Health Not Good for 20-30 Days of the Past 30	14%
Usual Activities Unattainable for 0-9 Days of the Past 30	4%
Usual Activities Unattainable for 10-30 Days of the Past 30	21%
Has Physical, Mental, or Emotional Problems	15%
Does Not Have Physical, Mental, or Emotional Problems	3%
Has a Disability with Special Equipment Needed	19%
Does Not Have a Disability with Special Equipment Needed	4%
Injured in a Fall in Past 12 Months (45+)	17%
Not Injured in a Fall in Past 12 Months (45+)	6%
Ever Been Tested for HIV	6%
Never Been Tested for HIV	5%

Source: South Dakota Behavioral Risk Factor Surveillance System, 2011-2014