

The following table shows the percent of those who have been diagnosed with kidney disease for various health behaviors and conditions. For example, 10 percent of respondents who have had a heart attack have kidney disease, while two percent of respondents who have not had a heart attack have kidney disease.

Kidney Disease Prevalence for Selected Health Behaviors and Conditions, 2011-2014

	Ever Diagnosed with Kidney Disease
Underweight (BMI < 18.5)	3%
Recommended Weight (BMI = 18.5-24.9)	1%
Overweight (BMI = 25.0-29.9)	2%
Obese - Class I (BMI = 30.0-34.9)	3%
Obese - Classes II & III (BMI = 35.0+)	4%
Watching Salt Intake	3%
Not Watching Salt Intake	1%
Participates in Leisure Time Exercise	2%
Does Not Participate in Leisure Time Exercise	3%
Current Smoker	2%
Former Smoker	3%
Never Smoked	2%
Uses Smokeless Tobacco	2%
Does Not Use Smokeless Tobacco	2%
Heard of "South Dakota Quitline"	2%
Not Heard of "South Dakota Quitline"	3%
Ever Diagnosed with Diabetes	7%
Never Diagnosed with Diabetes	2%
Has High Blood Pressure	5%
Does Not Have High Blood Pressure	1%
Has Health Insurance (18-64)	2%
Does Not Have Health Insurance (18-64)	2%
Employer Based Coverage (18-64)	1%
Private Health Plan (18-64)	1%
Medicare (18-64)	8%
Medicaid or Medical Assistance (18-64)	6%
The military, CHAMPUS, TriCare, or the VA (18-64)	2%
The Indian Health Service (18-64)	2%
Breast Exam Within Past Two Years	2%
No Breast Exam Within Past Two Years	3%
Mammogram Within Past Two Years (40+)	3%
No Mammogram Within Past Two Years (40+)	3%
Pap Smear Within Past Three Years	2%
No Pap Smear Within Past Three Years	2%
Blood Stool Test Within Past Two Years (50+)	7%
No Blood Stool Test Within Past Two Years (50+)	3%
Ever Had a Sigmoidoscopy or Colonoscopy (50+)	4%
Never Had a Sigmoidoscopy or Colonoscopy (50+)	2%
PSA Test Within Past Two Years (40+)	4%
No PSA Test Within Past Two Years (40+)	3%
Had a Flu Shot in Past 12 Months (65+)	6%
Did Not Have a Flu Shot in Past 12 Months (65+)	3%
Ever Had a Pneumonia Shot (65+)	6%
Never Had a Pneumonia Shot (65+)	2%
Ever Had a Shingles Shot (50+)	5%
Never Had a Shingles Shot (50+)	3%
Previously Had Heart Attack	10%
Never Had Heart Attack	2%

Kidney Disease Prevalence for Selected Health Behaviors and Conditions, 2011-2014

	Ever Diagnosed with Kidney Disease
Have Angina or Coronary Heart Disease	10%
Does Not Have Angina or Coronary Heart Disease	2%
Previously Had a Stroke	9%
Never Had a Stroke	2%
Ever Diagnosed with Cancer (Excluding Skin Cancer)	8%
Never Diagnosed with Cancer	2%
Ever Diagnosed with Skin Cancer	6%
Never Diagnosed with Skin Cancer	2%
Uses Sun Block Most of the Time	2%
Does Not Use Sun Block Most of the Time	2%
Current Asthma	4%
Former Asthma	2%
Never Had Asthma	2%
Ever Diagnosed with Arthritis	5%
Never Diagnosed with Arthritis	1%
Ever Diagnosed with COPD	7%
Never Diagnosed with COPD	2%
Ever Diagnosed with Depression	4%
Never Diagnosed with Depression	2%
Has Severe Vision Impairment	9%
Does Not Have Severe Vision Impairment	2%
Recent Increased Confusion/Memory Loss	9%
No Recent Increased Confusion/Memory Loss	2%
< 6 hours of sleep	5%
6+ hours of sleep	2%
Been to the Dentist in the Past Year	2%
Hasn't Been to the Dentist in the Past Year	3%
Always/Almost Always Wears Seat Belt	2%
Sometimes/Seldom/Never Wears Seat Belt	2%
Drank Alcohol in Past 30 Days	1%
Did Not Drink Alcohol in Past 30 Days	3%
Binge Drinker	1%
Not a Binge Drinker	3%
Heavy Drinker	1%
Not a Heavy Drinker	2%
Good/Very Good/Excellent Health Status	1%
Fair/Poor Health Status	7%
Physical Health Not Good for 0-29 Days of the Past 30	2%
Physical Health Not Good for 30 Days of the Past 30	8%
Mental Health Not Good for 0-19 Days of the Past 30	2%
Mental Health Not Good for 20-30 Days of the Past 30	4%
Usual Activities Unattainable for 0-9 Days of the Past 30	2%
Usual Activities Unattainable for 10-30 Days of the Past 30	8%
Has Physical, Mental, or Emotional Problems	6%
Does Not Have Physical, Mental, or Emotional Problems	1%
Has a Disability with Special Equipment Needed	9%
Does Not Have a Disability with Special Equipment Needed	2%
Injured in a Fall in Past 12 Months (45+)	6%
Not Injured in a Fall in Past 12 Months (45+)	3%
Ever Been Tested for HIV	2%
Never Been Tested for HIV	2%

Source: South Dakota Behavioral Risk Factor Surveillance System, 2011-2014