

The following table shows the percent of those, ages 40 or older, who have not had a PSA test within the past two years for various health behaviors and conditions. For example, 63 percent of respondents who have health insurance have not had a PSA test within the past two years, while 93 percent of respondents who do not have health insurance have not had a PSA test within the past two years.

Prevalence of No PSA Test Within the Past Two Years (Ages 40+) for Selected Health Behaviors and Conditions, 2011-2014

	No PSA Test Within Past Two Years (40+)
Participates in Leisure Time Exercise	55%
Does Not Participate in Leisure Time Exercise	57%
Watching Salt Intake	52%
Not Watching Salt Intake	59%
Current Smoker	69%
Former Smoker	47%
Never Smoked	58%
Uses Smokeless Tobacco	69%
Does Not Use Smokeless Tobacco	55%
Heard of "South Dakota Quitline"	56%
Not Heard of "South Dakota Quitline"	54%
Ever Diagnosed with Diabetes	47%
Never Diagnosed with Diabetes	57%
Has High Blood Pressure	48%
Does Not Have High Blood Pressure	61%
Has Health Insurance (18-64)	63%
Does Not Have Health Insurance (18-64)	93%
Blood Stool Test Within Past Two Years (50+)	28%
No Blood Stool Test Within Past Two Years (50+)	47%
Ever Had a Sigmoidoscopy or Colonoscopy (50+)	32%
Never Had a Sigmoidoscopy or Colonoscopy (50+)	68%
Had a Flu Shot in Past 12 Months (65+)	27%
Did Not Have a Flu Shot in Past 12 Months (65+)	44%
Ever Had a Pneumonia Shot (65+)	25%
Never Had a Pneumonia Shot (65+)	47%
Ever Had a Shingles Shot (50+)	26%
Never Had a Shingles Shot (50+)	52%
Previously Had Heart Attack	41%
Never Had Heart Attack	57%
Have Angina or Coronary Heart Disease	38%
Does Not Have Angina or Coronary Heart Disease	58%
Previously Had a Stroke	41%
Never Had a Stroke	56%
Ever Diagnosed with Cancer (Excluding Skin Cancer)	33%
Never Diagnosed with Cancer	58%
Ever Diagnosed with Skin Cancer	40%
Never Diagnosed with Skin Cancer	57%
Uses Sun Block Most of the Time	44%
Does Not Use Sun Block Most of the Time	57%
Ever Diagnosed with Arthritis	46%
Never Diagnosed with Arthritis	60%

Prevalence of No PSA Test Within the Past Two Years (Ages 40+) for Selected Health Behaviors and Conditions, 2011-2014

	No PSA Test Within Past Two Years (40+)
Ever Diagnosed with COPD	44%
Never Diagnosed with COPD	57%
Ever Diagnosed with Depression	55%
Never Diagnosed with Depression	56%
Ever Diagnosed with Kidney Disease	44%
Never Diagnosed with Kidney Disease	56%
Has Severe Vision Impairment	48%
Does Not Have Severe Vision Impairment	56%
< 6 hours of sleep	62%
6+ hours of sleep	55%
Been to the Dentist in the Past Year	51%
Hasn't Been to the Dentist in the Past Year	65%
Always/Almost Always Wears Seat Belt	53%
Sometimes/Seldom/Never Wears Seat Belt	64%
Drank Alcohol in Past 30 Days	56%
Did Not Drink Alcohol in Past 30 Days	55%
Binge Drinker	71%
Not a Binge Drinker	52%
Heavy Drinker	72%
Not a Heavy Drinker	55%
Good/Very Good/Excellent Health Status	56%
Fair/Poor Health Status	53%
Physical Health Not Good for 0-29 Days of the Past 30	57%
Physical Health Not Good for 30 Days of the Past 30	47%
Mental Health Not Good for 0-19 Days of the Past 30	56%
Mental Health Not Good for 20-30 Days of the Past 30	66%
Usual Activities Unattainable for 0-9 Days of the Past 30	56%
Usual Activities Unattainable for 10-30 Days of the Past 30	52%
Has Physical, Mental, or Emotional Problems	51%
Does Not Have Physical, Mental, or Emotional Problems	57%
Has a Disability with Special Equipment Needed	47%
Does Not Have a Disability with Special Equipment Needed	57%
Injured in a Fall in Past 12 Months (45+)	51%
Not Injured in a Fall in Past 12 Months (45+)	50%
Ever Been Tested for HIV	64%
Never Been Tested for HIV	54%

Source: South Dakota Behavioral Risk Factor Surveillance System, 2011-2014