

The following table shows the percent of those who sometimes, seldom, or never wear a seat belt for various health behaviors and conditions. For example, 29 percent of respondents who currently smoke say they sometimes, seldom, or never wear a seat belt, while 13 percent of respondents who have never smoked say they sometimes, seldom, or never wear a seat belt.

Prevalence of Lack of Seat Belt Use for Selected Health Behaviors and Conditions, 2011-2014

| | Sometimes/Seldom/Never Wears Seat Belt |
|---|---|
| Underweight (BMI < 18.5) | 23% |
| Recommended Weight (BMI = 18.5-24.9) | 15% |
| Overweight (BMI = 25.0-29.9) | 17% |
| Obese - Class I (BMI = 30.0-34.9) | 19% |
| Obese - Classes II & III (BMI = 35.0+) | 24% |
| Watching Salt Intake | 14% |
| Not Watching Salt Intake | 21% |
| Participates in Leisure Time Exercise | 15% |
| Does Not Participate in Leisure Time Exercise | 24% |
| Current Smoker | 29% |
| Former Smoker | 16% |
| Never Smoked | 13% |
| Uses Smokeless Tobacco | 38% |
| Does Not Use Smokeless Tobacco | 16% |
| Heard of "South Dakota Quitline" | 17% |
| Not Heard of "South Dakota Quitline" | 18% |
| Ever Diagnosed with Diabetes | 17% |
| Never Diagnosed with Diabetes | 17% |
| Has High Blood Pressure | 18% |
| Does Not Have High Blood Pressure | 17% |
| Has Health Insurance (18-64) | 17% |
| Does Not Have Health Insurance (18-64) | 29% |
| Employer Based Coverage (18-64) | 14% |
| Private Health Plan (18-64) | 23% |
| Medicare (18-64) | 25% |
| Medicaid or Medical Assistance (18-64) | 29% |
| The military, CHAMPUS, TriCare, or the VA (18-64) | 11% |
| The Indian Health Service (18-64) | 20% |
| Breast Exam Within Past Two Years | 9% |
| No Breast Exam Within Past Two Years | 16% |
| Mammogram Within Past Two Years (40+) | 7% |
| No Mammogram Within Past Two Years (40+) | 15% |
| Pap Smear Within Past Three Years | 10% |
| No Pap Smear Within Past Three Years | 14% |
| Blood Stool Test Within Past Two Years (50+) | 10% |
| No Blood Stool Test Within Past Two Years (50+) | 16% |
| Ever Had a Sigmoidoscopy or Colonoscopy (50+) | 13% |
| Never Had a Sigmoidoscopy or Colonoscopy (50+) | 20% |
| PSA Test Within Past Two Years (40+) | 18% |
| No PSA Test Within Past Two Years (40+) | 25% |
| Had a Flu Shot in Past 12 Months (65+) | 11% |
| Did Not Have a Flu Shot in Past 12 Months (65+) | 18% |
| Ever Had a Pneumonia Shot (65+) | 11% |
| Never Had a Pneumonia Shot (65+) | 17% |
| Ever Had a Shingles Shot (50+) | 10% |
| Never Had a Shingles Shot (50+) | 18% |
| Previously Had Heart Attack | 20% |
| Never Had Heart Attack | 17% |

Prevalence of Lack of Seat Belt Use for Selected Health Behaviors and Conditions, 2011-2014

| | Sometimes/Seldom/Never Wears Seat Belt |
|---|---|
| Have Angina or Coronary Heart Disease | 17% |
| Does Not Have Angina or Coronary Heart Disease | 17% |
| Previously Had a Stroke | 15% |
| Never Had a Stroke | 17% |
| Ever Diagnosed with Cancer (Excluding Skin Cancer) | 13% |
| Never Diagnosed with Cancer | 18% |
| Ever Diagnosed with Skin Cancer | 13% |
| Never Diagnosed with Skin Cancer | 18% |
| Uses Sun Block Most of the Time | 7% |
| Does Not Use Sun Block Most of the Time | 22% |
| Current Asthma | 21% |
| Former Asthma | 19% |
| Never Had Asthma | 17% |
| Ever Diagnosed with Arthritis | 16% |
| Never Diagnosed with Arthritis | 18% |
| Ever Diagnosed with COPD | 23% |
| Never Diagnosed with COPD | 17% |
| Ever Diagnosed with Depression | 17% |
| Never Diagnosed with Depression | 17% |
| Ever Diagnosed with Kidney Disease | 14% |
| Never Diagnosed with Kidney Disease | 17% |
| Has Severe Vision Impairment | 22% |
| Does Not Have Severe Vision Impairment | 17% |
| Recent Increased Confusion/Memory Loss | 22% |
| No Recent Increased Confusion/Memory Loss | 17% |
| < 6 hours of sleep | 25% |
| 6+ hours of sleep | 16% |
| Been to the Dentist in the Past Year | 14% |
| Hasn't Been to the Dentist in the Past Year | 25% |
| Drank Alcohol in Past 30 Days | 18% |
| Did Not Drink Alcohol in Past 30 Days | 16% |
| Binge Drinker | 27% |
| Not a Binge Drinker | 15% |
| Heavy Drinker | 31% |
| Not a Heavy Drinker | 16% |
| Good/Very Good/Excellent Health Status | 17% |
| Fair/Poor Health Status | 21% |
| Physical Health Not Good for 0-29 Days of the Past 30 | 17% |
| Physical Health Not Good for 30 Days of the Past 30 | 18% |
| Mental Health Not Good for 0-19 Days of the Past 30 | 17% |
| Mental Health Not Good for 20-30 Days of the Past 30 | 24% |
| Usual Activities Unattainable for 0-9 Days of the Past 30 | 17% |
| Usual Activities Unattainable for 10-30 Days of the Past 30 | 18% |
| Has Physical, Mental, or Emotional Problems | 20% |
| Does Not Have Physical, Mental, or Emotional Problems | 17% |
| Has a Disability with Special Equipment Needed | 18% |
| Does Not Have a Disability with Special Equipment Needed | 17% |
| Injured in a Fall in Past 12 Months (45+) | 16% |
| Not Injured in a Fall in Past 12 Months (45+) | 15% |
| Ever Been Tested for HIV | 18% |
| Never Been Tested for HIV | 17% |

Source: South Dakota Behavioral Risk Factor Surveillance System, 2011-2014