

The following table shows the percent of those who are obese (BMI 35+) for various health behaviors and conditions. For example, 17 percent of respondents who have been diagnosed with depression are obese (BMI 35+), while nine percent of respondents who have not been diagnosed with depression are obese (BMI 35+).

Class II-III Obesity Prevalence for Selected Health Behaviors and Conditions, 2011-2014

	Obese - Classes II-III (BMI = 35+)
Participates in Leisure Time Exercise	9%
Does Not Participate in Leisure Time Exercise	15%
Watching Salt Intake	12%
Not Watching Salt Intake	13%
Current Smoker	11%
Former Smoker	12%
Never Smoked	10%
Uses Smokeless Tobacco	11%
Does Not Use Smokeless Tobacco	11%
Heard of "South Dakota Quitline"	13%
Not Heard of "South Dakota Quitline"	10%
Ever Diagnosed with Diabetes	25%
Never Diagnosed with Diabetes	9%
Has High Blood Pressure	17%
Does Not Have High Blood Pressure	8%
Has Health Insurance (18-64)	12%
Does Not Have Health Insurance (18-64)	14%
Employer Based Coverage (18-64)	11%
Private Health Plan (18-64)	8%
Medicare (18-64)	21%
Medicaid or Medical Assistance (18-64)	17%
The military, CHAMPUS, TriCare, or the VA (18-64)	10%
The Indian Health Service (18-64)	15%
Breast Exam Within Past Two Years	12%
No Breast Exam Within Past Two Years	12%
Mammogram Within Past Two Years (40+)	12%
No Mammogram Within Past Two Years (40+)	12%
Pap Smear Within Past Three Years	12%
No Pap Smear Within Past Three Years	10%
Blood Stool Test Within Past Two Years (50+)	11%
No Blood Stool Test Within Past Two Years (50+)	12%
Ever Had a Sigmoidoscopy or Colonoscopy (50+)	11%
Never Had a Sigmoidoscopy or Colonoscopy (50+)	13%
PSA Test Within Past Two Years (40+)	13%
No PSA Test Within Past Two Years (40+)	12%
Had a Flu Shot in Past 12 Months (65+)	9%
Did Not Have a Flu Shot in Past 12 Months (65+)	8%
Ever Had a Pneumonia Shot (65+)	9%
Never Had a Pneumonia Shot (65+)	8%
Ever Had a Shingles Shot (50+)	9%
Never Had a Shingles Shot (50+)	14%
Previously Had Heart Attack	12%
Never Had Heart Attack	11%
Have Angina or Coronary Heart Disease	14%
Does Not Have Angina or Coronary Heart Disease	11%
Previously Had a Stroke	12%
Never Had a Stroke	11%
Ever Diagnosed with Cancer (Excluding Skin Cancer)	13%
Never Diagnosed with Cancer	11%

Class II-III Obesity Prevalence for Selected Health Behaviors and Conditions, 2011-2014

	Obese - Classes II-III (BMI = 35+)
Ever Diagnosed with Skin Cancer	8%
Never Diagnosed with Skin Cancer	11%
Uses Sun Block Most of the Time	9%
Does Not Use Sun Block Most of the Time	12%
Current Asthma	20%
Former Asthma	14%
Never Had Asthma	10%
Ever Diagnosed with Arthritis	15%
Never Diagnosed with Arthritis	9%
Ever Diagnosed with COPD	17%
Never Diagnosed with COPD	10%
Ever Diagnosed with Depression	17%
Never Diagnosed with Depression	9%
Ever Diagnosed with Kidney Disease	20%
Never Diagnosed with Kidney Disease	10%
Has Severe Vision Impairment	14%
Does Not Have Severe Vision Impairment	11%
Recent Increased Confusion/Memory Loss	17%
No Recent Increased Confusion/Memory Loss	12%
< 6 hours of sleep	16%
6+ hours of sleep	11%
Been to the Dentist in the Past Year	10%
Hasn't Been to the Dentist in the Past Year	14%
Always/Almost Always Wears Seat Belt	10%
Sometimes/Seldom/Never Wears Seat Belt	15%
Drank Alcohol in Past 30 Days	10%
Did Not Drink Alcohol in Past 30 Days	12%
Binge Drinker	10%
Not a Binge Drinker	11%
Heavy Drinker	10%
Not a Heavy Drinker	11%
Good/Very Good/Excellent Health Status	9%
Fair/Poor Health Status	20%
Physical Health Not Good for 0-29 Days of the Past 30	10%
Physical Health Not Good for 30 Days of the Past 30	17%
Mental Health Not Good for 0-19 Days of the Past 30	10%
Mental Health Not Good for 20-30 Days of the Past 30	18%
Usual Activities Unattainable for 0-9 Days of the Past 30	10%
Usual Activities Unattainable for 10-30 Days of the Past 30	19%
Has Physical, Mental, or Emotional Problems	17%
Does Not Have Physical, Mental, or Emotional Problems	9%
Has a Disability with Special Equipment Needed	21%
Does Not Have a Disability with Special Equipment Needed	10%
Injured in a Fall in Past 12 Months (45+)	16%
Not Injured in a Fall in Past 12 Months (45+)	11%
Ever Been Tested for HIV	12%
Never Been Tested for HIV	10%

Source: South Dakota Behavioral Risk Factor Surveillance System, 2011-2014