

The following table shows the percent of respondents, age 65+, who have not had a flu shot in the past 12 months for various health behaviors and conditions. For example, 27 percent of respondents who report they always wear a seatbelt have not had a flu shot in the past 12 months, while 39 percent of respondents who do not always wear a seatbelt have not had a flu shot in the past 12 months.

Prevalence of No Flu Shot (Ages 65+) For Selected Health Behaviors and Conditions, 2011-2015

	Did Not Have a Flu Shot in Past 12 Months (65+)
Underweight (BMI < 18.5)	44%
Recommended (BMI = 18.5-24.9)	33%
Overweight (BMI = 25.0-29.9)	29%
Obese - Class I (BMI = 30.0-34.9)	29%
Obese - Classes II & III (BMI = 35.0+)	30%
Participates in Leisure Time Exercise	29%
Does Not Participate in Leisure Time Exercise	34%
Meets Physical Activity Recommendations	28%
Does Not Meet Physical Activity Recommendations	31%
7+ Exercise Trips Per Week	48%
Less Than 7 Exercise Trips Per Week	27%
12+ Hours Sitting Per Day	18%
< 12 Hours Sitting Per Day	29%
Has 2+ Servings of Fruit per Day	25%
Has Less Than 2 Servings of Fruit per Day	32%
Has 3+ Servings of Vegetables per Day	39%
Has Less Than 3 Servings of Vegetables per Day	29%
Has 5+ Servings of Fruits and Vegetables per Day	27%
Has Less Than 5 Servings of Fruits and Vegetables per Day	30%
Current Smoker	41%
Former Smoker	28%
Never Smoked	31%
Uses Smokeless Tobacco	48%
Does Not Use Smokeless Tobacco	30%
Heard of "South Dakota Quitline"	27%
Not Heard of "South Dakota Quitline"	34%
Ever Diagnosed with Diabetes	24%
Never Diagnosed with Diabetes	32%
Has High Blood Pressure	26%
Does Not Have High Blood Pressure	36%
Has High Blood Cholesterol	26%
Does Not Have High Blood Cholesterol	30%
Ever Had a Pneumonia Shot (65+)	19%
Never Had a Pneumonia Shot (65+)	54%
Previously Had Heart Attack	27%
Never Had Heart Attack	31%
Knows Symptoms of Heart Attack	32%
Does Not Know Symptoms of Heart Attack	30%
Has Angina or Coronary Heart Disease	25%
Does Not Have Angina or Coronary Heart Disease	32%
Previously Had a Stroke	24%
Never Had a Stroke	31%
Ever Diagnosed with Cancer (Excluding Skin Cancer)	28%
Never Diagnosed with Cancer	31%
Ever Diagnosed with Skin Cancer	24%
Never Diagnosed with Skin Cancer	32%

Prevalence of No Flu Shot (Ages 65+) For Selected Health Behaviors and Conditions, 2011-2015

	Did Not Have a Flu Shot in Past 12 Months (65+)
Current Asthma	25%
Former Asthma	27%
Never Had Asthma	31%
Ever Diagnosed with Arthritis	27%
Never Diagnosed with Arthritis	34%
Ever Diagnosed with COPD	21%
Never Diagnosed with COPD	32%
Ever Diagnosed with Depression	30%
Never Diagnosed with Depression	31%
Ever Diagnosed with Kidney Disease	21%
Never Diagnosed with Kidney Disease	31%
Has Severe Vision Impairment	27%
Does Not Have Severe Vision Impairment	29%
Recent Increased Confusion/Memory Loss (45+)	27%
No Recent Increased Confusion/Memory Loss (45+)	29%
Always Wears Seat Belt	27%
Does Not Always Wear Seat Belt	39%
Drank Alcohol in Past 30 Days	29%
Did Not Drink Alcohol in Past 30 Days	31%
Binge Drinker	40%
Not a Binge Drinker	30%
Heavy Drinker	42%
Not a Heavy Drinker	30%
Has an Advance Directive	26%
Does Not Have an Advance Directive	35%
Good/Very Good/Excellent Health Status	32%
Fair/Poor Health Status	26%
Physical Health Not Good for 0-29 Days of the Past 30	31%
Physical Health Not Good for 30 Days of the Past 30	28%
Mental Health Not Good for 0-19 Days of the Past 30	31%
Mental Health Not Good for 20-30 Days of the Past 30	24%
Usual Activities Unattainable for 0-9 Days of the Past 30	31%
Usual Activities Unattainable for 10-30 Days of the Past 30	28%
Has Physical, Mental, or Emotional Problems	27%
Does Not Have Physical, Mental, or Emotional Problems	32%
Has a Disability with Special Equipment Needed	25%
Does Not Have a Disability with Special Equipment Needed	32%
Ever Been Tested for HIV	31%
Never Been Tested for HIV	31%

Source: South Dakota Behavioral Risk Factor Surveillance System, 2011-2015