

The following table shows the percent of respondents, age 65+, who have not had a pneumonia shot in the past 12 months for various health behaviors and conditions. For example, 21 percent of respondents who previously had a stroke have not had a pneumonia shot in the past 12 months, while 34 percent of respondents who have not had a stroke have not had a pneumonia shot in the past 12 months.

Prevalence of No Pneumonia Shot (65+) For Selected Health Behaviors and Conditions, 2011-2015

	Never Had a Pneumonia Shot (65+)
Underweight (BMI < 18.5)	35%
Recommended (BMI = 18.5-24.9)	35%
Overweight (BMI = 25.0-29.9)	34%
Obese - Class I (BMI = 30.0-34.9)	28%
Obese - Classes II & III (BMI = 35.0+)	27%
Participates in Leisure Time Exercise	33%
Does Not Participate in Leisure Time Exercise	33%
Meets Physical Activity Recommendations	32%
Does Not Meet Physical Activity Recommendations	32%
7+ Exercise Trips Per Week	40%
Less Than 7 Exercise Trips Per Week	29%
12+ Hours Sitting Per Day	23%
< 12 Hours Sitting Per Day	30%
Has 2+ Servings of Fruit per Day	29%
Has Less Than 2 Servings of Fruit per Day	34%
Has 3+ Servings of Vegetables per Day	35%
Has Less Than 3 Servings of Vegetables per Day	32%
Has 5+ Servings of Fruits and Vegetables per Day	28%
Has Less Than 5 Servings of Fruits and Vegetables per Day	33%
Current Smoker	40%
Former Smoker	29%
Never Smoked	35%
Uses Smokeless Tobacco	41%
Does Not Use Smokeless Tobacco	33%
Heard of "South Dakota Quitline"	31%
Not Heard of "South Dakota Quitline"	33%
Ever Diagnosed with Diabetes	23%
Never Diagnosed with Diabetes	35%
Has High Blood Pressure	27%
Does Not Have High Blood Pressure	41%
Has High Blood Cholesterol	28%
Does Not Have High Blood Cholesterol	33%
Had a Flu Shot in Past 12 Months (65+)	22%
Did Not Have a Flu Shot in Past 12 Months (65+)	58%
Previously Had Heart Attack	22%
Never Had Heart Attack	35%
Knows Symptoms of Heart Attack	28%
Does Not Know Symptoms of Heart Attack	33%
Has Angina or Coronary Heart Disease	20%
Does Not Have Angina or Coronary Heart Disease	35%
Previously Had a Stroke	21%
Never Had a Stroke	34%
Ever Diagnosed with Cancer (Excluding Skin Cancer)	24%
Never Diagnosed with Cancer	35%

Prevalence of No Pneumonia Shot (65+) For Selected Health Behaviors and Conditions, 2011-2015

	Never Had a Pneumonia Shot (65+)
Ever Diagnosed with Skin Cancer	24%
Never Diagnosed with Skin Cancer	35%
Current Asthma	14%
Former Asthma	37%
Never Had Asthma	34%
Ever Diagnosed with Arthritis	27%
Never Diagnosed with Arthritis	39%
Ever Diagnosed with COPD	13%
Never Diagnosed with COPD	35%
Ever Diagnosed with Depression	28%
Never Diagnosed with Depression	34%
Ever Diagnosed with Kidney Disease	15%
Never Diagnosed with Kidney Disease	34%
Has Severe Vision Impairment	22%
Does Not Have Severe Vision Impairment	32%
Recent Increased Confusion/Memory Loss (45+)	22%
No Recent Increased Confusion/Memory Loss (45+)	32%
Always Wears Seat Belt	31%
Does Not Always Wear Seat Belt	38%
Drank Alcohol in Past 30 Days	34%
Did Not Drink Alcohol in Past 30 Days	32%
Binge Drinker	50%
Not a Binge Drinker	32%
Heavy Drinker	41%
Not a Heavy Drinker	32%
Has an Advance Directive	26%
Does Not Have an Advance Directive	36%
Good/Very Good/Excellent Health Status	36%
Fair/Poor Health Status	23%
Physical Health Not Good for 0-29 Days of the Past 30	34%
Physical Health Not Good for 30 Days of the Past 30	22%
Mental Health Not Good for 0-19 Days of the Past 30	33%
Mental Health Not Good for 20-30 Days of the Past 30	29%
Usual Activities Unattainable for 0-9 Days of the Past 30	34%
Usual Activities Unattainable for 10-30 Days of the Past 30	23%
Has Physical, Mental, or Emotional Problems	25%
Does Not Have Physical, Mental, or Emotional Problems	37%
Has a Disability with Special Equipment Needed	22%
Does Not Have a Disability with Special Equipment Needed	35%
Ever Been Tested for HIV	27%
Never Been Tested for HIV	33%

Source: South Dakota Behavioral Risk Factor Surveillance System, 2011-2015