

The following table shows the percent of respondents who always wear a seatbelt for various health behaviors and conditions. For example, 50 percent of respondents who are heavy drinkers always wear a seatbelt, while 66 percent of respondents who are not heavy drinkers always wear a seatbelt.

**Prevalence of Seatbelt Use For Selected Health Behaviors and Conditions, 2011-2015**

	<b>Always Wears Seat Belt</b>
Underweight (BMI < 18.5)	68%
Recommended (BMI = 18.5-24.9)	69%
Overweight (BMI = 25.0-29.9)	64%
Obese - Class I (BMI = 30.0-34.9)	62%
Obese - Classes II & III (BMI = 35.0+)	58%
Participates in Leisure Time Exercise	67%
Does Not Participate in Leisure Time Exercise	57%
Meets Physical Activity Recommendations	71%
Does Not Meet Physical Activity Recommendations	61%
7+ Exercise Trips Per Week	64%
Less Than 7 Exercise Trips Per Week	69%
12+ Hours Sitting Per Day	69%
< 12 Hours Sitting Per Day	69%
Has 2+ Servings of Fruit per Day	76%
Has Less Than 2 Servings of Fruit per Day	63%
Has 3+ Servings of Vegetables per Day	76%
Has Less Than 3 Servings of Vegetables per Day	65%
Has 5+ Servings of Fruits and Vegetables per Day	77%
Has Less Than 5 Servings of Fruits and Vegetables per Day	65%
Current Smoker	53%
Former Smoker	66%
Never Smoked	69%
Uses Smokeless Tobacco	41%
Does Not Use Smokeless Tobacco	66%
Current E-Cigarette User	62%
Not E-Cigarette User	66%
Heard of "South Dakota Quitline"	65%
Not Heard of "South Dakota Quitline"	67%
Ever Diagnosed with Diabetes	66%
Never Diagnosed with Diabetes	64%
Has High Blood Pressure	65%
Does Not Have High Blood Pressure	65%
Has High Blood Cholesterol	69%
Does Not Have High Blood Cholesterol	70%
Has Health Insurance (18-64)	65%
Does Not Have Health Insurance (18-64)	52%
Employer Based Coverage (18-64)	67%
Private Health Plan (18-64)	55%
Medicare (18-64)	64%
Medicaid or Medical Assistance (18-64)	58%
The military, CHAMPUS, TriCare, or the VA (18-64)	73%
The Indian Health Service (18-64)	66%
Had a Flu Shot in Past 12 Months (65+)	75%
Did Not Have a Flu Shot in Past 12 Months (65+)	63%
Ever Had a Pneumonia Shot (65+)	74%
Never Had a Pneumonia Shot (65+)	67%
Previously Had Heart Attack	65%
Never Had Heart Attack	65%

**Prevalence of Seatbelt Use For Selected Health Behaviors and Conditions, 2011-2015**

	<b>Always Wears Seat Belt</b>
Knows Symptoms of Heart Attack	72%
Does Not Know Symptoms of Heart Attack	66%
Has Angina or Coronary Heart Disease	67%
Does Not Have Angina or Coronary Heart Disease	65%
Previously Had a Stroke	68%
Never Had a Stroke	65%
Ever Diagnosed with Cancer (Excluding Skin Cancer)	70%
Never Diagnosed with Cancer	64%
Ever Diagnosed with Skin Cancer	71%
Never Diagnosed with Skin Cancer	64%
Current Asthma	65%
Former Asthma	61%
Never Had Asthma	65%
Ever Diagnosed with Arthritis	67%
Never Diagnosed with Arthritis	64%
Ever Diagnosed with COPD	60%
Never Diagnosed with COPD	65%
Ever Diagnosed with Depression	65%
Never Diagnosed with Depression	65%
Ever Diagnosed with Kidney Disease	64%
Never Diagnosed with Kidney Disease	65%
Has Severe Vision Impairment	70%
Does Not Have Severe Vision Impairment	66%
Recent Increased Confusion/Memory Loss (45+)	63%
No Recent Increased Confusion/Memory Loss (45+)	69%
Drank Alcohol in Past 30 Days	62%
Did Not Drink Alcohol in Past 30 Days	68%
Binge Drinker	51%
Not a Binge Drinker	68%
Heavy Drinker	50%
Not a Heavy Drinker	66%
Has an Advance Directive	75%
Does Not Have an Advance Directive	66%
Good/Very Good/Excellent Health Status	65%
Fair/Poor Health Status	63%
Physical Health Not Good for 0-29 Days of the Past 30	64%
Physical Health Not Good for 30 Days of the Past 30	66%
Mental Health Not Good for 0-19 Days of the Past 30	65%
Mental Health Not Good for 20-30 Days of the Past 30	60%
Usual Activities Unattainable for 0-9 Days of the Past 30	65%
Usual Activities Unattainable for 10-30 Days of the Past 30	65%
Has Physical, Mental, or Emotional Problems	65%
Does Not Have Physical, Mental, or Emotional Problems	65%
Has a Disability with Special Equipment Needed	68%
Does Not Have a Disability with Special Equipment Needed	64%
Ever Been Tested for HIV	65%
Never Been Tested for HIV	64%

Source: South Dakota Behavioral Risk Factor Surveillance System, 2011-2015