




# South Dakota

## Demographic Information



South Dakota is located in the north central portion of the United States and averages 10.7 persons per square mile.

### 2016 Population Information

Subject	Number	Percent
Total population	865,454	100.0
White	737,070	85.2
Black or African American	17,302	2.0
American Indian & Alaska Native	77,711	9.0
Asian	12,767	1.5
Native Hawaiian & Other Pacific Islander	656	0.1
Two or More Races	19,948	2.3
Under 5 years	61,369	7.1
Under 18 years	213,287	24.6
65 years and over	138,805	16.0

Source: United States Census Bureau, 2016 Population Estimates

## Health Status Indicators 2012-2016

### Natality

● Percent of Low Birth Weight Infants	6.4
○ Percent of Mothers Receiving Care in 1st Trimester	72.0
○ Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	14.8
● Percent of Births Less Than 37 Wks. of Gestation	8.4
Average Age of Mother	27.7
○ Teenage Birth Rate <sup>2</sup>	13.0
Percent White Births	76.1
Percent American Indian Births	16.0
● Percent Unmarried	37.7
● Percent WIC births	34.7
Percent Breastfeeding at discharge	78.5
○ Percent Payment-Private Insurance	57.9
● Percent Payment-Medicaid	32.3
● Percent C-Section	25.3

●Denotes a health status indicator which is significantly lower than the national average.  
○Denotes a health status indicator which is significantly higher than the national average.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

● All Causes	705.4
● Heart Disease	152.1
Malignant Neoplasms (Cancer)	157.5
Trachea, Bronchus, & Lung	40.8
○ Colon, Rectum, & Anus	16.3
Female Breast	19.0
Pancreas	10.8
Prostate	18.9
○ Leukemia	7.4
○ Chronic Lower Respiratory Diseases	41.7
○ Alzheimer's Disease	36.5
Cerebrovascular Disease	36.5
○ Accidents	48.9
○ Motor Vehicle Accidents	16.8
○ Diabetes	23.1
○ Influenza and Pneumonia	16.9
○ Intentional Self-Harm (Suicide)	18.3
○ Chronic Liver Disease and Cirrhosis	15.0
○ Infant Mortality	6.63

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	8,408
2. Malignant Neoplasms (Cancer)	8,199
3. Accidents	2,272
4. Chronic Lower Respiratory Diseases	2,259
5. Alzheimer's Disease	2,185
6. Cerebrovascular Disease	2,064
7. Diabetes	1,216
8. Influenza and Pneumonia	962
9. Intentional Self-Harm (Suicide)	757
10. Chronic Liver Disease and Cirrhosis	657
Percent of Deaths due to tobacco use	17.2
Median age at death	80

●Denotes a health status indicator which is significantly lower than the national average.


○Denotes a health status indicator which is significantly higher than the national average.

<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.


Source: South Dakota Department of Health, Office of Health Statistics

# Aurora County

Demographic Information	Health Status Indicators 2012-2016																																																																																																																								
<div style="text-align: center;">  </div> <p>Aurora County is located in the south central portion of the state and averages 3.8 persons per square mile.</p> <h3 style="text-align: center;">2016 Population Information</h3> <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="text-align: left;">Subject</th> <th style="text-align: right;">Number</th> <th style="text-align: right;">Percent</th> </tr> </thead> <tbody> <tr> <td>Total population</td> <td style="text-align: right;">2,736</td> <td style="text-align: right;">100.0</td> </tr> <tr> <td>White</td> <td style="text-align: right;">2,594</td> <td style="text-align: right;">94.8</td> </tr> <tr> <td>Black or African American</td> <td style="text-align: right;">23</td> <td style="text-align: right;">0.8</td> </tr> <tr> <td>American Indian &amp; Alaska Native</td> <td style="text-align: right;">66</td> <td style="text-align: right;">2.4</td> </tr> <tr> <td>Asian</td> <td style="text-align: right;">19</td> <td style="text-align: right;">0.7</td> </tr> <tr> <td>Native Hawaiian &amp; Other Pacific Islander</td> <td style="text-align: right;">0</td> <td style="text-align: right;">0.0</td> </tr> <tr> <td>Two or More Races</td> <td style="text-align: right;">34</td> <td style="text-align: right;">1.2</td> </tr> <tr> <td>Under 5 years</td> <td style="text-align: right;">177</td> <td style="text-align: right;">6.5</td> </tr> <tr> <td>Under 18 years</td> <td style="text-align: right;">716</td> <td style="text-align: right;">26.2</td> </tr> <tr> <td>65 years and over</td> <td style="text-align: right;">557</td> <td style="text-align: right;">20.4</td> </tr> </tbody> </table> <p>Source: United States Census Bureau, 2016 Population Estimates</p>	Subject	Number	Percent	Total population	2,736	100.0	White	2,594	94.8	Black or African American	23	0.8	American Indian & Alaska Native	66	2.4	Asian	19	0.7	Native Hawaiian & Other Pacific Islander	0	0.0	Two or More Races	34	1.2	Under 5 years	177	6.5	Under 18 years	716	26.2	65 years and over	557	20.4	<h3 style="text-align: center;">Nativity</h3> <table border="1" style="width: 100%; border-collapse: collapse;"> <tbody> <tr> <td>Percent of Low Birth Weight Infants</td> <td style="text-align: right;">6.7</td> </tr> <tr> <td>Percent of Mothers Receiving Care in 1st Trimester</td> <td style="text-align: right;">69.9</td> </tr> <tr> <td>● Percent of Mothers Who Used Tobacco While Pregnant<sup>1</sup></td> <td style="text-align: right;">9.8</td> </tr> <tr> <td>Percent of Births Less Than 37 Wks. of Gestation</td> <td style="text-align: right;">9.8</td> </tr> <tr> <td>Average Age of Mother</td> <td style="text-align: right;">28.1</td> </tr> <tr> <td>Teenage Birth Rate<sup>2</sup></td> <td style="text-align: right;">LNE</td> </tr> <tr> <td>Percent White Births</td> <td style="text-align: right;">99.0</td> </tr> <tr> <td>Percent American Indian Births</td> <td style="text-align: right;">LNE</td> </tr> <tr> <td>● Percent Unmarried</td> <td style="text-align: right;">25.3</td> </tr> <tr> <td>Percent WIC births</td> <td style="text-align: right;">37.2</td> </tr> <tr> <td>Percent Breastfeeding at discharge</td> <td style="text-align: right;">76.3</td> </tr> <tr> <td>○ Percent Payment-Private Insurance</td> <td style="text-align: right;">73.6</td> </tr> <tr> <td>● Percent Payment-Medicaid</td> <td style="text-align: right;">17.1</td> </tr> <tr> <td>Percent C-Section</td> <td style="text-align: right;">30.4</td> </tr> </tbody> </table> <p>●Denotes a health status indicator which is significantly lower than the state average. ○Denotes a health status indicator which is significantly higher than the state average.</p> <p><sup>1</sup>Data for mothers who used tobacco are self-reported. <sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.</p>	Percent of Low Birth Weight Infants	6.7	Percent of Mothers Receiving Care in 1st Trimester	69.9	● Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	9.8	Percent of Births Less Than 37 Wks. of Gestation	9.8	Average Age of Mother	28.1	Teenage Birth Rate <sup>2</sup>	LNE	Percent White Births	99.0	Percent American Indian Births	LNE	● Percent Unmarried	25.3	Percent WIC births	37.2	Percent Breastfeeding at discharge	76.3	○ Percent Payment-Private Insurance	73.6	● Percent Payment-Medicaid	17.1	Percent C-Section	30.4	<h3 style="text-align: center;">Mortality<sup>3</sup></h3> <table border="1" style="width: 100%; border-collapse: collapse;"> <tbody> <tr> <td>● All Causes</td> <td style="text-align: right;">596.1</td> </tr> <tr> <td>Heart Disease</td> <td style="text-align: right;">140.2</td> </tr> <tr> <td>Malignant Neoplasms (Cancer)</td> <td style="text-align: right;">137.7</td> </tr> <tr> <td>Trachea, Bronchus, &amp; Lung</td> <td style="text-align: right;">48.7</td> </tr> <tr> <td>Colon, Rectum, &amp; Anus</td> <td style="text-align: right;">11.8</td> </tr> <tr> <td>Female Breast</td> <td style="text-align: right;">59.5</td> </tr> <tr> <td>Pancreas</td> <td style="text-align: right;">LNE</td> </tr> <tr> <td>Prostate</td> <td style="text-align: right;">39.5</td> </tr> <tr> <td>Leukemia</td> <td style="text-align: right;">LNE</td> </tr> <tr> <td>Chronic Lower Respiratory Diseases</td> <td style="text-align: right;">LNE</td> </tr> <tr> <td>Alzheimer's Disease</td> <td style="text-align: right;">37.9</td> </tr> <tr> <td>Cerebrovascular Disease</td> <td style="text-align: right;">58.1</td> </tr> <tr> <td>Accidents</td> <td style="text-align: right;">48.1</td> </tr> <tr> <td>Motor Vehicle Accidents</td> <td style="text-align: right;">33.3</td> </tr> <tr> <td>Diabetes</td> <td style="text-align: right;">26.1</td> </tr> <tr> <td>Influenza and Pneumonia</td> <td style="text-align: right;">21.5</td> </tr> <tr> <td>Intentional Self-Harm (Suicide)</td> <td style="text-align: right;">LNE</td> </tr> <tr> <td>Chronic Liver Disease and Cirrhosis</td> <td style="text-align: right;">LNE</td> </tr> <tr> <td>Infant Mortality</td> <td style="text-align: right;">LNE</td> </tr> </tbody> </table> <h3 style="text-align: center;">Leading Causes of Death</h3> <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="text-align: left;">Leading Causes of Death</th> <th style="text-align: right;">Total Deaths</th> </tr> </thead> <tbody> <tr> <td>1. Heart Disease</td> <td style="text-align: right;">33</td> </tr> <tr> <td>2. Malignant Neoplasms (Cancer)</td> <td style="text-align: right;">29</td> </tr> <tr> <td>3. Cerebrovascular Disease</td> <td style="text-align: right;">13</td> </tr> <tr> <td>4. Alzheimer's Disease</td> <td style="text-align: right;">10</td> </tr> <tr> <td>5. Accidents</td> <td style="text-align: right;">7</td> </tr> <tr> <td>T6. Diabetes</td> <td style="text-align: right;">6</td> </tr> <tr> <td>T6. Influenza and Pneumonia</td> <td style="text-align: right;">6</td> </tr> <tr> <td>Percent of Deaths due to tobacco use</td> <td style="text-align: right;">12.8</td> </tr> <tr> <td>Median age at death</td> <td style="text-align: right;">83</td> </tr> </tbody> </table> <p>●Denotes a health status indicator which is significantly lower than the state average. ○Denotes a health status indicator which is significantly higher than the state average. <sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.</p> <p>See technical notes for more information. Source: South Dakota Department of Health, Office of Health Statistics</p>	● All Causes	596.1	Heart Disease	140.2	Malignant Neoplasms (Cancer)	137.7	Trachea, Bronchus, & Lung	48.7	Colon, Rectum, & Anus	11.8	Female Breast	59.5	Pancreas	LNE	Prostate	39.5	Leukemia	LNE	Chronic Lower Respiratory Diseases	LNE	Alzheimer's Disease	37.9	Cerebrovascular Disease	58.1	Accidents	48.1	Motor Vehicle Accidents	33.3	Diabetes	26.1	Influenza and Pneumonia	21.5	Intentional Self-Harm (Suicide)	LNE	Chronic Liver Disease and Cirrhosis	LNE	Infant Mortality	LNE	Leading Causes of Death	Total Deaths	1. Heart Disease	33	2. Malignant Neoplasms (Cancer)	29	3. Cerebrovascular Disease	13	4. Alzheimer's Disease	10	5. Accidents	7	T6. Diabetes	6	T6. Influenza and Pneumonia	6	Percent of Deaths due to tobacco use	12.8	Median age at death	83
Subject	Number	Percent																																																																																																																							
Total population	2,736	100.0																																																																																																																							
White	2,594	94.8																																																																																																																							
Black or African American	23	0.8																																																																																																																							
American Indian & Alaska Native	66	2.4																																																																																																																							
Asian	19	0.7																																																																																																																							
Native Hawaiian & Other Pacific Islander	0	0.0																																																																																																																							
Two or More Races	34	1.2																																																																																																																							
Under 5 years	177	6.5																																																																																																																							
Under 18 years	716	26.2																																																																																																																							
65 years and over	557	20.4																																																																																																																							
Percent of Low Birth Weight Infants	6.7																																																																																																																								
Percent of Mothers Receiving Care in 1st Trimester	69.9																																																																																																																								
● Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	9.8																																																																																																																								
Percent of Births Less Than 37 Wks. of Gestation	9.8																																																																																																																								
Average Age of Mother	28.1																																																																																																																								
Teenage Birth Rate <sup>2</sup>	LNE																																																																																																																								
Percent White Births	99.0																																																																																																																								
Percent American Indian Births	LNE																																																																																																																								
● Percent Unmarried	25.3																																																																																																																								
Percent WIC births	37.2																																																																																																																								
Percent Breastfeeding at discharge	76.3																																																																																																																								
○ Percent Payment-Private Insurance	73.6																																																																																																																								
● Percent Payment-Medicaid	17.1																																																																																																																								
Percent C-Section	30.4																																																																																																																								
● All Causes	596.1																																																																																																																								
Heart Disease	140.2																																																																																																																								
Malignant Neoplasms (Cancer)	137.7																																																																																																																								
Trachea, Bronchus, & Lung	48.7																																																																																																																								
Colon, Rectum, & Anus	11.8																																																																																																																								
Female Breast	59.5																																																																																																																								
Pancreas	LNE																																																																																																																								
Prostate	39.5																																																																																																																								
Leukemia	LNE																																																																																																																								
Chronic Lower Respiratory Diseases	LNE																																																																																																																								
Alzheimer's Disease	37.9																																																																																																																								
Cerebrovascular Disease	58.1																																																																																																																								
Accidents	48.1																																																																																																																								
Motor Vehicle Accidents	33.3																																																																																																																								
Diabetes	26.1																																																																																																																								
Influenza and Pneumonia	21.5																																																																																																																								
Intentional Self-Harm (Suicide)	LNE																																																																																																																								
Chronic Liver Disease and Cirrhosis	LNE																																																																																																																								
Infant Mortality	LNE																																																																																																																								
Leading Causes of Death	Total Deaths																																																																																																																								
1. Heart Disease	33																																																																																																																								
2. Malignant Neoplasms (Cancer)	29																																																																																																																								
3. Cerebrovascular Disease	13																																																																																																																								
4. Alzheimer's Disease	10																																																																																																																								
5. Accidents	7																																																																																																																								
T6. Diabetes	6																																																																																																																								
T6. Influenza and Pneumonia	6																																																																																																																								
Percent of Deaths due to tobacco use	12.8																																																																																																																								
Median age at death	83																																																																																																																								

# Beadle County

## Demographic Information



Beadle County is located in the center of eastern South Dakota and averages 13.8 persons per square mile.

### 2016 Population Information

Subject	Number	Percent
Total population	18,101	100.0
White	15,680	86.6
Black or African American	266	1.5
American Indian & Alaska Native	276	1.5
Asian	1,502	8.3
Native Hawaiian & Other Pacific Islander	41	0.2
Two or More Races	336	1.9
Under 5 years	1,617	8.9
Under 18 years	4,916	27.2
65 years and over	3,124	17.3

Source: United States Census Bureau, 2016 Population Estimates

## Health Status Indicators 2012-2016

### Nativity

Percent of Low Birth Weight Infants	7.2
● Percent of Mothers Receiving Care in 1st Trimester	55.9
Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	13.2
Percent of Births Less Than 37 Wks. of Gestation	8.3
● Average Age of Mother	26.6
○ Teenage Birth Rate <sup>2</sup>	22.5
Percent White Births	77.6
Percent American Indian Births	1.3
Percent Unmarried	38.9
○ Percent WIC births	49.7
Percent Breastfeeding at discharge	75.7
● Percent Payment-Private Insurance	49.6
○ Percent Payment-Medicaid	38.4
○ Percent C-Section	31.3

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

All Causes	721.1
Heart Disease	149.9
Malignant Neoplasms (Cancer)	162.5
Trachea, Bronchus, & Lung	37.4
Colon, Rectum, & Anus	12.5
Female Breast	14.4
Pancreas	11.3
Prostate	21.5
Leukemia	5.0
Chronic Lower Respiratory Diseases	33.5
Alzheimer's Disease	45.7
Cerebrovascular Disease	42.9
Accidents	42.1
Motor Vehicle Accidents	13.1
○ Diabetes	38.5
○ Influenza and Pneumonia	31.0
Intentional Self-Harm (Suicide)	15.9
Chronic Liver Disease and Cirrhosis	13.3
Infant Mortality	10.75

### Leading Causes of Death

	Total Deaths
1. Heart Disease	212
2. Malignant Neoplasms (Cancer)	206
3. Alzheimer's Disease	75
4. Cerebrovascular Disease	65
5. Diabetes	52
6. Accidents	46
7. Influenza and Pneumonia	45
8. Chronic Lower Respiratory Diseases	43
9. Nephritis, nephrotic syndrome, and Nephrosis	14
10. Intentional Self-Harm (Suicide)	13
Percent of Deaths due to tobacco use	19.7
Median age at death	82

●Denotes a health status indicator which is significantly lower than the state average.


○Denotes a health status indicator which is significantly higher than the state average.

<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.


Source: South Dakota Department of Health, Office of Health Statistics

# Bennett County

Demographic Information	Health Status Indicators 2012-2016																																																				
<div style="text-align: center;">  </div> <p>Bennett County is located on the southern border in western South Dakota and averages 2.9 persons per square mile.</p> <h3 style="text-align: center;">2016 Population Information</h3> <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="text-align: left;">Subject</th> <th style="text-align: right;">Number</th> <th style="text-align: right;">Percent</th> </tr> </thead> <tbody> <tr> <td>Total population</td> <td style="text-align: right;">3,460</td> <td style="text-align: right;">100.0</td> </tr> <tr> <td>White</td> <td style="text-align: right;">1,190</td> <td style="text-align: right;">34.4</td> </tr> <tr> <td>Black or African American</td> <td style="text-align: right;">17</td> <td style="text-align: right;">0.5</td> </tr> <tr> <td>American Indian &amp; Alaska Native</td> <td style="text-align: right;">2,045</td> <td style="text-align: right;">59.1</td> </tr> <tr> <td>Asian</td> <td style="text-align: right;">19</td> <td style="text-align: right;">0.5</td> </tr> <tr> <td>Native Hawaiian &amp; Other Pacific Islander</td> <td style="text-align: right;">2</td> <td style="text-align: right;">0.1</td> </tr> <tr> <td>Two or More Races</td> <td style="text-align: right;">187</td> <td style="text-align: right;">5.4</td> </tr> <tr> <td>Under 5 years</td> <td style="text-align: right;">343</td> <td style="text-align: right;">9.9</td> </tr> <tr> <td>Under 18 years</td> <td style="text-align: right;">1,161</td> <td style="text-align: right;">33.6</td> </tr> <tr> <td>65 years and over</td> <td style="text-align: right;">431</td> <td style="text-align: right;">12.5</td> </tr> </tbody> </table>	Subject	Number	Percent	Total population	3,460	100.0	White	1,190	34.4	Black or African American	17	0.5	American Indian & Alaska Native	2,045	59.1	Asian	19	0.5	Native Hawaiian & Other Pacific Islander	2	0.1	Two or More Races	187	5.4	Under 5 years	343	9.9	Under 18 years	1,161	33.6	65 years and over	431	12.5	<h3 style="text-align: center;">Nativity</h3> <ul style="list-style-type: none"> <li>○ Percent of Low Birth Weight Infants 4.6</li> <li>● Percent of Mothers Receiving Care in 1st Trimester 52.4</li> <li>○ Percent of Mothers Who Used Tobacco While Pregnant<sup>1</sup> 25.8</li> <li>○ Percent of Births Less Than 37 Wks. of Gestation 9.6</li> <li>● Average Age of Mother 25.3</li> <li>○ Teenage Birth Rate<sup>2</sup> 41.2</li> <li>○ Percent White Births 11.6</li> <li>○ Percent American Indian Births 80.9</li> <li>○ Percent Unmarried 75.1</li> <li>○ Percent WIC births 70.7</li> <li>● Percent Breastfeeding at discharge 51.2</li> <li>● Percent Payment-Private Insurance 12.5</li> <li>○ Percent Payment-Medicaid 58.1</li> <li>○ Percent C-Section 21.1</li> </ul>	<h3 style="text-align: center;">Mortality<sup>3</sup></h3> <ul style="list-style-type: none"> <li>○ All Causes 1,008.3</li> <li>Heart Disease 182.4</li> <li>Malignant Neoplasms (Cancer) 212.4 <ul style="list-style-type: none"> <li>Trachea, Bronchus, &amp; Lung 25.0</li> <li>Colon, Rectum, &amp; Anus 35.3</li> <li>Female Breast LNE</li> <li>Pancreas 17.6</li> <li>Prostate 36.3</li> <li>Leukemia LNE</li> </ul> </li> <li>Chronic Lower Respiratory Diseases 70.9</li> <li>Alzheimer's Disease 24.1</li> <li>Cerebrovascular Disease 33.9 <ul style="list-style-type: none"> <li>○ Accidents 99.5 <ul style="list-style-type: none"> <li>Motor Vehicle Accidents 37.0</li> </ul> </li> <li>○ Diabetes 65.1</li> </ul> </li> <li>Influenza and Pneumonia 32.4</li> <li>Intentional Self-Harm (Suicide) 23.9</li> <li>Chronic Liver Disease and Cirrhosis 50.1</li> <li>Infant Mortality 20.23</li> </ul> <table border="1" style="width: 100%; border-collapse: collapse; margin-top: 10px;"> <thead> <tr> <th style="text-align: left;">Leading Causes of Death</th> <th style="text-align: right;">Total Deaths</th> </tr> </thead> <tbody> <tr> <td>1. Malignant Neoplasms (Cancer)</td> <td style="text-align: right;">36</td> </tr> <tr> <td>2. Heart Disease</td> <td style="text-align: right;">30</td> </tr> <tr> <td>3. Accidents</td> <td style="text-align: right;">17</td> </tr> <tr> <td>4. Chronic Lower Respiratory Diseases</td> <td style="text-align: right;">12</td> </tr> <tr> <td>5. Diabetes</td> <td style="text-align: right;">11</td> </tr> <tr> <td>6. Chronic Liver Disease and Cirrhosis</td> <td style="text-align: right;">7</td> </tr> <tr> <td>T7. Cerebrovascular Disease</td> <td style="text-align: right;">6</td> </tr> <tr> <td>T7. Influenza and Pneumonia</td> <td style="text-align: right;">6</td> </tr> </tbody> </table> <ul style="list-style-type: none"> <li>Percent of Deaths due to tobacco use 24.1</li> <li>Median age at death 74</li> </ul>	Leading Causes of Death	Total Deaths	1. Malignant Neoplasms (Cancer)	36	2. Heart Disease	30	3. Accidents	17	4. Chronic Lower Respiratory Diseases	12	5. Diabetes	11	6. Chronic Liver Disease and Cirrhosis	7	T7. Cerebrovascular Disease	6	T7. Influenza and Pneumonia	6
Subject	Number	Percent																																																			
Total population	3,460	100.0																																																			
White	1,190	34.4																																																			
Black or African American	17	0.5																																																			
American Indian & Alaska Native	2,045	59.1																																																			
Asian	19	0.5																																																			
Native Hawaiian & Other Pacific Islander	2	0.1																																																			
Two or More Races	187	5.4																																																			
Under 5 years	343	9.9																																																			
Under 18 years	1,161	33.6																																																			
65 years and over	431	12.5																																																			
Leading Causes of Death	Total Deaths																																																				
1. Malignant Neoplasms (Cancer)	36																																																				
2. Heart Disease	30																																																				
3. Accidents	17																																																				
4. Chronic Lower Respiratory Diseases	12																																																				
5. Diabetes	11																																																				
6. Chronic Liver Disease and Cirrhosis	7																																																				
T7. Cerebrovascular Disease	6																																																				
T7. Influenza and Pneumonia	6																																																				
<p>Source: United States Census Bureau, 2016 Population Estimates</p>	<p>●Denotes a health status indicator which is significantly lower than the state average. ○Denotes a health status indicator which is significantly higher than the state average.</p> <p><sup>1</sup>Data for mothers who used tobacco are self-reported. <sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.</p>	<p>○Denotes a health status indicator which is significantly lower than the state average. ○Denotes a health status indicator which is significantly higher than the state average.</p> <p><sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.</p> <p>See technical notes for more information. Source: South Dakota Department of Health, Office of Health Statistics</p>																																																			

# Bon Homme County

## Demographic Information



Bon Homme County is located on the Nebraska border in eastern South Dakota and averages 12.5 persons per square mile.

### 2016 Population Information

Subject	Number	Percent
Total population	6,984	100.0
White	6,225	89.1
Black or African American	90	1.3
American Indian & Alaska Native	554	7.9
Asian	17	0.2
Native Hawaiian & Other Pacific Islander	1	0.0
Two or More Races	97	1.4
Under 5 years	353	5.1
Under 18 years	1,317	18.9
65 years and over	1,422	20.4

Source: United States Census Bureau, 2016 Population Estimates

## Health Status Indicators 2012-2016

### Nativity

Percent of Low Birth Weight Infants	4.8
Percent of Mothers Receiving Care in 1st Trimester	80.2
● Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	9.9
Percent of Births Less Than 37 Wks. of Gestation	8.1
Average Age of Mother	28.1
Teenage Birth Rate <sup>2</sup>	LNE
Percent White Births	94.3
Percent American Indian Births	3.0
● Percent Unmarried	19.5
Percent WIC births	32.9
Percent Breastfeeding at discharge	80.5
○ Percent Payment-Private Insurance	73.9
● Percent Payment-Medicaid	19.8
Percent C-Section	26.3

●Denotes a health status indicator which is significantly lower than the state average.  
○Denotes a health status indicator which is significantly higher than the state average.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

● All Causes	584.6
● Heart Disease	118.4
Malignant Neoplasms (Cancer)	132.7
Trachea, Bronchus, & Lung	31.0
Colon, Rectum, & Anus	12.8
Female Breast	29.0
Pancreas	6.1
Prostate	8.6
Leukemia	11.1
Chronic Lower Respiratory Diseases	41.2
Alzheimer's Disease	24.0
Cerebrovascular Disease	32.3
Accidents	38.0
Motor Vehicle Accidents	18.3
Diabetes	17.9
Influenza and Pneumonia	18.1
Intentional Self-Harm (Suicide)	14.1
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	LNE

### Leading Causes of Death

	Total Deaths
1. Heart Disease	82
2. Malignant Neoplasms (Cancer)	76
T3. Chronic Lower Respiratory Diseases	26
T3. Cerebrovascular Disease	26
5. Alzheimer's Disease	18
6. Accidents	16
7. Influenza and Pneumonia	14
8. Essential (Primary) Hypertension and Hypertensive Renal Disease	13
9. Diabetes	11
10. Unspecified Dementia	10
Percent of Deaths due to tobacco use	12.4
Median age at death	85

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Brookings County

## Demographic Information



Brookings County is located on the eastern border of the state and averages 40.3 persons per square mile.

### 2016 Population Information

Subject	Number	Percent
Total population	34,135	100.0
White	31,770	93.1
Black or African American	558	1.6
American Indian & Alaska Native	385	1.1
Asian	936	2.7
Native Hawaiian & Other Pacific Islander	28	0.1
Two or More Races	458	1.3
Under 5 years	2,177	6.4
Under 18 years	6,925	20.3
65 years and over	3,807	11.2

Source: United States Census Bureau, 2016 Population Estimates

## Health Status Indicators 2012-2016

### Natality

Percent of Low Birth Weight Infants	5.3
○ Percent of Mothers Receiving Care in 1st Trimester	80.7
● Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	10.3
● Percent of Births Less Than 37 Wks. of Gestation	6.9
○ Average Age of Mother	28.6
● Teenage Birth Rate <sup>2</sup>	4.1
Percent White Births	89.1
Percent American Indian Births	2.0
● Percent Unmarried	22.5
● Percent WIC births	23.1
○ Percent Breastfeeding at discharge	86.4
○ Percent Payment-Private Insurance	75.7
● Percent Payment-Medicaid	17.2
● Percent C-Section	20.4

●Denotes a health status indicator which is significantly lower than the state average.  
○Denotes a health status indicator which is significantly higher than the state average.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

All Causes	668.3
Heart Disease	148.6
Malignant Neoplasms (Cancer)	167.0
Trachea, Bronchus, & Lung	35.4
Colon, Rectum, & Anus	18.8
Female Breast	12.6
Pancreas	11.1
Prostate	18.8
Leukemia	10.4
Chronic Lower Respiratory Diseases	32.6
Alzheimer's Disease	33.4
Cerebrovascular Disease	48.9
● Accidents	33.9
● Motor Vehicle Accidents	8.5
Diabetes	21.7
Influenza and Pneumonia	12.3
Intentional Self-Harm (Suicide)	12.1
Chronic Liver Disease and Cirrhosis	9.8
Infant Mortality	8.00

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Malignant Neoplasms (Cancer)	241
2. Heart Disease	225
3. Cerebrovascular Disease	75
T4. Accidents	54
T4. Alzheimer's Disease	54
6. Chronic Lower Respiratory Diseases	48
7. Diabetes	31
8. Influenza and Pneumonia	19
9. Intentional Self-Harm (Suicide)	17
10. Chronic Liver Disease and Cirrhosis	13

Percent of Deaths due to tobacco use	15.1
Median age at death	82

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Brown County

## Demographic Information



Brown County is located on the North Dakota border in the eastern part of the state and averages 21.3 persons per square mile.

### 2016 Population Information

Subject	Number	Percent
Total population	39,128	100.0
White	35,175	89.9
Black or African American	662	1.7
American Indian & Alaska Native	1,351	3.5
Asian	1,126	2.7
Native Hawaiian & Other Pacific Islander	62	0.2
Two or More Races	752	1.9
Under 5 years	2,547	6.5
Under 18 years	8,325	23.8
65 years and over	6,344	16.2

Source: United States Census Bureau, 2016 Population Estimates

## Health Status Indicators 2012-2016

### Nativity

Percent of Low Birth Weight Infants	6.3
Percent of Mothers Receiving Care in 1st Trimester	71.4
○ Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	17.1
Percent of Births Less Than 37 Wks. of Gestation	8.0
Average Age of Mother	27.7
● Teenage Birth Rate <sup>2</sup>	7.5
Percent White Births	87.1
Percent American Indian Births	5.2
● Percent Unmarried	33.6
● Percent WIC births	31.2
Percent Breastfeeding at discharge	81.7
○ Percent Payment-Private Insurance	65.6
● Percent Payment-Medicaid	29.1
Percent C-Section	26.6

●Denotes a health status indicator which is significantly lower than the state average.  
○Denotes a health status indicator which is significantly higher than the state average.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

All Causes	703.5
Heart Disease	147.8
Malignant Neoplasms (Cancer)	165.9
Trachea, Bronchus, & Lung	51.3
Colon, Rectum, & Anus	16.4
Female Breast	27.6
Pancreas	11.6
Prostate	16.3
Leukemia	6.5
Chronic Lower Respiratory Diseases	35.7
Alzheimer's Disease	41.1
Cerebrovascular Disease	40.0
Accidents	40.1
● Motor Vehicle Accidents	9.9
Diabetes	28.2
Influenza and Pneumonia	18.5
Intentional Self-Harm (Suicide)	12.9
● Chronic Liver Disease and Cirrhosis	7.0
Infant Mortality	5.16

### Leading Causes of Death

	Total Deaths
1. Heart Disease	432
2. Malignant Neoplasms (Cancer)	412
3. Alzheimer's Disease	132
4. Cerebrovascular Disease	114
5. Chronic Lower Respiratory Diseases	96
6. Accidents	92
7. Diabetes	72
8. Influenza and Pneumonia	55
9. Pneumonitis Due to Solids and Liquids	32
10. Unspecified Dementia	31

Percent of Deaths due to tobacco use	14.9
Median age at death	83

●Denotes a health status indicator which is significantly lower than the state average.  
○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Brule County

## Demographic Information



Brule County is located in the south central part of the state and averages 6.4 persons per square mile.

## 2016 Population Information

Subject	Number	Percent
Total population	5,238	100.0
White	4,511	86.1
Black or African American	21	0.4
American Indian & Alaska Native	500	9.5
Asian	22	0.3
Native Hawaiian & Other Pacific Islander	4	0.1
Two or More Races	180	3.4
Under 5 years	364	6.9
Under 18 years	1,370	26.2
65 years and over	943	18.0

Source: United States Census Bureau, 2016 Population Estimates

## Health Status Indicators 2012-2016

### Nativity

Percent of Low Birth Weight Infants	5.1
Percent of Mothers Receiving Care in 1st Trimester	65.5
Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	17.7
Percent of Births Less Than 37 Wks. of Gestation	9.2
Average Age of Mother	27.4
● Teenage Birth Rate <sup>2</sup>	4.9
Percent White Births	78.0
Percent American Indian Births	18.2
Percent Unmarried	35.0
Percent WIC births	37.8
Percent Breastfeeding at discharge	70.3
Percent Payment-Private Insurance	63.1
Percent Payment-Medicaid	32.6
Percent C-Section	29.1

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

All Causes	726.6
Heart Disease	165.9
Malignant Neoplasms (Cancer)	171.6
Trachea, Bronchus, & Lung	38.0
Colon, Rectum, & Anus	20.3
Female Breast	20.0
Pancreas	12.3
Prostate	47.9
Leukemia	15.7
Chronic Lower Respiratory Diseases	32.9
Alzheimer's Disease	35.4
Cerebrovascular Disease	21.7
Accidents	52.1
Motor Vehicle Accidents	13.3
Diabetes	25.2
Influenza and Pneumonia	20.5
Intentional Self-Harm (Suicide)	15.7
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	8.09

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	70
2. Malignant Neoplasms (Cancer)	64
3. Accidents	19
4. Alzheimer's Disease	18
5. Chronic Lower Respiratory Diseases	14
6. Diabetes	10
T7. Cerebrovascular Disease	9
T7. Influenza and Pneumonia	9
9. Pneumonitis Due to Solids and Liquids	8

Percent of Deaths due to tobacco use	17.0
Median age at death	81

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Buffalo County

## Demographic Information



Buffalo County is located in the central portion of the state and averages 4.1 persons per square mile.

### 2016 Population Information

Subject	Number	Percent
Total population	2,043	100.0
White	317	15.5
Black or African American	17	0.8
American Indian & Alaska Native	1,658	81.2
Asian	1	0.1
Native Hawaiian & Other Pacific Islander	0	0.0
Two or More Races	50	2.4
Under 5 years	224	11.0
Under 18 years	789	38.6
65 years and over	162	7.9

Source: United States Census Bureau, 2016 Population Estimates

## Health Status Indicators 2012-2016

### Nativity

Percent of Low Birth Weight Infants	8.4
● Percent of Mothers Receiving Care in 1st Trimester	31.7
○ Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	33.6
Percent of Births Less Than 37 Wks. of Gestation	9.6
● Average Age of Mother	25.8
○ Teenage Birth Rate <sup>2</sup>	71.7
Percent White Births	15.9
Percent American Indian Births	81.0
○ Percent Unmarried	72.2
○ Percent WIC births	66.5
● Percent Breastfeeding at discharge	38.8
● Percent Payment-Private Insurance	16.3
○ Percent Payment-Medicaid	71.4
Percent C-Section	22.2

●Denotes a health status indicator which is significantly lower than the state average.  
○Denotes a health status indicator which is significantly higher than the state average.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

○ All Causes	1,486.6
Heart Disease	189.4
Malignant Neoplasms (Cancer)	244.2
Trachea, Bronchus, & Lung	116.5
Colon, Rectum, & Anus	66.1
Female Breast	LNE
Pancreas	LNE
Prostate	LNE
Leukemia	LNE
Chronic Lower Respiratory Diseases	LNE
Alzheimer's Disease	LNE
Cerebrovascular Disease	68.3
Accidents	114.3
Motor Vehicle Accidents	47.0
○ Diabetes	104.3
Influenza and Pneumonia	LNE
○ Intentional Self-Harm (Suicide)	98.6
○ Chronic Liver Disease and Cirrhosis	146.7
Infant Mortality	19.84

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Malignant Neoplasms (Cancer)	16
2. Heart Disease	14
3. Chronic Liver Disease and Cirrhosis	12
4. Accidents	10
5. Intentional Self-Harm (Suicide)	8
6. Diabetes	7
Percent of Deaths due to tobacco use	18.1
Median age at death	61

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Butte County

## Demographic Information



Butte County is located on the western border of the state and averages 4.5 people per square mile.

### 2016 Population Information

Subject	Number	Percent
Total population	10,205	100.0
White	9,611	94.2
Black or African American	61	0.6
American Indian & Alaska Native	236	2.3
Asian	36	0.4
Native Hawaiian & Other Pacific Islander	9	0.1
Two or More Races	252	2.5
Under 5 years	618	6.1
Under 18 years	2,469	24.2
65 years and over	2,004	19.6

Source: United States Census Bureau, 2016 Population Estimates

## Health Status Indicators 2012-2016

### Nativity

Percent of Low Birth Weight Infants	7.5
Percent of Mothers Receiving Care in 1st Trimester	78.1
○ Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	22.2
Percent of Births Less Than 37 Wks. of Gestation	9.3
● Average Age of Mother	26.9
Teenage Birth Rate <sup>2</sup>	11.1
Percent White Births	94.1
Percent American Indian Births	2.7
Percent Unmarried	36.2
○ Percent WIC births	47.3
Percent Breastfeeding at discharge	83.7
Percent Payment-Private Insurance	53.2
○ Percent Payment-Medicaid	40.7
Percent C-Section	23.5

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

All Causes	718.4
○ Heart Disease	197.6
Malignant Neoplasms (Cancer)	148.7
Trachea, Bronchus, & Lung	38.2
Colon, Rectum, & Anus	20.5
Female Breast	8.2
Pancreas	6.8
Prostate	19.4
Leukemia	3.9
Chronic Lower Respiratory Diseases	55.0
Alzheimer's Disease	32.7
Cerebrovascular Disease	34.1
Accidents	59.0
Motor Vehicle Accidents	28.2
Diabetes	15.0
● Influenza and Pneumonia	8.6
Intentional Self-Harm (Suicide)	16.7
Chronic Liver Disease and Cirrhosis	15.4
Infant Mortality	4.76

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	150
2. Malignant Neoplasms (Cancer)	108
3. Chronic Lower Respiratory Diseases	40
4. Accidents	33
5. Alzheimer's Disease	26
6. Cerebrovascular Disease	25
7. Diabetes	12
T8. Intentional Self-Harm (Suicide)	9
T8. Chronic Liver Disease and Cirrhosis	9
T10. Influenza and Pneumonia	6
T10. Septicemia	6
T10. Neoplasms of Uncertain or Unknown Behavior	6

Percent of Deaths due to tobacco use	18.8
Median age at death	79

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Campbell County

## Demographic Information



Campbell County is located in the north central portion of the state and averages 2.0 persons per square mile.

### 2016 Population Information

Subject	Number	Percent
Total population	1,378	100.0
White	1,342	97.4
Black or African American	5	0.4
American Indian & Alaska Native	13	0.9
Asian	8	0.6
Native Hawaiian & Other Pacific Islander	0	0.0
Two or More Races	10	0.7
Under 5 years	55	4.0
Under 18 years	234	17.0
65 years and over	351	25.5

Source: United States Census Bureau, 2016 Population Estimates

## Health Status Indicators 2012-2016

### Nativity

Percent of Low Birth Weight Infants	LNE
Percent of Mothers Receiving Care in 1st Trimester	75.4
● Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	7.0
Percent of Births Less Than 37 Wks. of Gestation	5.4
○ Average Age of Mother	29.2
Teenage Birth Rate <sup>2</sup>	LNE
Percent White Births	94.5
Percent American Indian Births	LNE
● Percent Unmarried	15.8
● Percent WIC births	19.6
Percent Breastfeeding at discharge	83.6
Percent Payment-Private Insurance	74.1
● Percent Payment-Medicaid	14.8
Percent C-Section	35.1

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

All Causes	624.5
Heart Disease	116.7
Malignant Neoplasms (Cancer)	139.8
Trachea, Bronchus, & Lung	39.8
Colon, Rectum, & Anus	LNE
Female Breast	LNE
Pancreas	LNE
Prostate	LNE
Leukemia	29.6
Chronic Lower Respiratory Diseases	LNE
Alzheimer's Disease	19.2
Cerebrovascular Disease	38.9
Accidents	136.2
Motor Vehicle Accidents	66.7
Diabetes	38.0
Influenza and Pneumonia	LNE
Intentional Self-Harm (Suicide)	LNE
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	LNE

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Malignant Neoplasms (Cancer)	19
2. Heart Disease	17
T3. Accidents	6
T3. Cerebrovascular Disease	6
5. Diabetes	5
Percent of Deaths due to tobacco use	20.0
Median age at death	77

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Charles Mix County

## Demographic Information



Charles Mix County is located in the south central area of the state and averages 8.3 persons per square mile.

### 2016 Population Information

Subject	Number	Percent
Total population	9,396	100.0
White	6,027	64.1
Black or African American	41	0.4
American Indian & Alaska Native	2,994	31.9
Asian	29	0.3
Native Hawaiian & Other Pacific Islander	1	0.0
Two or More Races	304	3.2
Under 5 years	858	9.1
Under 18 years	2,794	29.7
65 years and over	1,668	17.8

Source: United States Census Bureau, 2016 Population Estimates

## Health Status Indicators 2012-2016

### Nativity

Percent of Low Birth Weight Infants	6.2
● Percent of Mothers Receiving Care in 1st Trimester	65.0
○ Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	23.0
Percent of Births Less Than 37 Wks. of Gestation	8.2
● Average Age of Mother	26.3
○ Teenage Birth Rate <sup>2</sup>	24.8
Percent White Births	41.8
Percent American Indian Births	52.5
○ Percent Unmarried	57.7
○ Percent WIC births	60.2
● Percent Breastfeeding at discharge	66.9
● Percent Payment-Private Insurance	35.6
○ Percent Payment-Medicaid	54.4
○ Percent C-Section	30.8

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

○ All Causes	807.8
Heart Disease	173.3
Malignant Neoplasms (Cancer)	169.3
Trachea, Bronchus, & Lung	42.3
Colon, Rectum, & Anus	10.6
Female Breast	34.2
Pancreas	11.7
Prostate	26.1
Leukemia	5.9
Chronic Lower Respiratory Diseases	37.0
○ Alzheimer's Disease	55.8
Cerebrovascular Disease	37.5
Accidents	70.5
Motor Vehicle Accidents	15.2
Diabetes	35.7
Influenza and Pneumonia	18.0
○ Intentional Self-Harm (Suicide)	42.6
Chronic Liver Disease and Cirrhosis	31.1
Infant Mortality	LNE

### Leading Causes of Death

	Total Deaths
1. Heart Disease	114
2. Malignant Neoplasms (Cancer)	112
3. Alzheimer's Disease	45
4. Accidents	36
5. Cerebrovascular Disease	26
6. Chronic Lower Respiratory Diseases	25
7. Diabetes	19
8. Intentional Self-Harm (Suicide)	16
9. Influenza and Pneumonia	14
T10. Chronic Liver Disease and Cirrhosis	12
T10. Assault (Homicide)	12

Percent of Deaths due to tobacco use	15.3
Median age at death	78

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Clark County

## Demographic Information



Clark County is located in east central South Dakota and averages 3.9 persons per square mile.

### 2016 Population Information

Subject	Number	Percent
Total population	3,656	100.0
White	3,519	96.3
Black or African American	74	2.0
American Indian & Alaska Native	8	0.2
Asian	20	0.5
Native Hawaiian & Other Pacific Islander	0	0.0
Two or More Races	35	1.0
Under 5 years	349	9.5
Under 18 years	910	24.9
65 years and over	751	20.5

Source: United States Census Bureau, 2016 Population Estimates

## Health Status Indicators 2012-2016

### Nativity

Percent of Low Birth Weight Infants	5.2
Percent of Mothers Receiving Care in 1st Trimester	64.4
● Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	9.4
Percent of Births Less Than 37 Wks. of Gestation	6.3
Average Age of Mother	28.3
Teenage Birth Rate <sup>2</sup>	9.4
Percent White Births	98.2
Percent American Indian Births	1.4
● Percent Unmarried	16.0
● Percent WIC births	19.9
Percent Breastfeeding at discharge	87.8
○ Percent Payment-Private Insurance	82.6
● Percent Payment-Medicaid	13.9
Percent C-Section	20.9

●Denotes a health status indicator which is significantly lower than the state average.  
○Denotes a health status indicator which is significantly higher than the state average.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

All Causes	679.8
Heart Disease	131.3
Malignant Neoplasms (Cancer)	130.0
Trachea, Bronchus, & Lung	29.9
Colon, Rectum, & Anus	10.7
Female Breast	17.5
Pancreas	LNE
Prostate	20.9
Leukemia	13.8
Chronic Lower Respiratory Diseases	37.3
Alzheimer's Disease	27.9
Cerebrovascular Disease	50.0
Accidents	60.7
Motor Vehicle Accidents	18.4
Diabetes	30.0
Influenza and Pneumonia	20.4
Intentional Self-Harm (Suicide)	16.7
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	10.45

### Leading Causes of Death

Leading Causes of Death	Total Deaths
T1. Heart Disease	43
T1. Malignant Neoplasms (Cancer)	43
T3. Accidents	18
T3. Cerebrovascular Disease	18
5. Alzheimer's Disease	12
T6. Diabetes	8
T6. Influenza and Pneumonia	8
8. Unspecified Dementia	5
Percent of Deaths due to tobacco use	13.9
Median age at death	84

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Clay County

## Demographic Information



Clay County is located in the southeastern part of the state and averages 33.6 persons per square mile.

### 2016 Population Information

Subject	Number	Percent
Total population	14,086	100.0
White	12,697	90.1
Black or African American	250	1.8
American Indian & Alaska Native	451	3.2
Asian	330	2.3
Native Hawaiian & Other Pacific Islander	15	0.1
Two or More Races	343	2.4
Under 5 years	708	5.0
Under 18 years	2,451	17.4
65 years and over	1,660	11.8

Source: United States Census Bureau, 2016 Population Estimates

## Health Status Indicators 2012-2016

### Nativity

Percent of Low Birth Weight Infants	5.6
○ Percent of Mothers Receiving Care in 1st Trimester	80.4
● Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	10.0
Percent of Births Less Than 37 Wks. of Gestation	7.3
○ Average Age of Mother	28.6
● Teenage Birth Rate <sup>2</sup>	3.9
Percent White Births	86.1
Percent American Indian Births	5.9
● Percent Unmarried	28.9
● Percent WIC births	30.2
Percent Breastfeeding at discharge	80.7
○ Percent Payment-Private Insurance	68.0
Percent Payment-Medicaid	28.7
○ Percent C-Section	33.5

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

All Causes	777.3
○ Heart Disease	240.8
Malignant Neoplasms (Cancer)	178.4
Trachea, Bronchus, & Lung	50.8
Colon, Rectum, & Anus	14.0
Female Breast	17.7
Pancreas	9.0
Prostate	39.4
Leukemia	9.9
Chronic Lower Respiratory Diseases	41.8
● Alzheimer's Disease	20.4
Cerebrovascular Disease	44.4
● Accidents	24.0
Motor Vehicle Accidents	9.3
Diabetes	28.0
Influenza and Pneumonia	24.5
Intentional Self-Harm (Suicide)	15.4
Chronic Liver Disease and Cirrhosis	8.2
Infant Mortality	6.65

### Leading Causes of Death

	Total Deaths
1. Heart Disease	168
2. Malignant Neoplasms (Cancer)	112
3. Cerebrovascular Disease	30
4. Chronic Lower Respiratory Diseases	27
5. Diabetes	18
6. Influenza and Pneumonia	17
T7. Accidents	15
T7. Alzheimer's Disease	15
9. Intentional Self-Harm (Suicide)	9
T10. Nephritis, nephrotic syndrome, and Nephrosis	6
T10. Congenital Malformations, Deformations, and Chromosomal Abnormalities	6
Percent of Deaths due to tobacco use	16.3
Median age at death	81

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Codington County

## Demographic Information



Codington County is located in eastern South Dakota and averages 39.5 persons per square mile.

### 2016 Population Information

Subject	Number	Percent
Total population	28,063	100.0
White	26,494	94.4
Black or African American	214	0.8
American Indian & Alaska Native	671	2.4
Asian	280	1.0
Native Hawaiian & Other Pacific Islander	2	0.0
Two or More Races	402	1.4
Under 5 years	1,869	6.7
Under 18 years	6,866	24.5
65 years and over	4,572	16.3

Source: United States Census Bureau, 2016 Population Estimates

## Health Status Indicators 2012-2016

### Natality

Percent of Low Birth Weight Infants	6.7
○ Percent of Mothers Receiving Care in 1st Trimester	79.4
○ Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	21.9
Percent of Births Less Than 37 Wks. of Gestation	7.2
● Average Age of Mother	27.3
Teenage Birth Rate <sup>2</sup>	9.0
Percent White Births	90.4
Percent American Indian Births	5.1
Percent Unmarried	37.9
Percent WIC births	36.7
Percent Breastfeeding at discharge	77.1
○ Percent Payment-Private Insurance	65.3
Percent Payment-Medicaid	31.5
Percent C-Section	23.8

### Mortality<sup>3</sup>

● All Causes	651.8
Heart Disease	144.2
○ Malignant Neoplasms (Cancer)	185.1
Trachea, Bronchus, & Lung	49.7
Colon, Rectum, & Anus	20.9
Female Breast	26.2
Pancreas	13.1
Prostate	21.3
Leukemia	10.7
Chronic Lower Respiratory Diseases	43.0
Alzheimer's Disease	31.7
Cerebrovascular Disease	32.2
● Accidents	33.4
Motor Vehicle Accidents	13.2
Diabetes	24.4
Influenza and Pneumonia	12.6
Intentional Self-Harm (Suicide)	12.9
● Chronic Liver Disease and Cirrhosis	5.7
Infant Mortality	3.67

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Malignant Neoplasms (Cancer)	328
2. Heart Disease	277
3. Chronic Lower Respiratory Diseases	78
4. Alzheimer's Disease	64
5. Cerebrovascular Disease	62
6. Accidents	52
7. Diabetes	45
8. Influenza and Pneumonia	24
9. Unspecified Dementia	19
10. Intentional Self-Harm (Suicide)	17
Percent of Deaths due to tobacco use	21.0
Median age at death	81

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Corson County

## Demographic Information



Corson County is located in the north central portion of the state and averages 1.6 persons per square mile.

### 2016 Population Information

Subject	Number	Percent
Total population	4,132	100.0
White	1,298	31.4
Black or African American	13	0.3
American Indian & Alaska Native	2,652	64.2
Asian	22	0.5
Native Hawaiian & Other Pacific Islander	2	0.1
Two or More Races	145	3.5
Under 5 years	403	9.8
Under 18 years	1,462	35.4
65 years and over	459	11.1

Source: United States Census Bureau, 2016 Population Estimates

## Health Status Indicators 2012-2016

### Nativity

Percent of Low Birth Weight Infants	8.4
● Percent of Mothers Receiving Care in 1st Trimester	43.8
○ Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	23.3
○ Percent of Births Less Than 37 Wks. of Gestation	13.0
● Average Age of Mother	26.4
○ Teenage Birth Rate <sup>2</sup>	29.2
Percent White Births	20.9
Percent American Indian Births	73.6
○ Percent Unmarried	70.5
○ Percent WIC births	69.1
● Percent Breastfeeding at discharge	55.0
● Percent Payment-Private Insurance	17.6
○ Percent Payment-Medicaid	74.3
Percent C-Section	28.3

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

○ All Causes	1,236.5
○ Heart Disease	303.7
Malignant Neoplasms (Cancer)	201.5
Trachea, Bronchus, & Lung	24.9
Colon, Rectum, & Anus	44.7
Female Breast	LNE
Pancreas	LNE
Prostate	LNE
Leukemia	LNE
Chronic Lower Respiratory Diseases	38.7
Alzheimer's Disease	19.8
Cerebrovascular Disease	28.9
○ Accidents	117.5
Motor Vehicle Accidents	26.9
○ Diabetes	73.8
Influenza and Pneumonia	18.0
○ Intentional Self-Harm (Suicide)	60.2
○ Chronic Liver Disease and Cirrhosis	63.4
Infant Mortality	15.35

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	56
2. Malignant Neoplasms (Cancer)	39
3. Accidents	23
T4. Diabetes	13
T4. Chronic Liver Disease and Cirrhosis	13
6. Intentional Self-Harm (Suicide)	12
7. Chronic Lower Respiratory Diseases	7
T8. Nephritis, Nephrotic Syndrome, and Nephrosis	6
T8. Ill-Defined and Unknown Causes of Mortality	6
10. Cerebrovascular Disease	5
Percent of Deaths due to tobacco use	16.5
Median age at death	62

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Custer County

## Demographic Information



Custer County is located in the southern Black Hills and averages 5.3 persons per square mile.

## 2016 Population Information

Subject	Number	Percent
Total population	8,596	100.0
White	7,986	92.9
Black or African American	52	0.6
American Indian & Alaska Native	322	3.7
Asian	43	0.5
Native Hawaiian & Other Pacific Islander	2	0.0
Two or More Races	191	2.2
Under 5 years	327	3.8
Under 18 years	1,371	15.9
65 years and over	2,328	27.1

Source: United States Census Bureau, 2016 Population Estimates

## Health Status Indicators 2012-2016

### Nativity

Percent of Low Birth Weight Infants	8.7
Percent of Mothers Receiving Care in 1st Trimester	70.1
Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	17.7
Percent of Births Less Than 37 Wks. of Gestation	6.7
Average Age of Mother	28.2
Teenage Birth Rate <sup>2</sup>	7.0
Percent White Births	87.5
Percent American Indian Births	8.0
Percent Unmarried	36.5
Percent WIC births	33.5
Percent Breastfeeding at discharge	85.7
Percent Payment-Private Insurance	52.0
Percent Payment-Medicaid	37.6
Percent C-Section	21.3

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

All Causes	644.6
Heart Disease	127.2
Malignant Neoplasms (Cancer)	140.1
Trachea, Bronchus, & Lung	30.2
Colon, Rectum, & Anus	14.4
Female Breast	20.7
Pancreas	7.9
Prostate	14.8
Leukemia	11.4
Chronic Lower Respiratory Diseases	36.3
● Alzheimer's Disease	21.9
Cerebrovascular Disease	38.3
Accidents	62.0
Motor Vehicle Accidents	25.3
Diabetes	13.8
Influenza and Pneumonia	16.9
Intentional Self-Harm (Suicide)	19.5
Chronic Liver Disease and Cirrhosis	13.5
Infant Mortality	LNE

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Malignant Neoplasms (Cancer)	110
2. Heart Disease	97
3. Accidents	29
T4. Chronic Lower Respiratory Diseases	27
T4. Cerebrovascular Disease	27
6. Alzheimer's Disease	16
7. Influenza and Pneumonia	12
8. Intentional Self-Harm (Suicide)	11
T9. Diabetes	10
T9. Chronic Liver Disease and Cirrhosis	10

Percent of Deaths due to tobacco use	17.0
Median age at death	78

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Davison County

## Demographic Information



Davison County is located in eastern South Dakota and averages 44.8 persons per square mile.

### 2016 Population Information

Subject	Number	Percent
Total population	19,903	100.0
White	18,653	93.7
Black or African American	171	0.9
American Indian & Alaska Native	605	3.0
Asian	133	0.7
Native Hawaiian & Other Pacific Islander	23	0.1
Two or More Races	318	1.6
Under 5 years	1,271	6.4
Under 18 years	4,632	23.3
65 years and over	3,582	18.0

Source: United States Census Bureau, 2016 Population Estimates

## Health Status Indicators 2012-2016

### Nativity

Percent of Low Birth Weight Infants	5.1
○ Percent of Mothers Receiving Care in 1st Trimester	80.7
○ Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	19.0
Percent of Births Less Than 37 Wks. of Gestation	8.6
● Average Age of Mother	27.0
Teenage Birth Rate <sup>2</sup>	11.4
Percent White Births	87.9
Percent American Indian Births	8.5
Percent Unmarried	40.7
Percent WIC births	36.6
Percent Breastfeeding at discharge	73.3
○ Percent Payment-Private Insurance	64.7
Percent Payment-Medicaid	30.8
Percent C-Section	28.5

●Denotes a health status indicator which is significantly lower than the state average.  
○Denotes a health status indicator which is significantly higher than the state average.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

All Causes	702.4
○ Heart Disease	187.0
Malignant Neoplasms (Cancer)	166.0
Trachea, Bronchus, & Lung	52.6
Colon, Rectum, & Anus	16.1
Female Breast	13.4
Pancreas	10.3
Prostate	9.9
Leukemia	8.5
● Chronic Lower Respiratory Diseases	28.1
● Alzheimer's Disease	25.7
Cerebrovascular Disease	35.2
Accidents	40.8
● Motor Vehicle Accidents	8.8
Diabetes	24.8
Influenza and Pneumonia	24.9
Intentional Self-Harm (Suicide)	17.7
Chronic Liver Disease and Cirrhosis	8.7
Infant Mortality	5.40

### Leading Causes of Death

	Total Deaths
1. Heart Disease	285
2. Malignant Neoplasms (Cancer)	239
3. Cerebrovascular Disease	65
4. Alzheimer's Disease	50
5. Accidents	49
6. Influenza and Pneumonia	46
7. Chronic Lower Respiratory Diseases	45
8. Unspecified Dementia	34
9. Diabetes	31
10. Essential (Primary) Hypertension and Hypertensive Renal Disease	18
Percent of Deaths due to tobacco use	15.9
Median age at death	82

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Day County

## Demographic Information



Day County is located in the northeastern part of the state and averages 5.6 persons per square mile.

## 2016 Population Information

Subject	Number	Percent
Total population	5,571	100.0
White	4,877	87.5
Black or African American	25	0.4
American Indian & Alaska Native	540	9.7
Asian	27	0.5
Native Hawaiian & Other Pacific Islander	0	0.0
Two or More Races	102	1.8
Under 5 years	329	5.9
Under 18 years	1,248	22.4
65 years and over	1,408	25.3

Source: United States Census Bureau, 2016 Population Estimates

## Health Status Indicators 2012-2016

### Natality

Percent of Low Birth Weight Infants	6.8
Percent of Mothers Receiving Care in 1st Trimester	71.6
○ Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	26.1
Percent of Births Less Than 37 Wks. of Gestation	8.5
Average Age of Mother	27.7
Teenage Birth Rate <sup>2</sup>	14.5
Percent White Births	76.2
Percent American Indian Births	18.6
Percent Unmarried	35.3
Percent WIC births	40.5
Percent Breastfeeding at discharge	73.9
Percent Payment-Private Insurance	63.5
Percent Payment-Medicaid	31.6
Percent C-Section	27.0

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

All Causes	660.5
Heart Disease	160.0
Malignant Neoplasms (Cancer)	128.5
Trachea, Bronchus, & Lung	32.1
● Colon, Rectum, & Anus	6.6
Female Breast	21.7
Pancreas	12.3
Prostate	28.3
Leukemia	LNE
Chronic Lower Respiratory Diseases	34.2
Alzheimer's Disease	35.8
Cerebrovascular Disease	41.7
Accidents	71.6
Motor Vehicle Accidents	25.9
Diabetes	18.3
Influenza and Pneumonia	11.5
Intentional Self-Harm (Suicide)	14.8
● Chronic Liver Disease and Cirrhosis	6.2
Infant Mortality	LNE

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	96
2. Malignant Neoplasms (Cancer)	63
3. Cerebrovascular Disease	28
T4. Accidents	25
T4. Alzheimer's Disease	25
6. Chronic Lower Respiratory Diseases	20
7. Diabetes	10
8. Parkinson's Disease	9
T9. Influenza and Pneumonia	7
T9. Essential (Primary) Hypertension and Hypertensive Renal Disease	7

Percent of Deaths due to tobacco use	15.0
Median age at death	83

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Deuel County

## Demographic Information



Deuel County borders Minnesota and averages 7.0 persons per square mile.

## 2016 Population Information

Subject	Number	Percent
Total population	4,231	100.0
White	4,131	97.6
Black or African American	34	0.8
American Indian & Alaska Native	19	0.4
Asian	5	0.1
Native Hawaiian & Other Pacific Islander	0	0.0
Two or More Races	42	1.0
Under 5 years	232	5.5
Under 18 years	978	23.1
65 years and over	915	21.6

Source: United States Census Bureau, 2016 Population Estimates

## Health Status Indicators 2012-2016

### Nativity

Percent of Low Birth Weight Infants	4.5
Percent of Mothers Receiving Care in 1st Trimester	80.6
Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	12.2
Percent of Births Less Than 37 Wks. of Gestation	5.3
Average Age of Mother	27.9
Teenage Birth Rate <sup>2</sup>	LNE
Percent White Births	100
Percent American Indian Births	LNE
● Percent Unmarried	20.6
Percent WIC births	28.9
Percent Breastfeeding at discharge	85.0
○ Percent Payment-Private Insurance	71.4
● Percent Payment-Medicaid	19.6
Percent C-Section	22.7

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

● All Causes	586.6
Heart Disease	148.0
Malignant Neoplasms (Cancer)	124.4
Trachea, Bronchus, & Lung	28.9
Colon, Rectum, & Anus	8.0
Female Breast	LNE
Pancreas	LNE
Prostate	LNE
Leukemia	LNE
Chronic Lower Respiratory Diseases	34.3
● Alzheimer's Disease	18.3
Cerebrovascular Disease	40.7
Accidents	42.6
Motor Vehicle Accidents	14.6
● Diabetes	10.9
Influenza and Pneumonia	23.8
Intentional Self-Harm (Suicide)	LNE
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	LNE

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	53
2. Malignant Neoplasms (Cancer)	44
3. Cerebrovascular Disease	16
4. Chronic Lower Respiratory Diseases	13
5. Accidents	12
6. Influenza and Pneumonia	11
T7. Alzheimer's Disease	7
T7. Septicemia	7
Percent of Deaths due to tobacco use	16.7
Median age at death	83

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Dewey County

## Demographic Information



Dewey County is located in the north central region of the state and averages 2.3 persons per square mile.

### 2016 Population Information

Subject	Number	Percent
Total population	5,742	100.0
White	1,234	21.5
Black or African American	25	0.4
American Indian & Alaska Native	4,205	73.2
Asian	15	0.3
Native Hawaiian & Other Pacific Islander	2	0.0
Two or More Races	261	4.5
Under 5 years	705	12.3
Under 18 years	2,104	36.6
65 years and over	553	9.6

Source: United States Census Bureau, 2016 Population Estimates

## Health Status Indicators 2012-2016

### Nativity

Percent of Low Birth Weight Infants	8.0
● Percent of Mothers Receiving Care in 1st Trimester	49.4
○ Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	21.5
○ Percent of Births Less Than 37 Wks. of Gestation	13.2
● Average Age of Mother	25.7
○ Teenage Birth Rate <sup>2</sup>	47.1
Percent White Births	10.6
Percent American Indian Births	84.3
○ Percent Unmarried	69.3
○ Percent WIC births	79.6
● Percent Breastfeeding at discharge	54.2
● Percent Payment-Private Insurance	16.8
○ Percent Payment-Medicaid	72.6
○ Percent C-Section	33.4

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

○ All Causes	1,376.1
○ Heart Disease	238.0
○ Malignant Neoplasms (Cancer)	222.1
Trachea, Bronchus, & Lung	60.5
Colon, Rectum, & Anus	25.8
Female Breast	26.1
Pancreas	16.3
Prostate	31.6
Leukemia	12.0
Chronic Lower Respiratory Diseases	52.2
Alzheimer's Disease	LNE
Cerebrovascular Disease	28.2
○ Accidents	162.9
○ Motor Vehicle Accidents	79.6
○ Diabetes	91.9
○ Influenza and Pneumonia	54.1
○ Intentional Self-Harm (Suicide)	37.2
○ Chronic Liver Disease and Cirrhosis	90.6
Infant Mortality	11.80

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	54
2. Malignant Neoplasms (Cancer)	52
3. Accidents	42
4. Chronic Liver Disease and Cirrhosis	21
5. Diabetes	20
6. Influenza and Pneumonia	13
7. Chronic Lower Respiratory Diseases	12
8. Intentional Self-Harm (Suicide)	10
9. Nephritis, Nephrotic Syndrome, and Nephrosis	9
10. Cerebrovascular Disease	7
Percent of Deaths due to tobacco use	16.3
Median age at death	65

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Douglas County

## Demographic Information



Douglas County is located in south central South Dakota and averages 7.0 persons per square mile.

### 2016 Population Information

Subject	Number	Percent
Total population	2,932	100.0
White	2,805	95.7
Black or African American	16	0.5
American Indian & Alaska Native	71	2.4
Asian	7	0.2
Native Hawaiian & Other Pacific Islander	0	0.0
Two or More Races	33	1.1
Under 5 years	197	6.7
Under 18 years	672	22.9
65 years and over	749	25.5

Source: United States Census Bureau, 2016 Population Estimates

## Health Status Indicators 2012-2016

### Nativity

Percent of Low Birth Weight Infants	5.5
Percent of Mothers Receiving Care in 1st Trimester	74.4
• Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	9.0
Percent of Births Less Than 37 Wks. of Gestation	5.5
Average Age of Mother	27.4
Teenage Birth Rate <sup>2</sup>	LNE
Percent White Births	98.0
Percent American Indian Births	LNE
• Percent Unmarried	13.1
• Percent WIC births	23.7
Percent Breastfeeding at discharge	89.8
Percent Payment-Private Insurance	69.9
• Percent Payment-Medicaid	17.9
Percent C-Section	29.1

•Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

All Causes	625.9
Heart Disease	121.4
Malignant Neoplasms (Cancer)	157.1
Trachea, Bronchus, & Lung	37.5
Colon, Rectum, & Anus	21.5
Female Breast	54.5
Pancreas	13.0
Prostate	LNE
Leukemia	LNE
Chronic Lower Respiratory Diseases	50.6
Alzheimer's Disease	48.4
Cerebrovascular Disease	25.5
Accidents	45.2
Motor Vehicle Accidents	LNE
Diabetes	14.8
Influenza and Pneumonia	12.5
Intentional Self-Harm (Suicide)	LNE
Chronic Liver Disease and Cirrhosis	28.6
Infant Mortality	LNE

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	47
2. Malignant Neoplasms (Cancer)	44
3. Alzheimer's Disease	20
4. Chronic Lower Respiratory Diseases	16
5. Accidents	15
6. Cerebrovascular Disease	11
T7. Diabetes	6
T7. Influenza and Pneumonia	6
9. Essential (Primary) Hypertension and Hypertensive Renal Disease	5
Percent of Deaths due to tobacco use	15.2
Median age at death	86

•Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Edmunds County

## Demographic Information



Edmunds County is located in the north central region of the state and averages 3.6 persons per square mile.

### 2016 Population Information

Subject	Number	Percent
Total population	3,952	100.0
White	3,842	97.2
Black or African American	17	0.4
American Indian & Alaska Native	34	0.9
Asian	15	0.4
Native Hawaiian & Other Pacific Islander	0	0.0
Two or More Races	44	1.1
Under 5 years	282	57.1
Under 18 years	917	23.2
65 years and over	851	21.5

Source: United States Census Bureau, 2016 Population Estimates

## Health Status Indicators 2012-2016

### Nativity

● Percent of Low Birth Weight Infants	2.1
○ Percent of Mothers Receiving Care in 1st Trimester	66.8
● Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	8.8
● Percent of Births Less Than 37 Wks. of Gestation	4.6
Average Age of Mother	28.3
Teenage Birth Rate <sup>2</sup>	LNE
Percent White Births	97.5
Percent American Indian Births	1.3
● Percent Unmarried	15.9
● Percent WIC births	18.8
Percent Breastfeeding at discharge	86.6
○ Percent Payment-Private Insurance	79.5
● Percent Payment-Medicaid	15.9
● Percent C-Section	19.2

●Denotes a health status indicator which is significantly lower than the state average.  
○Denotes a health status indicator which is significantly higher than the state average.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

● All Causes	607.7
Heart Disease	120.7
Malignant Neoplasms (Cancer)	130.3
Trachea, Bronchus, & Lung	40.1
Colon, Rectum, & Anus	10.0
Female Breast	LNE
Pancreas	16.3
Prostate	23.8
Leukemia	LNE
Chronic Lower Respiratory Diseases	32.9
Alzheimer's Disease	24.8
Cerebrovascular Disease	28.8
Accidents	52.4
Motor Vehicle Accidents	21.7
Diabetes	31.8
Influenza and Pneumonia	24.3
Intentional Self-Harm (Suicide)	LNE
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	LNE

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	45
2. Malignant Neoplasms (Cancer)	43
3. Accidents	15
4. Chronic Lower Respiratory Diseases	13
T5. Alzheimer's Disease	11
T5. Cerebrovascular Disease	11
T5. Influenza and Pneumonia	11
8. Essential (Primary) Hypertension and Hypertensive Renal Disease	10
9. Diabetes	9
10. Unspecified Dementia	7
Percent of Deaths due to tobacco use	17.2
Median age at death	84

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Fall River County

## Demographic Information



Fall River County is located in the southwestern corner of the state and averages 4.1 persons per square mile.

### 2016 Population Information

Subject	Number	Percent
Total population	6,849	100.0
White	6,002	87.6
Black or African American	93	1.4
American Indian & Alaska Native	471	6.9
Asian	53	0.8
Native Hawaiian & Other Pacific Islander	5	0.1
Two or More Races	225	3.3
Under 5 years	313	4.6
Under 18 years	1,226	17.9
65 years and over	1,899	27.7

Source: United States Census Bureau, 2016 Population Estimates

## Health Status Indicators 2012-2016

### Nativity

Percent of Low Birth Weight Infants	7.9
Percent of Mothers Receiving Care in 1st Trimester	67.6
Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	17.6
Percent of Births Less Than 37 Wks. of Gestation	8.0
Average Age of Mother	27.1
Teenage Birth Rate <sup>2</sup>	12.2
Percent White Births	79.8
Percent American Indian Births	10.3
Percent Unmarried	43.5
○ Percent WIC births	45.5
Percent Breastfeeding at discharge	79.0
● Percent Payment-Private Insurance	47.1
○ Percent Payment-Medicaid	40.4
● Percent C-Section	17.1

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

○ All Causes	957.3
○ Heart Disease	212.5
Malignant Neoplasms (Cancer)	186.4
Trachea, Bronchus, & Lung	39.8
Colon, Rectum, & Anus	27.1
Female Breast	28.4
Pancreas	5.6
Prostate	13.5
Leukemia	5.6
○ Chronic Lower Respiratory Diseases	93.7
Alzheimer's Disease	30.7
Cerebrovascular Disease	37.7
Accidents	74.9
Motor Vehicle Accidents	26.1
○ Diabetes	41.5
Influenza and Pneumonia	17.8
Intentional Self-Harm (Suicide)	23.1
Chronic Liver Disease and Cirrhosis	20.9
Infant Mortality	LNE

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	139
2. Malignant Neoplasms (Cancer)	119
3. Chronic Lower Respiratory Diseases	66
4. Accidents	32
T5. Cerebrovascular Disease	25
T5. Diabetes	25
7. Alzheimer's Disease	22
T8. Influenza and Pneumonia	10
T8. Intentional Self-Harm (Suicide)	10
T8. Unspecified Dementia	10
Percent of Deaths due to tobacco use	23.3
Median age at death	79

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Faulk County

## Demographic Information



Faulk County is located in north central South Dakota and averages 2.4 persons per square mile.

### 2016 Population Information

Subject	Number	Percent
Total population	2,354	100.0
White	2,312	98.2
Black or African American	9	0.4
American Indian & Alaska Native	7	0.3
Asian	4	0.2
Native Hawaiian & Other Pacific Islander	0	0.0
Two or More Races	22	0.9
Under 5 years	188	8.0
Under 18 years	555	23.6
65 years and over	547	23.2

Source: United States Census Bureau, 2016 Population Estimates

## Health Status Indicators 2012-2016

### Nativity

Percent of Low Birth Weight Infants	7.9
● Percent of Mothers Receiving Care in 1st Trimester	58.6
● Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	2.0
Percent of Births Less Than 37 Wks. of Gestation	5.9
○ Average Age of Mother	29.2
Teenage Birth Rate <sup>2</sup>	LNE
Percent White Births	100
Percent American Indian Births	LNE
● Percent Unmarried	5.3
● Percent WIC births	13.2
Percent Breastfeeding at discharge	90.1
○ Percent Payment-Private Insurance	84.9
● Percent Payment-Medicaid	7.2
Percent C-Section	18.4

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

All Causes	690.3
Heart Disease	138.5
Malignant Neoplasms (Cancer)	173.3
Trachea, Bronchus, & Lung	34.4
Colon, Rectum, & Anus	LNE
Female Breast	35.4
Pancreas	15.8
Prostate	26.4
Leukemia	LNE
Chronic Lower Respiratory Diseases	44.4
Alzheimer's Disease	38.6
● Cerebrovascular Disease	17.3
Accidents	33.9
Motor Vehicle Accidents	LNE
Diabetes	36.9
Influenza and Pneumonia	LNE
Intentional Self-Harm (Suicide)	25.8
Chronic Liver Disease and Cirrhosis	29.1
Infant Mortality	LNE

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Malignant Neoplasms (Cancer)	37
2. Heart Disease	36
3. Alzheimer's Disease	13
4. Chronic Lower Respiratory Diseases	11
T5. Accidents	8
T5. Diabetes	8
T7. Cerebrovascular Disease	5
T7. Essential (Primary) Hypertension and Hypertensive Renal Disease	5
Percent of Deaths due to tobacco use	11.9
Median age at death	83

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Grant County

## Demographic Information



Grant County borders Minnesota in northeastern South Dakota and averages 10.8 persons per square mile.

### 2016 Population Information

Subject	Number	Percent
Total population	7,148	100.0
White	6,905	96.6
Black or African American	46	0.6
American Indian & Alaska Native	73	1.0
Asian	23	0.3
Native Hawaiian & Other Pacific Islander	3	0.0
Two or More Races	98	1.4
Under 5 years	441	6.2
Under 18 years	1,621	22.7
65 years and over	1,511	21.1

Source: United States Census Bureau, 2016 Population Estimates

## Health Status Indicators 2012-2016

### Natality

Percent of Low Birth Weight Infants	5.0
Percent of Mothers Receiving Care in 1st Trimester	78.9
Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	12.9
Percent of Births Less Than 37 Wks. of Gestation	6.3
Average Age of Mother	27.9
● Teenage Birth Rate <sup>2</sup>	5.2
Percent White Births	95.1
Percent American Indian Births	1.2
● Percent Unmarried	27.8
Percent WIC births	32.7
Percent Breastfeeding at discharge	82.3
○ Percent Payment-Private Insurance	70.1
● Percent Payment-Medicaid	19.9
Percent C-Section	25.8

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

All Causes	726.6
Heart Disease	160.7
Malignant Neoplasms (Cancer)	127.5
Trachea, Bronchus, & Lung	39.1
● Colon, Rectum, & Anus	7.8
Female Breast	LNE
Pancreas	10.4
Prostate	LNE
Leukemia	LNE
Chronic Lower Respiratory Diseases	60.0
Alzheimer's Disease	43.2
○ Cerebrovascular Disease	63.2
Accidents	38.1
Motor Vehicle Accidents	14.1
Diabetes	30.3
Influenza and Pneumonia	21.9
Intentional Self-Harm (Suicide)	13.2
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	LNE

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	101
2. Malignant Neoplasms (Cancer)	72
3. Cerebrovascular Disease	42
4. Chronic Lower Respiratory Diseases	37
5. Alzheimer's Disease	29
6. Accidents	19
7. Diabetes	18
8. Influenza and Pneumonia	13
T9. Essential (Primary) Hypertension and Hypertensive Renal Disease	7
T9. Nephritis, Nephrotic Syndrome, and Nephrosis	7
Percent of Deaths due to tobacco use	16.7
Median age at death	83

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Gregory County

## Demographic Information



Gregory County borders the west bank of the Missouri River and the state of Nebraska and averages 4.2 persons per square mile.

### 2016 Population Information

Subject	Number	Percent
Total population	4,171	100.0
White	3,698	88.6
Black or African American	12	0.3
American Indian & Alaska Native	325	7.8
Asian	17	0.4
Native Hawaiian & Other Pacific Islander	3	0.1
Two or More Races	116	2.8
Under 5 years	259	6.2
Under 18 years	977	23.4
65 years and over	1,019	24.4

Source: United States Census Bureau, 2016 Population Estimates

## Health Status Indicators 2012-2016

### Natality

● Percent of Low Birth Weight Infants	3.5
● Percent of Mothers Receiving Care in 1st Trimester	60.8
Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	12.1
Percent of Births Less Than 37 Wks. of Gestation	5.8
Average Age of Mother	27.9
Teenage Birth Rate <sup>2</sup>	18.3
Percent White Births	77.4
Percent American Indian Births	17.1
Percent Unmarried	34.7
Percent WIC births	41.7
Percent Breastfeeding at discharge	79.1
Percent Payment-Private Insurance	57.5
Percent Payment-Medicaid	37.5
Percent C-Section	27.0

●Denotes a health status indicator which is significantly lower than the state average.  
○Denotes a health status indicator which is significantly higher than the state average.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

○ All Causes	808.6
Heart Disease	179.3
Malignant Neoplasms (Cancer)	169.6
Trachea, Bronchus, & Lung	38.0
Colon, Rectum, & Anus	13.8
Female Breast	20.4
Pancreas	20.7
Prostate	LNE
Leukemia	LNE
Chronic Lower Respiratory Diseases	58.4
Alzheimer's Disease	25.0
Cerebrovascular Disease	35.4
○ Accidents	92.5
○ Motor Vehicle Accidents	60.5
Diabetes	34.5
Influenza and Pneumonia	8.8
Intentional Self-Harm (Suicide)	LNE
Chronic Liver Disease and Cirrhosis	34.1
Infant Mortality	15.44

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	80
2. Malignant Neoplasms (Cancer)	67
3. Chronic Lower Respiratory Diseases	29
4. Accidents	20
5. Cerebrovascular Disease	17
6. Diabetes	15
7. Alzheimer's Disease	14
8. Disorders of Lipoprotein Metabolism and Other Lipidemias	9
9. Chronic Liver Disease and Cirrhosis	8
10. Influenza and Pneumonia	5

Percent of Deaths due to tobacco use	19.6
Median age at death	82

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Haakon County

## Demographic Information



Haakon County is located in the west central region of the state and averages 1.1 persons per square mile.

### 2016 Population Information

Subject	Number	Percent
Total population	1,892	100.0
White	1,772	93.7
Black or African American	13	0.7
American Indian & Alaska Native	33	1.7
Asian	11	0.6
Native Hawaiian & Other Pacific Islander	1	0.1
Two or More Races	62	3.3
Under 5 years	105	5.5
Under 18 years	427	22.6
65 years and over	442	23.4

Source: United States Census Bureau, 2016 Population Estimates

## Health Status Indicators 2012-2016

### Natality

Percent of Low Birth Weight Infants	5.0
Percent of Mothers Receiving Care in 1st Trimester	70.0
Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	15.0
Percent of Births Less Than 37 Wks. of Gestation	5.0
Average Age of Mother	28.4
Teenage Birth Rate <sup>2</sup>	LNE
Percent White Births	90.0
Percent American Indian Births	LNE
• Percent Unmarried	21.0
Percent WIC births	33.7
Percent Breastfeeding at discharge	86.0
Percent Payment-Private Insurance	74.0
• Percent Payment-Medicaid	20.0
Percent C-Section	23.0

•Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

All Causes	706.6
Heart Disease	155.8
Malignant Neoplasms (Cancer)	157.2
Trachea, Bronchus, & Lung	34.2
Colon, Rectum, & Anus	40.0
Female Breast	LNE
Pancreas	LNE
Prostate	LNE
Leukemia	27.1
Chronic Lower Respiratory Diseases	77.0
Alzheimer's Disease	LNE
Cerebrovascular Disease	37.8
• Accidents	19.0
Motor Vehicle Accidents	LNE
Diabetes	LNE
Influenza and Pneumonia	35.3
Intentional Self-Harm (Suicide)	LNE
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	LNE

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	28
2. Malignant Neoplasms (Cancer)	24
3. Chronic Lower Respiratory Diseases	16
4. Cerebrovascular Disease	10
5. Influenza and Pneumonia	8
6. Pneumonitis Due to Solids and Liquids	7
7. Urinary Tract Infection, Site Not Specified	5

Percent of Deaths due to tobacco use	12.6
Median age at death	85

•Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Hamlin County

## Demographic Information



Hamlin County is located in the northeastern region of the state and averages 11.6 persons per square mile.

### 2016 Population Information

Subject	Number	Percent
Total population	6,028	100.0
White	5,898	97.8
Black or African American	27	0.4
American Indian & Alaska Native	36	0.6
Asian	19	0.3
Native Hawaiian & Other Pacific Islander	1	0.0
Two or More Races	47	0.8
Under 5 years	557	9.2
Under 18 years	1,871	31.0
65 years and over	1,070	17.8

Source: United States Census Bureau, 2016 Population Estimates

## Health Status Indicators 2012-2016

### Nativity

Percent of Low Birth Weight Infants	4.8
Percent of Mothers Receiving Care in 1st Trimester	73.2
Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	12.9
Percent of Births Less Than 37 Wks. of Gestation	6.3
Average Age of Mother	27.6
● Teenage Birth Rate <sup>2</sup>	5.6
Percent White Births	98.8
Percent American Indian Births	LNE
● Percent Unmarried	13.5
Percent WIC births	34.3
Percent Breastfeeding at discharge	86.7
○ Percent Payment-Private Insurance	76.3
● Percent Payment-Medicaid	18.5
● Percent C-Section	17.6

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

All Causes	762.9
Heart Disease	131.4
Malignant Neoplasms (Cancer)	145.6
Trachea, Bronchus, & Lung	47.8
Colon, Rectum, & Anus	13.0
Female Breast	LNE
Pancreas	12.6
Prostate	33.1
Leukemia	7.4
Chronic Lower Respiratory Diseases	37.5
○ Alzheimer's Disease	90.1
Cerebrovascular Disease	38.2
Accidents	49.9
Motor Vehicle Accidents	19.5
Diabetes	27.0
Influenza and Pneumonia	19.6
Intentional Self-Harm (Suicide)	14.1
Chronic Liver Disease and Cirrhosis	13.3
Infant Mortality	4.99

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Malignant Neoplasms (Cancer)	60
2. Heart Disease	57
3. Alzheimer's Disease	45
T4. Chronic Lower Respiratory Diseases	17
T4. Cerebrovascular Disease	17
6. Accidents	16
7. Vascular Dementia	13
8. Diabetes	12
9. Influenza and Pneumonia	10
10. Essential (Primary) Hypertension and Hypertensive Renal Disease	8
Percent of Deaths due to tobacco use	14.2
Median age at death	84

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Hand County

## Demographic Information



Hand County is located in central South Dakota and averages 2.4 persons per square mile.

### 2016 Population Information

Subject	Number	Percent
Total population	3,319	100.0
White	3,246	97.8
Black or African American	4	0.1
American Indian & Alaska Native	18	0.5
Asian	15	0.5
Native Hawaiian & Other Pacific Islander	3	0.1
Two or More Races	33	1.0
Under 5 years	183	5.5
Under 18 years	694	20.9
65 years and over	840	25.3

Source: United States Census Bureau, 2016 Population Estimates

## Health Status Indicators 2012-2016

### Nativity

Percent of Low Birth Weight Infants	6.4
● Percent of Mothers Receiving Care in 1st Trimester	57.7
● Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	7.6
Percent of Births Less Than 37 Wks. of Gestation	10.5
Average Age of Mother	28.2
Teenage Birth Rate <sup>2</sup>	LNE
Percent White Births	97.7
Percent American Indian Births	LNE
● Percent Unmarried	16.4
● Percent WIC births	19.3
Percent Breastfeeding at discharge	84.7
○ Percent Payment-Private Insurance	82.5
● Percent Payment-Medicaid	13.5
Percent C-Section	33.9

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

All Causes	624.2
● Heart Disease	113.4
Malignant Neoplasms (Cancer)	140.9
● Trachea, Bronchus, & Lung	21.1
Colon, Rectum, & Anus	15.1
Female Breast	LNE
Pancreas	21.0
Prostate	40.3
Leukemia	LNE
Chronic Lower Respiratory Diseases	56.8
Alzheimer's Disease	26.0
Cerebrovascular Disease	53.3
Accidents	45.9
● Motor Vehicle Accidents	6.7
Diabetes	26.6
Influenza and Pneumonia	18.9
Intentional Self-Harm (Suicide)	23.9
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	LNE

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Malignant Neoplasms (Cancer)	45
2. Heart Disease	44
3. Cerebrovascular Disease	25
4. Chronic Lower Respiratory Diseases	21
T5. Accidents	12
T5. Alzheimer's Disease	12
7. Diabetes	10
8. Atherosclerosis	9
9. Influenza and Pneumonia	7
10. Essential (Primary) Hypertension and Hypertensive Renal Disease	6

Percent of Deaths due to tobacco use 24.2

Median age at death 84

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Hanson County

## Demographic Information



Hanson County is located in southeastern South Dakota and averages 7.7 people per square mile.

### 2016 Population Information

Subject	Number	Percent
Total population	3,374	100.0
White	3,291	97.5
Black or African American	18	0.5
American Indian & Alaska Native	15	0.4
Asian	21	0.6
Native Hawaiian & Other Pacific Islander	2	0.1
Two or More Races	27	0.8
Under 5 years	284	8.4
Under 18 years	1,076	31.9
65 years and over	467	13.8

Source: United States Census Bureau, 2016 Population Estimates

## Health Status Indicators 2012-2016

### Nativity

Percent of Low Birth Weight Infants	6.9
Percent of Mothers Receiving Care in 1st Trimester	65.4
● Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	6.1
Percent of Births Less Than 37 Wks. of Gestation	5.6
○ Average Age of Mother	29.5
Teenage Birth Rate <sup>2</sup>	6.8
Percent White Births	100
Percent American Indian Births	LNE
● Percent Unmarried	13.0
● Percent WIC births	14.5
Percent Breastfeeding at discharge	85.3
○ Percent Payment-Private Insurance	79.7
● Percent Payment-Medicaid	7.4
Percent C-Section	23.4

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

All Causes	738.6
Heart Disease	207.6
Malignant Neoplasms (Cancer)	207.1
Trachea, Bronchus, & Lung	57.0
Colon, Rectum, & Anus	37.2
Female Breast	LNE
Pancreas	LNE
Prostate	LNE
Leukemia	LNE
Chronic Lower Respiratory Diseases	LNE
Alzheimer's Disease	33.0
Cerebrovascular Disease	23.0
Accidents	45.4
Motor Vehicle Accidents	35.4
Diabetes	LNE
Influenza and Pneumonia	LNE
Intentional Self-Harm (Suicide)	18.8
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	21.65

### Leading Causes of Death

1. Malignant Neoplasms (Cancer)	30
2. Heart Disease	29
3. Accidents	7
4. Certain Conditions Originating in the Perinatal Period	5

Percent of Deaths due to tobacco use	13.0
Median age at death	74

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Harding County

## Demographic Information



Harding County is located in the northwestern corner of the state and averages 0.5 persons per square mile.

### 2016 Population Information

Subject	Number	Percent
Total population	1,278	100.0
White	1,209	94.6
Black or African American	11	0.9
American Indian & Alaska Native	28	2.2
Asian	3	0.2
Native Hawaiian & Other Pacific Islander	0	0.0
Two or More Races	27	2.1
Under 5 years	81	6.3
Under 18 years	306	23.9
65 years and over	211	16.5

Source: United States Census Bureau, 2016 Population Estimates

## Health Status Indicators 2012-2016

### Nativity

Percent of Low Birth Weight Infants	4.5
Percent of Mothers Receiving Care in 1st Trimester	74.7
● Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	6.8
● Percent of Births Less Than 37 Wks. of Gestation	3.4
Average Age of Mother	28.7
Teenage Birth Rate <sup>2</sup>	LNE
Percent White Births	93.2
Percent American Indian Births	LNE
● Percent Unmarried	14.9
● Percent WIC births	8.0
Percent Breastfeeding at discharge	95.3
○ Percent Payment-Private Insurance	83.7
● Percent Payment-Medicaid	9.3
Percent C-Section	18.2

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

● All Causes	413.5
Heart Disease	100.4
● Malignant Neoplasms (Cancer)	59.9
Trachea, Bronchus, & Lung	LNE
Colon, Rectum, & Anus	LNE
Female Breast	LNE
Pancreas	LNE
Prostate	LNE
Leukemia	LNE
Chronic Lower Respiratory Diseases	LNE
Alzheimer's Disease	LNE
Cerebrovascular Disease	39.3
Accidents	46.2
Motor Vehicle Accidents	LNE
Diabetes	64.5
Influenza and Pneumonia	LNE
Intentional Self-Harm (Suicide)	LNE
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	LNE

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	8
T2. Malignant Neoplasms (Cancer)	5
T2. Diabetes	5
Percent of Deaths due to tobacco use	18.2
Median age at death	75

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Hughes County

## Demographic Information



Hughes County is located in the center of the state and averages 23.0 persons per square mile.

### 2016 Population Information

Subject	Number	Percent
Total population	17,600	100.0
White	14,890	84.6
Black or African American	148	0.8
American Indian & Alaska Native	1,958	11.1
Asian	121	0.7
Native Hawaiian & Other Pacific Islander	5	0.0
Two or More Races	478	2.7
Under 5 years	1,227	7.0
Under 18 years	4,174	23.7
65 years and over	2,863	16.3

Source: United States Census Bureau, 2016 Population Estimates

## Health Status Indicators 2012-2016

### Nativity

Percent of Low Birth Weight Infants	6.4
● Percent of Mothers Receiving Care in 1st Trimester	53.5
○ Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	18.2
Percent of Births Less Than 37 Wks. of Gestation	9.0
Average Age of Mother	27.7
● Teenage Birth Rate <sup>2</sup>	7.3
Percent White Births	75.7
Percent American Indian Births	18.7
Percent Unmarried	40.7
Percent WIC births	34.5
Percent Breastfeeding at discharge	77.4
Percent Payment-Private Insurance	59.9
Percent Payment-Medicaid	35.5
○ Percent C-Section	31.6

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

● All Causes	637.3
● Heart Disease	125.6
Malignant Neoplasms (Cancer)	156.7
Trachea, Bronchus, & Lung	36.5
○ Colon, Rectum, & Anus	28.7
Female Breast	23.1
Pancreas	10.2
● Prostate	7.3
Leukemia	4.2
Chronic Lower Respiratory Diseases	47.6
● Alzheimer's Disease	25.7
Cerebrovascular Disease	46.9
Accidents	38.2
Motor Vehicle Accidents	9.9
Diabetes	32.9
Influenza and Pneumonia	17.5
Intentional Self-Harm (Suicide)	13.8
● Chronic Liver Disease and Cirrhosis	7.7
Infant Mortality	10.87

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Malignant Neoplasms (Cancer)	168
2. Heart Disease	141
3. Cerebrovascular Disease	54
4. Chronic Lower Respiratory Diseases	51
5. Accidents	38
6. Diabetes	36
7. Alzheimer's Disease	31
8. Influenza and Pneumonia	21
9. Intentional Self-Harm (Suicide)	13
10. Unspecified Dementia	10
Percent of Deaths due to tobacco use	17.5
Median age at death	80

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Hutchinson County

## Demographic Information



Hutchinson County is located in the southeastern region of the state and averages 9.0 persons per square mile.

### 2016 Population Information

Subject	Number	Percent
Total population	7,368	100.0
White	7,103	96.4
Black or African American	78	1.1
American Indian & Alaska Native	98	1.3
Asian	17	0.2
Native Hawaiian & Other Pacific Islander	2	0.0
Two or More Races	70	1.0
Under 5 years	596	8.1
Under 18 years	1,807	24.5
65 years and over	1,668	22.6

Source: United States Census Bureau, 2016 Population Estimates

## Health Status Indicators 2012-2016

### Nativity

Percent of Low Birth Weight Infants	5.2
Percent of Mothers Receiving Care in 1st Trimester	69.2
● Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	8.5
Percent of Births Less Than 37 Wks. of Gestation	8.0
○ Average Age of Mother	28.6
● Teenage Birth Rate <sup>2</sup>	4.7
Percent White Births	97.5
Percent American Indian Births	0.9
● Percent Unmarried	20.4
● Percent WIC births	19.2
Percent Breastfeeding at discharge	83.6
○ Percent Payment-Private Insurance	78.1
● Percent Payment-Medicaid	16.2
Percent C-Section	25.0

●Denotes a health status indicator which is significantly lower than the state average.  
○Denotes a health status indicator which is significantly higher than the state average.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

All Causes	689.4
Heart Disease	164.0
● Malignant Neoplasms (Cancer)	116.6
● Trachea, Bronchus, & Lung	25.3
Colon, Rectum, & Anus	15.5
Female Breast	9.5
Pancreas	6.6
Prostate	23.9
Leukemia	LNE
Chronic Lower Respiratory Diseases	34.8
Alzheimer's Disease	42.1
○ Cerebrovascular Disease	56.7
Accidents	59.0
Motor Vehicle Accidents	23.9
Diabetes	15.7
Influenza and Pneumonia	24.8
Intentional Self-Harm (Suicide)	LNE
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	5.61

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	152
2. Malignant Neoplasms (Cancer)	84
3. Alzheimer's Disease	49
4. Cerebrovascular Disease	48
5. Accidents	31
6. Chronic Lower Respiratory Diseases	25
7. Influenza and Pneumonia	23
8. Diabetes	13
9. Essential (Primary) Hypertension and Hypertensive Renal Disease	12
10. Pneumonitis Due to Solids and Liquids	11

Percent of Deaths due to tobacco use	10.1
Median age at death	86

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Hyde County

## Demographic Information



Hyde County is located in the central region of the state and averages 1.7 people per square mile.

### 2016 Population Information

Subject	Number	Percent
Total population	1,352	100.0
White	1,200	88.8
Black or African American	5	0.4
American Indian & Alaska Native	115	8.5
Asian	3	0.2
Native Hawaiian & Other Pacific Islander	0	0.0
Two or More Races	29	2.1
Under 5 years	70	5.2
Under 18 years	273	20.2
65 years and over	337	24.9

Source: United States Census Bureau, 2016 Population Estimates

## Health Status Indicators 2012-2016

### Nativity

Percent of Low Birth Weight Infants	5.2
Percent of Mothers Receiving Care in 1st Trimester	64.5
Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	10.4
Percent of Births Less Than 37 Wks. of Gestation	11.7
Average Age of Mother	27.4
Teenage Birth Rate <sup>2</sup>	LNE
Percent White Births	85.7
Percent American Indian Births	11.7
Percent Unmarried	28.6
Percent WIC births	26.3
Percent Breastfeeding at discharge	80.3
Percent Payment-Private Insurance	76.6
● Percent Payment-Medicaid	20.8
Percent C-Section	19.5

●Denotes a health status indicator which is significantly lower than the state average.  
○Denotes a health status indicator which is significantly higher than the state average.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

All Causes	745.7
○ Heart Disease	253.7
Malignant Neoplasms (Cancer)	108.4
Trachea, Bronchus, & Lung	24.2
Colon, Rectum, & Anus	LNE
Female Breast	LNE
Pancreas	LNE
Prostate	LNE
Leukemia	LNE
Chronic Lower Respiratory Diseases	19.2
Alzheimer's Disease	28.5
Cerebrovascular Disease	78.3
Accidents	60.6
Motor Vehicle Accidents	LNE
Diabetes	LNE
Influenza and Pneumonia	28.5
Intentional Self-Harm (Suicide)	67.5
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	LNE

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	37
2. Malignant Neoplasms (Cancer)	15
3. Cerebrovascular Disease	14
4. Accidents	6
T5. Alzheimer's Disease	5
T5. Influenza and Pneumonia	5
Percent of Deaths due to tobacco use	10.5
Median age at death	85

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Jackson County

## Demographic Information



Jackson County is located in western South Dakota and averages 1.6 persons per square mile.

### 2016 Population Information

Subject	Number	Percent
Total population	3,326	100.0
White	1,424	42.8
Black or African American	29	0.9
American Indian & Alaska Native	1,718	51.7
Asian	6	0.2
Native Hawaiian & Other Pacific Islander	0	0.0
Two or More Races	149	4.5
Under 5 years	352	10.6
Under 18 years	1,115	33.5
65 years and over	448	13.5

Source: United States Census Bureau, 2016 Population Estimates

## Health Status Indicators 2012-2016

### Nativity

Percent of Low Birth Weight Infants	7.4
● Percent of Mothers Receiving Care in 1st Trimester	53.2
○ Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	19.7
○ Percent of Births Less Than 37 Wks. of Gestation	13.3
● Average Age of Mother	26.0
○ Teenage Birth Rate <sup>2</sup>	39.5
Percent White Births	19.2
Percent American Indian Births	77.3
○ Percent Unmarried	74.0
○ Percent WIC births	67.4
● Percent Breastfeeding at discharge	60.9
● Percent Payment-Private Insurance	16.5
○ Percent Payment-Medicaid	57.7
Percent C-Section	23.2

● Denotes a health status indicator which is significantly lower than the state average.  
○ Denotes a health status indicator which is significantly higher than the state average.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

○ All Causes	1,011.7
Heart Disease	203.8
Malignant Neoplasms (Cancer)	193.1
Trachea, Bronchus, & Lung	65.0
Colon, Rectum, & Anus	LNE
Female Breast	LNE
Pancreas	LNE
Prostate	32.7
Leukemia	LNE
Chronic Lower Respiratory Diseases	59.2
Alzheimer's Disease	LNE
Cerebrovascular Disease	22.1
○ Accidents	110.7
○ Motor Vehicle Accidents	67.9
Diabetes	55.4
Influenza and Pneumonia	19.6
Intentional Self-Harm (Suicide)	36.7
○ Chronic Liver Disease and Cirrhosis	55.0
Infant Mortality	10.93

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	36
2. Malignant Neoplasms (Cancer)	35
3. Accidents	17
4. Chronic Lower Respiratory Diseases	11
5. Diabetes	9
6. Chronic Liver Disease and Cirrhosis	8
7. Intentional Self-Harm (Suicide)	6
Percent of Deaths due to tobacco use	11.9
Median age at death	68

● Denotes a health status indicator which is significantly lower than the state average.

○ Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Jerauld County

## Demographic Information



Jerauld County is located in the central region of the state and averages 3.9 persons per square mile.

### 2016 Population Information

Subject	Number	Percent
Total population	2,004	100.0
White	1,956	97.6
Black or African American	3	0.1
American Indian & Alaska Native	13	0.6
Asian	5	0.2
Native Hawaiian & Other Pacific Islander	4	0.2
Two or More Races	23	1.1
Under 5 years	145	7.2
Under 18 years	482	24.1
65 years and over	506	25.2

Source: United States Census Bureau, 2016 Population Estimates

## Health Status Indicators 2012-2016

### Nativity

Percent of Low Birth Weight Infants	6.1
Percent of Mothers Receiving Care in 1st Trimester	72.2
Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	15.7
Percent of Births Less Than 37 Wks. of Gestation	8.7
○ Average Age of Mother	28.8
Teenage Birth Rate <sup>2</sup>	LNE
Percent White Births	99.1
Percent American Indian Births	LNE
Percent Unmarried	27.8
● Percent WIC births	20.9
Percent Breastfeeding at discharge	81.7
○ Percent Payment-Private Insurance	75.7
● Percent Payment-Medicaid	18.3
Percent C-Section	19.1

●Denotes a health status indicator which is significantly lower than the state average.  
○Denotes a health status indicator which is significantly higher than the state average.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

All Causes	623.7
Heart Disease	134.2
Malignant Neoplasms (Cancer)	156.2
Trachea, Bronchus, & Lung	29.9
Colon, Rectum, & Anus	22.5
Female Breast	LNE
Pancreas	28.2
Prostate	LNE
Leukemia	14.0
● Chronic Lower Respiratory Diseases	17.7
○ Alzheimer's Disease	93.2
Cerebrovascular Disease	43.2
Accidents	67.1
Motor Vehicle Accidents	34.8
Diabetes	LNE
Influenza and Pneumonia	16.2
Intentional Self-Harm (Suicide)	LNE
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	LNE

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Malignant Neoplasms (Cancer)	32
T2. Heart Disease	28
T2. Alzheimer's Disease	28
4. Cerebrovascular Disease	11
5. Accidents	8
6. Chronic Lower Respiratory Diseases	5
Percent of Deaths due to tobacco use	11.9
Median age at death	86

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Jones County

## Demographic Information



Jones County is located in western South Dakota and averages 1.0 persons per square mile.

### 2016 Population Information

Subject	Number	Percent
Total population	927	100.0
White	846	91.3
Black or African American	6	0.6
American Indian & Alaska Native	36	3.9
Asian	2	0.2
Native Hawaiian & Other Pacific Islander	8	0.9
Two or More Races	29	3.1
Under 5 years	43	4.6
Under 18 years	187	20.2
65 years and over	208	22.4

Source: United States Census Bureau, 2016 Population Estimates

## Health Status Indicators 2012-2016

### Natality

Percent of Low Birth Weight Infants	7.5
Percent of Mothers Receiving Care in 1st Trimester	56.6
Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	15.1
Percent of Births Less Than 37 Wks. of Gestation	11.3
Average Age of Mother	27.7
Teenage Birth Rate <sup>2</sup>	LNE
Percent White Births	92.5
Percent American Indian Births	7.5
Percent Unmarried	26.4
Percent WIC births	30.2
Percent Breastfeeding at discharge	92.5
Percent Payment-Private Insurance	67.9
Percent Payment-Medicaid	26.4
Percent C-Section	18.9

●Denotes a health status indicator which is significantly lower than the state average.  
○Denotes a health status indicator which is significantly higher than the state average.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

All Causes	567.9
● Heart Disease	87.2
Malignant Neoplasms (Cancer)	115.5
Trachea, Bronchus, & Lung	47.7
Colon, Rectum, & Anus	LNE
Female Breast	LNE
Pancreas	LNE
Prostate	LNE
Leukemia	LNE
Chronic Lower Respiratory Diseases	33.6
Alzheimer's Disease	LNE
Cerebrovascular Disease	LNE
Accidents	LNE
Motor Vehicle Accidents	LNE
Diabetes	LNE
Influenza and Pneumonia	LNE
Intentional Self-Harm (Suicide)	LNE
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	LNE

Leading Causes of Death	Total Deaths
1. Malignant Neoplasms (Cancer)	10
2. Heart Disease	8

Percent of Deaths due to tobacco use	19.6
Median age at death	76

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Kingsbury County

## Demographic Information



Kingsbury County is located in east central South Dakota and averages 6.2 persons per square mile.

### 2016 Population Information

Subject	Number	Percent
Total population	5,001	100.0
White	4,836	96.7
Black or African American	24	0.5
American Indian & Alaska Native	43	0.9
Asian	29	0.6
Native Hawaiian & Other Pacific Islander	1	0.0
Two or More Races	68	1.4
Under 5 years	297	5.9
Under 18 years	1,091	21.8
65 years and over	1,134	22.7

Source: United States Census Bureau, 2016 Population Estimates

## Health Status Indicators 2012-2016

### Nativity

Percent of Low Birth Weight Infants	6.7
Percent of Mothers Receiving Care in 1st Trimester	79.4
Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	12.8
Percent of Births Less Than 37 Wks. of Gestation	7.1
○ Average Age of Mother	28.4
Teenage Birth Rate <sup>2</sup>	6.2
Percent White Births	98.4
Percent American Indian Births	LNE
● Percent Unmarried	21.2
● Percent WIC births	22.4
Percent Breastfeeding at discharge	85.2
○ Percent Payment-Private Insurance	80.0
● Percent Payment-Medicaid	16.8
● Percent C-Section	17.6

●Denotes a health status indicator which is significantly lower than the state average.  
○Denotes a health status indicator which is significantly higher than the state average.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

All Causes	768.9
Heart Disease	161.7
Malignant Neoplasms (Cancer)	192.7
Trachea, Bronchus, & Lung	46.1
Colon, Rectum, & Anus	14.7
Female Breast	30.9
○ Pancreas	30.0
Prostate	LNE
Leukemia	21.3
Chronic Lower Respiratory Diseases	34.2
Alzheimer's Disease	26.9
Cerebrovascular Disease	39.6
Accidents	60.6
○ Motor Vehicle Accidents	46.0
Diabetes	24.3
Influenza and Pneumonia	23.0
Intentional Self-Harm (Suicide)	32.0
Chronic Liver Disease and Cirrhosis	12.8
Infant Mortality	9.62

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	86
2. Malignant Neoplasms (Cancer)	81
3. Cerebrovascular Disease	20
T4. Accidents	18
T4. Chronic Lower Respiratory Diseases	18
6. Alzheimer's Disease	15
T7. Diabetes	10
T7. Influenza and Pneumonia	10
9. Intentional Self-Harm (Suicide)	8
10. Parkinson's Disease	7
Percent of Deaths due to tobacco use	11.5
Median age at death	83

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Lake County

## Demographic Information



Lake County is located in the east central region of the state and averages 19.9 persons per square mile.

### 2016 Population Information

Subject	Number	Percent
Total population	12,909	100.0
White	12,281	95.1
Black or African American	134	1.0
American Indian & Alaska Native	134	1.0
Asian	186	1.4
Native Hawaiian & Other Pacific Islander	3	0.0
Two or More Races	171	1.3
Under 5 years	715	5.5
Under 18 years	2,566	19.9
65 years and over	2,831	21.9

Source: United States Census Bureau, 2016 Population Estimates

## Health Status Indicators 2012-2016

### Nativity

Percent of Low Birth Weight Infants	4.9
Percent of Mothers Receiving Care in 1st Trimester	74.6
● Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	11.4
Percent of Births Less Than 37 Wks. of Gestation	7.5
○ Average Age of Mother	28.4
Teenage Birth Rate <sup>2</sup>	7.9
Percent White Births	95.2
Percent American Indian Births	1.7
● Percent Unmarried	25.3
● Percent WIC births	25.3
Percent Breastfeeding at discharge	82.0
○ Percent Payment-Private Insurance	74.7
● Percent Payment-Medicaid	18.3
Percent C-Section	23.3

●Denotes a health status indicator which is significantly lower than the state average.  
○Denotes a health status indicator which is significantly higher than the state average.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

● All Causes	622.6
Heart Disease	136.3
● Malignant Neoplasms (Cancer)	129.4
Trachea, Bronchus, & Lung	35.4
Colon, Rectum, & Anus	16.4
Female Breast	12.5
Pancreas	5.4
Prostate	18.1
Leukemia	5.0
Chronic Lower Respiratory Diseases	41.9
Alzheimer's Disease	42.1
○ Cerebrovascular Disease	57.9
● Accidents	29.7
Motor Vehicle Accidents	12.8
Diabetes	30.0
Influenza and Pneumonia	10.6
Intentional Self-Harm (Suicide)	15.6
Chronic Liver Disease and Cirrhosis	11.6
Infant Mortality	6.09

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	134
2. Malignant Neoplasms (Cancer)	125
3. Cerebrovascular Disease	52
4. Alzheimer's Disease	39
5. Chronic Lower Respiratory Diseases	38
6. Diabetes	29
7. Accidents	22
8. Unspecified Dementia	19
9. Influenza and Pneumonia	10
10. Chronic Liver Disease and Cirrhosis	9

Percent of Deaths due to tobacco use	22.3
Median age at death	80

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Lawrence County

## Demographic Information



Lawrence County is located along the Wyoming border and averages 30.1 persons per square mile.

### 2016 Population Information

Subject	Number	Percent
Total population	25,281	100.0
White	23,670	93.6
Black or African American	249	1.0
American Indian & Alaska Native	579	2.3
Asian	251	1.0
Native Hawaiian & Other Pacific Islander	15	0.1
Two or More Races	517	2.0
Under 5 years	1,180	4.7
Under 18 years	4,604	18.2
65 years and over	4,963	19.6

Source: United States Census Bureau, 2016 Population Estimates

## Health Status Indicators 2012-2016

### Natality

○ Percent of Low Birth Weight Infants	8.3
○ Percent of Mothers Receiving Care in 1st Trimester	80.1
Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	17.5
○ Percent of Births Less Than 37 Wks. of Gestation	10.5
Average Age of Mother	27.8
Teenage Birth Rate <sup>2</sup>	9.1
Percent White Births	92.5
Percent American Indian Births	2.1
Percent Unmarried	37.8
Percent WIC births	31.8
Percent Breastfeeding at discharge	84.1
Percent Payment-Private Insurance	61.3
Percent Payment-Medicaid	31.4
● Percent C-Section	20.9

### Mortality<sup>3</sup>

● All Causes	659.5
Heart Disease	134.3
Malignant Neoplasms (Cancer)	147.9
Trachea, Bronchus, & Lung	35.0
Colon, Rectum, & Anus	16.5
Female Breast	17.7
Pancreas	9.2
Prostate	24.9
● Leukemia	3.1
○ Chronic Lower Respiratory Diseases	55.2
Alzheimer's Disease	29.1
Cerebrovascular Disease	31.5
Accidents	42.1
Motor Vehicle Accidents	13.1
Diabetes	23.9
Influenza and Pneumonia	14.8
Intentional Self-Harm (Suicide)	14.8
Chronic Liver Disease and Cirrhosis	11.7
Infant Mortality	6.80

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Malignant Neoplasms (Cancer)	257
2. Heart Disease	243
3. Chronic Lower Respiratory Diseases	103
4. Accidents	65
5. Cerebrovascular Disease	58
6. Alzheimer's Disease	57
7. Diabetes	45
8. Influenza and Pneumonia	27
9. Vascular Dementia	20
10. Intentional Self-Harm (Suicide)	19

Percent of Deaths due to tobacco use	22.0
Median age at death	81

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Lincoln County

## Demographic Information



Lincoln County is located in southeastern South Dakota and averages 77.7 persons per square mile.

### 2016 Population Information

Subject	Number	Percent
Total population	54,469	100.0
White	51,820	95.1
Black or African American	740	1.4
American Indian & Alaska Native	316	0.6
Asian	705	1.3
Native Hawaiian & Other Pacific Islander	33	0.1
Two or More Races	855	1.6
Under 5 years	4,125	7.6
Under 18 years	15,558	28.6
65 years and over	6,340	11.6

Source: United States Census Bureau, 2016 Population Estimates

## Health Status Indicators 2012-2016

### Nativity

Percent of Low Birth Weight Infants	5.4
○ Percent of Mothers Receiving Care in 1st Trimester	85.4
● Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	5.9
Percent of Births Less Than 37 Wks. of Gestation	7.4
○ Average Age of Mother	29.4
● Teenage Birth Rate <sup>2</sup>	4.2
Percent White Births	96.7
Percent American Indian Births	0.4
● Percent Unmarried	16.3
● Percent WIC births	8.7
○ Percent Breastfeeding at discharge	84.0
○ Percent Payment-Private Insurance	85.1
● Percent Payment-Medicaid	10.4
Percent C-Section	26.0

●Denotes a health status indicator which is significantly lower than the state average.  
○Denotes a health status indicator which is significantly higher than the state average.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

● All Causes	470.0
● Heart Disease	98.7
● Malignant Neoplasms (Cancer)	116.1
● Trachea, Bronchus, & Lung	28.7
● Colon, Rectum, & Anus	9.9
Female Breast	17.7
Pancreas	14.9
Prostate	16.6
Leukemia	5.0
● Chronic Lower Respiratory Diseases	24.0
Alzheimer's Disease	36.7
Cerebrovascular Disease	29.1
● Accidents	22.2
● Motor Vehicle Accidents	6.5
● Diabetes	10.3
Influenza and Pneumonia	11.4
● Intentional Self-Harm (Suicide)	9.3
● Chronic Liver Disease and Cirrhosis	2.6
Infant Mortality	5.85

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Malignant Neoplasms (Cancer)	263
2. Heart Disease	229
3. Alzheimer's Disease	82
4. Cerebrovascular Disease	66
5. Accidents	56
6. Chronic Lower Respiratory Diseases	55
7. Unspecified Dementia	28
8. Influenza and Pneumonia	26
9. Diabetes	23
10. Intentional Self-Harm (Suicide)	21
Percent of Deaths due to tobacco use	16.1
Median age at death	80

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Lyman County

## Demographic Information



Lyman County is located in central South Dakota and averages 2.3 persons per square mile.

### 2016 Population Information

Subject	Number	Percent
Total population	3,894	100.0
White	2,177	55.9
Black or African American	24	0.6
American Indian & Alaska Native	1,545	39.7
Asian	13	0.3
Native Hawaiian & Other Pacific Islander	1	0.0
Two or More Races	134	3.4
Under 5 years	352	9.0
Under 18 years	1,134	29.1
65 years and over	574	14.7

Source: United States Census Bureau, 2016 Population Estimates

## Health Status Indicators 2012-2016

### Nativity

Percent of Low Birth Weight Infants	6.7
● Percent of Mothers Receiving Care in 1st Trimester	47.8
○ Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	25.1
Percent of Births Less Than 37 Wks. of Gestation	10.2
● Average Age of Mother	26.3
Teenage Birth Rate <sup>2</sup>	28.1
Percent White Births	39.9
Percent American Indian Births	54.5
○ Percent Unmarried	60.1
○ Percent WIC births	60.3
● Percent Breastfeeding at discharge	62.3
● Percent Payment-Private Insurance	35.6
○ Percent Payment-Medicaid	57.1
Percent C-Section	31.2

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

○ All Causes	869.9
Heart Disease	165.7
Malignant Neoplasms (Cancer)	172.8
Trachea, Bronchus, & Lung	49.0
Colon, Rectum, & Anus	26.7
Female Breast	42.1
Pancreas	LNE
Prostate	56.0
Leukemia	LNE
Chronic Lower Respiratory Diseases	37.7
Alzheimer's Disease	46.3
Cerebrovascular Disease	39.7
○ Accidents	104.6
○ Motor Vehicle Accidents	53.5
Diabetes	35.0
Influenza and Pneumonia	23.2
Intentional Self-Harm (Suicide)	55.6
Chronic Liver Disease and Cirrhosis	22.4
Infant Mortality	LNE

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Malignant Neoplasms (Cancer)	39
2. Heart Disease	34
3. Accidents	18
T4. Alzheimer's Disease	9
T4. Diabetes	9
T4. Intentional Self-Harm (Suicide)	9
T7. Chronic Lower Respiratory Diseases	8
T7. Cerebrovascular Disease	8
Percent of Deaths due to tobacco use	27.5
Median age at death	72

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# McCook County

## Demographic Information



McCook County is located in eastern South Dakota and averages 9.8 persons per square mile.

### 2016 Population Information

Subject	Number	Percent
Total population	5,625	100.0
White	5,462	97.1
Black or African American	26	0.5
American Indian & Alaska Native	57	1.0
Asian	11	0.2
Native Hawaiian & Other Pacific Islander	3	0.1
Two or More Races	66	1.2
Under 5 years	424	7.5
Under 18 years	1,498	26.6
65 years and over	1,117	19.9

Source: United States Census Bureau, 2016 Population Estimates

## Health Status Indicators 2012-2016

### Nativity

Percent of Low Birth Weight Infants	5.0
Percent of Mothers Receiving Care in 1st Trimester	74.0
● Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	9.5
Percent of Births Less Than 37 Wks. of Gestation	8.2
○ Average Age of Mother	28.6
Teenage Birth Rate <sup>2</sup>	7.2
Percent White Births	97.6
Percent American Indian Births	1.1
● Percent Unmarried	24.1
● Percent WIC births	27.1
Percent Breastfeeding at discharge	79.4
○ Percent Payment-Private Insurance	75.0
● Percent Payment-Medicaid	18.1
Percent C-Section	25.7

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

○ All Causes	895.0
○ Heart Disease	200.5
○ Malignant Neoplasms (Cancer)	232.8
Trachea, Bronchus, & Lung	59.2
Colon, Rectum, & Anus	29.7
Female Breast	22.7
Pancreas	17.4
Prostate	40.0
Leukemia	LNE
● Chronic Lower Respiratory Diseases	20.4
○ Alzheimer's Disease	74.5
Cerebrovascular Disease	47.5
Accidents	63.3
Motor Vehicle Accidents	30.5
Diabetes	28.3
Influenza and Pneumonia	11.9
Intentional Self-Harm (Suicide)	15.1
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	LNE

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Malignant Neoplasms (Cancer)	95
2. Heart Disease	94
3. Alzheimer's Disease	41
4. Cerebrovascular Disease	24
5. Accidents	19
6. Diabetes	13
7. Septicemia	11
8. Chronic Lower Respiratory Diseases	9
9. Influenza and Pneumonia	7
10. Vascular Dementia	6
Percent of Deaths due to tobacco use	11.2
Median age at death	81

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# McPherson County

## Demographic Information



McPherson County is located in the north central region of the state and averages 2.2 persons per square mile.

### 2016 Population Information

Subject	Number	Percent
Total population	2,438	100.0
White	2,400	98.4
Black or African American	7	0.3
American Indian & Alaska Native	5	0.2
Asian	6	0.2
Native Hawaiian & Other Pacific Islander	1	0.0
Two or More Races	19	0.8
Under 5 years	174	7.1
Under 18 years	565	23.2
65 years and over	679	27.9

Source: United States Census Bureau, 2016 Population Estimates

## Health Status Indicators 2012-2016

### Nativity

Percent of Low Birth Weight Infants	7.0
● Percent of Mothers Receiving Care in 1st Trimester	52.8
Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	10.2
Percent of Births Less Than 37 Wks. of Gestation	9.4
○ Average Age of Mother	28.9
Teenage Birth Rate <sup>2</sup>	LNE
Percent White Births	98.4
Percent American Indian Births	LNE
● Percent Unmarried	16.4
Percent WIC births	25.4
Percent Breastfeeding at discharge	82.8
Percent Payment-Private Insurance	72.4
● Percent Payment-Medicaid	20.5
Percent C-Section	25.0

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

All Causes	718.4
○ Heart Disease	230.7
Malignant Neoplasms (Cancer)	125.5
Trachea, Bronchus, & Lung	31.8
Colon, Rectum, & Anus	14.7
Female Breast	LNE
Pancreas	LNE
Prostate	LNE
Leukemia	LNE
● Chronic Lower Respiratory Diseases	22.6
Alzheimer's Disease	30.3
Cerebrovascular Disease	41.6
Accidents	58.2
Motor Vehicle Accidents	27.6
Diabetes	35.9
Influenza and Pneumonia	LNE
Intentional Self-Harm (Suicide)	52.8
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	LNE

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	67
2. Malignant Neoplasms (Cancer)	35
3. Cerebrovascular Disease	15
T4. Accidents	12
T4. Alzheimer's Disease	12
T6. Chronic Lower Respiratory Diseases	8
T6. Diabetes	8

Percent of Deaths due to tobacco use	9.8
Median age at death	84

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Marshall County

## Demographic Information



Marshall County is located in the northeastern part of the state and averages 5.6 persons per square mile.

### 2016 Population Information

Subject	Number	Percent
Total population	4,801	100.0
White	4,232	88.1
Black or African American	50	1.0
American Indian & Alaska Native	418	8.7
Asian	12	0.2
Native Hawaiian & Other Pacific Islander	0	0.0
Two or More Races	89	1.9
Under 5 years	344	7.2
Under 18 years	1,086	22.6
65 years and over	969	20.2

Source: United States Census Bureau, 2016 Population Estimates

## Health Status Indicators 2012-2016

### Nativity

Percent of Low Birth Weight Infants	6.6
● Percent of Mothers Receiving Care in 1st Trimester	57.7
Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	16.5
Percent of Births Less Than 37 Wks. of Gestation	7.2
○ Average Age of Mother	28.6
Teenage Birth Rate <sup>2</sup>	7.8
Percent White Births	83.2
Percent American Indian Births	11.1
● Percent Unmarried	29.1
● Percent WIC births	26.4
Percent Breastfeeding at discharge	82.0
Percent Payment-Private Insurance	67.2
● Percent Payment-Medicaid	22.3
Percent C-Section	26.9

●Denotes a health status indicator which is significantly lower than the state average.  
○Denotes a health status indicator which is significantly higher than the state average.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

All Causes	702.7
Heart Disease	150.6
Malignant Neoplasms (Cancer)	159.1
Trachea, Bronchus, & Lung	44.6
Colon, Rectum, & Anus	13.2
Female Breast	29.4
Pancreas	20.9
Prostate	32.0
Leukemia	7.3
Chronic Lower Respiratory Diseases	47.1
Alzheimer's Disease	24.4
○ Cerebrovascular Disease	64.5
Accidents	61.5
Motor Vehicle Accidents	22.5
Diabetes	13.1
Influenza and Pneumonia	18.5
Intentional Self-Harm (suicide)	25.9
Chronic Liver Disease and Cirrhosis	7.4
Infant Mortality	11.98

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	59
2. Malignant Neoplasms (Cancer)	56
3. Cerebrovascular Disease	25
4. Chronic Lower Respiratory Diseases	19
5. Accidents	17
6. Alzheimer's Disease	10
T7. Influenza and Pneumonia	7
T7. Intentional Self-Harm (Suicide)	7
Percent of Deaths due to tobacco use	18.1
Median age at death	81

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Meade County

## Demographic Information



Meade County is located in west central South Dakota and averages 7.3 persons per square mile.

### 2016 Population Information

Subject	Number	Percent
Total population	27,693	100.0
White	25,217	91.1
Black or African American	525	1.9
American Indian & Alaska Native	793	2.9
Asian	326	1.2
Native Hawaiian & Other Pacific Islander	26	0.1
Two or More Races	806	2.9
Under 5 years	1,689	6.1
Under 18 years	6,519	23.5
65 years and over	4,057	14.6

Source: United States Census Bureau, 2016 Population Estimates

## Health Status Indicators 2012-2016

### Nativity

Percent of Low Birth Weight Infants	7.2
Percent of Mothers Receiving Care in 1st Trimester	77.1
Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	15.3
Percent of Births Less Than 37 Wks. of Gestation	8.9
Average Age of Mother	27.9
Teenage Birth Rate <sup>2</sup>	8.6
Percent White Births	92.0
Percent American Indian Births	3.4
● Percent Unmarried	28.7
● Percent WIC births	29.7
○ Percent Breastfeeding at discharge	86.3
Percent Payment-Private Insurance	55.8
● Percent Payment-Medicaid	25.2
● Percent C-Section	20.3

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

All Causes	683.5
Heart Disease	142.2
Malignant Neoplasms (Cancer)	153.8
Trachea, Bronchus, & Lung	47.1
Colon, Rectum, & Anus	15.4
Female Breast	12.4
Pancreas	9.9
Prostate	20.9
Leukemia	6.3
○ Chronic Lower Respiratory Diseases	68.3
● Alzheimer's Disease	20.4
Cerebrovascular Disease	33.0
● Accidents	33.2
Motor Vehicle Accidents	12.2
● Diabetes	14.3
Influenza and Pneumonia	19.7
Intentional Self-Harm (suicide)	23.2
Chronic Liver Disease and Cirrhosis	13.5
Infant Mortality	3.25

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Malignant Neoplasms (Cancer)	224
2. Heart Disease	195
3. Chronic Lower Respiratory Diseases	95
4. Cerebrovascular Disease	45
5. Accidents	44
6. Intentional Self-Harm (Suicide)	30
T7. Alzheimer's Disease	27
T7. Influenza and Pneumonia	27
9. Unspecified Dementia	25
10. Diabetes	21
Percent of Deaths due to tobacco use	19.9
Median age at death	78

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Mellette County

## Demographic Information



Mellette County is located in the south central region of the state and averages 1.6 persons per square mile.

### 2016 Population Information

Subject	Number	Percent
Total population	2,102	100.0
White	830	39.5
Black or African American	9	0.4
American Indian & Alaska Native	1,145	54.5
Asian	6	0.3
Native Hawaiian & Other Pacific Islander	0	0.0
Two or More Races	112	5.3
Under 5 years	174	8.3
Under 18 years	650	30.9
65 years and over	338	16.1

Source: United States Census Bureau, 2016 Population Estimates

## Health Status Indicators 2012-2016

### Nativity

Percent of Low Birth Weight Infants	5.5
● Percent of Mothers Receiving Care in 1st Trimester	48.9
Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	20.7
Percent of Births Less Than 37 Wks. of Gestation	7.1
● Average Age of Mother	25.4
Teenage Birth Rate <sup>2</sup>	35.7
Percent White Births	17.7
Percent American Indian Births	71.3
○ Percent Unmarried	74.3
○ Percent WIC births	60.9
Percent Breastfeeding at discharge	71.3
● Percent Payment-Private Insurance	14.2
○ Percent Payment-Medicaid	60.1
○ Percent C-Section	40.4

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

○ All Causes	1,017.4
Heart Disease	131.0
Malignant Neoplasms (Cancer)	146.2
Trachea, Bronchus, & Lung	37.4
Colon, Rectum, & Anus	LNE
Female Breast	LNE
Pancreas	LNE
Prostate	LNE
Leukemia	LNE
Chronic Lower Respiratory Diseases	76.0
Alzheimer's Disease	38.1
Cerebrovascular Disease	23.0
Accidents	106.0
Motor Vehicle Accidents	69.1
○ Diabetes	96.2
Influenza and Pneumonia	LNE
Intentional Self-Harm (suicide)	LNE
○ Chronic Liver Disease and Cirrhosis	111.6
Infant Mortality	16.39

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Malignant Neoplasms (Cancer)	19
2. Heart Disease	17
T3. Accidents	10
T3. Chronic Lower Respiratory Diseases	10
T3. Diabetes	10
6. Chronic Liver Disease and Cirrhosis	9
7. Alzheimer's Disease	5
Percent of Deaths due to tobacco use	17.2
Median age at death	75

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Miner County

## Demographic Information



Miner County is located in the east central region of the state and averages 4.2 persons per square mile.

### 2016 Population Information

Subject	Number	Percent
Total population	2,281	100.0
White	2,201	96.5
Black or African American	26	1.9
American Indian & Alaska Native	12	0.5
Asian	10	0.4
Native Hawaiian & Other Pacific Islander	0	0.0
Two or More Races	32	1.4
Under 5 years	128	5.6
Under 18 years	537	23.5
65 years and over	497	21.8

Source: United States Census Bureau, 2016 Population Estimates

## Health Status Indicators 2012-2016

### Nativity

Percent of Low Birth Weight Infants	4.9
Percent of Mothers Receiving Care in 1st Trimester	68.6
● Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	6.5
Percent of Births Less Than 37 Wks. of Gestation	5.7
○ Average Age of Mother	29.5
Teenage Birth Rate <sup>2</sup>	LNE
Percent White Births	97.6
Percent American Indian Births	LNE
● Percent Unmarried	13.8
● Percent WIC births	13.0
Percent Breastfeeding at discharge	82.8
○ Percent Payment-Private Insurance	78.7
● Percent Payment-Medicaid	17.2
Percent C-Section	28.5

●Denotes a health status indicator which is significantly lower than the state average.  
○Denotes a health status indicator which is significantly higher than the state average.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

All Causes	774.2
Heart Disease	184.1
Malignant Neoplasms (Cancer)	172.9
Trachea, Bronchus, & Lung	49.1
Colon, Rectum, & Anus	LNE
Female Breast	LNE
Pancreas	12.6
Prostate	LNE
Leukemia	LNE
Chronic Lower Respiratory Diseases	39.5
Alzheimer's Disease	57.0
Cerebrovascular Disease	41.6
Accidents	56.6
Motor Vehicle Accidents	26.5
Diabetes	38.6
Influenza and Pneumonia	13.5
Intentional Self-Harm (Suicide)	LNE
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	LNE

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	40
2. Malignant Neoplasms (Cancer)	35
3. Alzheimer's Disease	15
4. Cerebrovascular Disease	10
T5. Accidents	8
T5. Chronic Lower Respiratory Diseases	8
T5. Diabetes	8
8. Unspecified Dementia	5
Percent of Deaths due to tobacco use	20.5
Median age at death	83

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Minnehaha County

## Demographic Information



Minnehaha County is located in southeastern South Dakota and averages 210.0 persons per square mile.

### 2016 Population Information

Subject	Number	Percent
Total population	187,318	100.0
White	163,461	87.3
Black or African American	9,770	5.2
American Indian & Alaska Native	5,059	2.7
Asian	4,261	2.3
Native Hawaiian & Other Pacific Islander	203	0.1
Two or More Races	4,564	2.4
Under 5 years	14,622	7.8
Under 18 years	47,340	25.3
65 years and over	24,108	12.9

Source: United States Census Bureau, 2016 Population Estimates

## Health Status Indicators 2012-2016

### Nativity

Percent of Low Birth Weight Infants	6.6
○ Percent of Mothers Receiving Care in 1st Trimester	76.6
● Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	11.8
Percent of Births Less Than 37 Wks. of Gestation	7.9
○ Average Age of Mother	28.2
Teenage Birth Rate <sup>2</sup>	12.3
Percent White Births	80.7
Percent American Indian Births	5.0
● Percent Unmarried	33.8
● Percent WIC births	26.8
Percent Breastfeeding at discharge	80.2
○ Percent Payment-Private Insurance	65.4
● Percent Payment-Medicaid	29.8
Percent C-Section	25.4

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

All Causes	699.9
Heart Disease	145.5
Malignant Neoplasms (Cancer)	167.1
Trachea, Bronchus, & Lung	44.4
Colon, Rectum, & Anus	17.5
Female Breast	21.3
Pancreas	11.1
Prostate	19.7
Leukemia	7.0
Chronic Lower Respiratory Diseases	44.2
○ Alzheimer's Disease	42.9
Cerebrovascular Disease	37.5
Accidents	43.7
● Motor Vehicle Accidents	10.2
● Diabetes	15.7
● Influenza and Pneumonia	13.2
Intentional Self-Harm (Suicide)	17.0
● Chronic Liver Disease and Cirrhosis	9.8
● Infant Mortality	4.66

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Malignant Neoplasms (Cancer)	1,549
2. Heart Disease	1,386
3. Alzheimer's Disease	414
4. Chronic Lower Respiratory Diseases	410
5. Accidents	404
6. Cerebrovascular Disease	351
7. Intentional Self-Harm (Suicide)	152
8. Diabetes	143
9. Influenza and Pneumonia	130
10. Chronic Liver Disease and Cirrhosis	91
Percent of Deaths due to tobacco use	17.5
Median age at death	78

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Moody County

## Demographic Information



Moody County is located on the Minnesota border and averages 12.5 persons per square mile.

## 2015 Population Information

Subject	Number	Percent
Total population	6,505	100.0
White	5,170	79.5
Black or African American	114	1.8
American Indian & Alaska Native	904	13.9
Asian	115	1.8
Native Hawaiian & Other Pacific Islander	0	0.0
Two or More Races	202	3.1
Under 5 years	482	7.4
Under 18 years	1,686	25.9
65 years and over	1,121	17.2

Source: United States Census Bureau, 2016 Population Estimates

## Health Status Indicators 2012-2016

### Nativity

Percent of Low Birth Weight Infants	6.3
Percent of Mothers Receiving Care in 1st Trimester	71.4
Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	16.4
Percent of Births Less Than 37 Wks. of Gestation	9.5
Average Age of Mother	27.8
Teenage Birth Rate <sup>2</sup>	16.6
Percent White Births	73.6
Percent American Indian Births	22.1
Percent Unmarried	40.8
Percent WIC births	35.5
Percent Breastfeeding at discharge	77.7
Percent Payment-Private Insurance	60.0
Percent Payment-Medicaid	30.5
Percent C-Section	24.0

●Denotes a health status indicator which is significantly lower than the state average.  
○Denotes a health status indicator which is significantly higher than the state average.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

● All Causes	621.0
Heart Disease	154.7
Malignant Neoplasms (Cancer)	158.3
Trachea, Bronchus, & Lung	27.3
Colon, Rectum, & Anus	21.6
Female Breast	LNE
Pancreas	13.9
Prostate	28.4
Leukemia	LNE
● Chronic Lower Respiratory Diseases	17.4
Alzheimer's Disease	47.7
Cerebrovascular Disease	30.1
Accidents	45.5
Motor Vehicle Accidents	16.5
Diabetes	17.8
Influenza and Pneumonia	LNE
Intentional Self-Harm (Suicide)	18.1
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	6.48

Leading Causes of Death	Total Deaths
1. Heart Disease	69
2. Malignant Neoplasms (Cancer)	65
3. Alzheimer's Disease	23
4. Accidents	15
5. Cerebrovascular Disease	13
T6. Chronic Lower Respiratory Diseases	8
T6. Diabetes	8
T8. Vascular Dementia	7
T8. Peripheral Vascular Disease, Unspecified	7
10. Intentional Self-Harm (Suicide)	6
Percent of Deaths due to tobacco use	16.5
Median age at death	80

●Denotes a health status indicator which is significantly lower than the state average.


○Denotes a health status indicator which is significantly higher than the state average.

<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.


Source: South Dakota Department of Health, Office of Health Statistics

# Oglala Lakota County

Demographic Information	Health Status Indicators 2012-2016																																																										
 <p>Oglala Lakota County (formerly known as Shannon County) is located in the southwestern part of the state, along the Nebraska border and averages 6.5 persons per square mile.</p> <p><b>2015 Population Information</b></p> <table border="1"> <thead> <tr> <th>Subject</th> <th>Number</th> <th>Percent</th> </tr> </thead> <tbody> <tr> <td>Total population</td> <td>14,415</td> <td>100.0</td> </tr> <tr> <td>White</td> <td>733</td> <td>5.1</td> </tr> <tr> <td>Black or African American</td> <td>46</td> <td>0.3</td> </tr> <tr> <td>American Indian &amp; Alaska Native</td> <td>13,382</td> <td>92.8</td> </tr> <tr> <td>Asian</td> <td>25</td> <td>0.2</td> </tr> <tr> <td>Native Hawaiian &amp; Other Pacific Islander</td> <td>7</td> <td>0.1</td> </tr> <tr> <td>Two or More Races</td> <td>222</td> <td>1.5</td> </tr> <tr> <td>Under 5 years</td> <td>1,589</td> <td>11.0</td> </tr> <tr> <td>Under 18 years</td> <td>5,432</td> <td>37.7</td> </tr> <tr> <td>65 years and over</td> <td>993</td> <td>6.9</td> </tr> </tbody> </table> <p>Source: United States Census Bureau, 2016 Population Estimates</p>	Subject	Number	Percent	Total population	14,415	100.0	White	733	5.1	Black or African American	46	0.3	American Indian & Alaska Native	13,382	92.8	Asian	25	0.2	Native Hawaiian & Other Pacific Islander	7	0.1	Two or More Races	222	1.5	Under 5 years	1,589	11.0	Under 18 years	5,432	37.7	65 years and over	993	6.9	<p><b>Natality</b></p> <ul style="list-style-type: none"> <li>○ Percent of Low Birth Weight Infants 8.1</li> <li>● Percent of Mothers Receiving Care in 1st Trimester 58.3</li> <li>○ Percent of Mothers Who Used Tobacco While Pregnant<sup>1</sup> 18.9</li> <li>○ Percent of Births Less Than 37 Wks. of Gestation 11.4</li> <li>● Average Age of Mother 24.9</li> <li>○ Teenage Birth Rate<sup>2</sup> 42.0</li> <li>○ Percent White Births 1.5</li> <li>○ Percent American Indian Births 97.4</li> <li>○ Percent Unmarried 88.2</li> <li>○ Percent WIC births 73.5</li> <li>● Percent Breastfeeding at discharge 55.1</li> <li>● Percent Payment-Private Insurance 3.8</li> <li>○ Percent Payment-Medicaid 61.7</li> <li>● Percent C-Section 22.2</li> </ul> <p>●Denotes a health status indicator which is significantly lower than the state average. ○Denotes a health status indicator which is significantly higher than the state average.</p> <p><sup>1</sup>Data for mothers who used tobacco are self-reported. <sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.</p>	<p><b>Mortality<sup>3</sup></b></p> <ul style="list-style-type: none"> <li>○ All Causes 1,378.4</li> <li>Heart Disease 162.5</li> <li>○ Malignant Neoplasms (Cancer) 221.3 <ul style="list-style-type: none"> <li>Trachea, Bronchus, &amp; Lung 60.2</li> <li>Colon, Rectum, &amp; Anus 23.1</li> <li>Female Breast 9.4</li> <li>Pancreas 17.4</li> <li>Prostate LNE</li> <li>Leukemia LNE</li> </ul> </li> <li>Chronic Lower Respiratory Diseases 47.5</li> <li>● Alzheimer's Disease 14.8</li> <li>Cerebrovascular Disease 41.4 <ul style="list-style-type: none"> <li>○ Accidents 151.6 <ul style="list-style-type: none"> <li>○ Motor Vehicle Accidents 67.2</li> </ul> </li> <li>○ Diabetes 105.0</li> <li>○ Influenza and Pneumonia 42.0</li> <li>○ Intentional Self-Harm (Suicide) 46.4</li> <li>○ Chronic Liver Disease and Cirrhosis 160.9</li> <li>○ Infant Mortality 18.96</li> </ul> </li> </ul> <p><b>Leading Causes of Death</b></p> <table border="1"> <thead> <tr> <th>Leading Causes of Death</th> <th>Total Deaths</th> </tr> </thead> <tbody> <tr> <td>1. Accidents</td> <td>93</td> </tr> <tr> <td>2. Malignant Neoplasms (Cancer)</td> <td>92</td> </tr> <tr> <td>3. Chronic Liver Disease and Cirrhosis</td> <td>85</td> </tr> <tr> <td>4. Heart Disease</td> <td>67</td> </tr> <tr> <td>5. Diabetes</td> <td>51</td> </tr> <tr> <td>6. Intentional Self-Harm (Suicide)</td> <td>37</td> </tr> <tr> <td>7. Septicemia</td> <td>20</td> </tr> <tr> <td>T8. Cerebrovascular Disease</td> <td>19</td> </tr> <tr> <td>T8. Assault (Homicide)</td> <td>19</td> </tr> <tr> <td>T10. Chronic Lower Respiratory Diseases</td> <td>17</td> </tr> <tr> <td>T10. Influenza and Pneumonia</td> <td>17</td> </tr> </tbody> </table> <p>Percent of Deaths due to tobacco use 10.9 Median age at death 54</p> <p>●Denotes a health status indicator which is significantly lower than the state average. ○Denotes a health status indicator which is significantly higher than the state average. <sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.</p> <p>See technical notes for more information. Source: South Dakota Department of Health, Office of Health Statistics</p>	Leading Causes of Death	Total Deaths	1. Accidents	93	2. Malignant Neoplasms (Cancer)	92	3. Chronic Liver Disease and Cirrhosis	85	4. Heart Disease	67	5. Diabetes	51	6. Intentional Self-Harm (Suicide)	37	7. Septicemia	20	T8. Cerebrovascular Disease	19	T8. Assault (Homicide)	19	T10. Chronic Lower Respiratory Diseases	17	T10. Influenza and Pneumonia	17
Subject	Number	Percent																																																									
Total population	14,415	100.0																																																									
White	733	5.1																																																									
Black or African American	46	0.3																																																									
American Indian & Alaska Native	13,382	92.8																																																									
Asian	25	0.2																																																									
Native Hawaiian & Other Pacific Islander	7	0.1																																																									
Two or More Races	222	1.5																																																									
Under 5 years	1,589	11.0																																																									
Under 18 years	5,432	37.7																																																									
65 years and over	993	6.9																																																									
Leading Causes of Death	Total Deaths																																																										
1. Accidents	93																																																										
2. Malignant Neoplasms (Cancer)	92																																																										
3. Chronic Liver Disease and Cirrhosis	85																																																										
4. Heart Disease	67																																																										
5. Diabetes	51																																																										
6. Intentional Self-Harm (Suicide)	37																																																										
7. Septicemia	20																																																										
T8. Cerebrovascular Disease	19																																																										
T8. Assault (Homicide)	19																																																										
T10. Chronic Lower Respiratory Diseases	17																																																										
T10. Influenza and Pneumonia	17																																																										

# Pennington County

## Demographic Information



Pennington County is located on the Wyoming border and averages 36.4 persons per square mile.

## 2016 Population Information

Subject	Number	Percent
Total population	109,372	100.0
White	91,462	83.6
Black or African American	1,456	1.3
American Indian & Alaska Native	11,019	10.1
Asian	1,255	1.1
Native Hawaiian & Other Pacific Islander	99	0.1
Two or More Races	4,801	3.7
Under 5 years	7,330	6.7
Under 18 years	25,566	23.4
65 years and over	18,548	17.0

Source: United States Census Bureau, 2016 Population Estimates

## Health Status Indicators 2012-2016

### Nativity

Percent of Low Birth Weight Infants	6.8
Percent of Mothers Receiving Care in 1st Trimester	73.0
○ Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	16.7
Percent of Births Less Than 37 Wks. of Gestation	9.0
● Average Age of Mother	27.3
Teenage Birth Rate <sup>2</sup>	15.1
Percent White Births	71.5
Percent American Indian Births	18.8
○ Percent Unmarried	43.4
○ Percent WIC births	39.6
○ Percent Breastfeeding at discharge	83.4
● Percent Payment-Private Insurance	41.7
○ Percent Payment-Medicaid	38.8
● Percent C-Section	20.0

●Denotes a health status indicator which is significantly lower than the state average.  
○Denotes a health status indicator which is significantly higher than the state average.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

● All Causes	641.3
Heart Disease	146.6
Malignant Neoplasms (Cancer)	150.6
Trachea, Bronchus, & Lung	39.8
● Colon, Rectum, & Anus	11.6
Female Breast	14.6
Pancreas	11.2
Prostate	17.0
Leukemia	9.4
● Chronic Lower Respiratory Diseases	30.7
● Alzheimer's Disease	27.6
● Cerebrovascular Disease	26.7
Accidents	47.4
Motor Vehicle Accidents	13.4
● Diabetes	13.0
● Influenza and Pneumonia	11.8
Intentional Self-Harm (Suicide)	18.7
Chronic Liver Disease and Cirrhosis	16.2
Infant Mortality	6.25

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	984
2. Malignant Neoplasms (Cancer)	975
3. Accidents	269
4. Chronic Lower Respiratory Diseases	206
5. Alzheimer's Disease	194
6. Cerebrovascular Disease	180
7. Unspecified Dementia	116
8. Intentional Self-Harm (Suicide)	98
9. Chronic Liver Disease and Cirrhosis	92
10. Diabetes	85
Percent of Deaths due to tobacco use	17.8
Median age at death	77

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Perkins County

## Demographic Information



Perkins County is located in northwestern South Dakota and averages 1.0 persons per square mile.

### 2016 Population Information

Subject	Number	Percent
Total population	2,983	100.0
White	2,869	96.2
Black or African American	17	0.6
American Indian & Alaska Native	55	1.8
Asian	8	0.3
Native Hawaiian & Other Pacific Islander	2	0.1
Two or More Races	32	1.1
Under 5 years	168	5.6
Under 18 years	626	21.0
65 years and over	711	23.8

Source: United States Census Bureau, 2016 Population Estimates

## Health Status Indicators 2012-2016

### Natality

Percent of Low Birth Weight Infants	5.6
Percent of Mothers Receiving Care in 1st Trimester	67.4
Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	12.9
Percent of Births Less Than 37 Wks. of Gestation	6.7
○ Average Age of Mother	28.8
Teenage Birth Rate <sup>2</sup>	LNE
Percent White Births	97.1
Percent American Indian Births	2.3
● Percent Unmarried	23.6
Percent WIC births	33.9
Percent Breastfeeding at discharge	83.2
Percent Payment-Private Insurance	62.3
● Percent Payment-Medicaid	22.8
Percent C-Section	29.2

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

All Causes	726.3
Heart Disease	126.3
Malignant Neoplasms (Cancer)	156.6
Trachea, Bronchus, & Lung	24.2
Colon, Rectum, & Anus	18.0
Female Breast	39.0
Pancreas	LNE
Prostate	LNE
Leukemia	LNE
Chronic Lower Respiratory Diseases	56.9
Alzheimer's Disease	38.0
Cerebrovascular Disease	29.6
Accidents	51.5
Motor Vehicle Accidents	24.6
Diabetes	16.8
Influenza and Pneumonia	27.1
Intentional Self-Harm (Suicide)	LNE
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	LNE

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Malignant Neoplasms (Cancer)	42
2. Heart Disease	40
3. Chronic Lower Respiratory Diseases	19
4. Alzheimer's Disease	15
5. Cerebrovascular Disease	11
6. Accidents	9
7. Diabetes	6
T8. Influenza and Pneumonia	5
T8. Aortic Aneurysm and Dissection	5
Percent of Deaths due to tobacco use	19.7
Median age at death	83

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Potter County

## Demographic Information



Potter County is located in north central South Dakota and averages 2.7 persons per square mile.

### 2015 Population Information

Subject	Number	Percent
Total population	2,299	100.0
White	2,194	95.4
Black or African American	9	0.4
American Indian & Alaska Native	48	2.1
Asian	12	0.5
Native Hawaiian & Other Pacific Islander	1	0.0
Two or More Races	35	1.5
Under 5 years	113	4.9
Under 18 years	476	20.7
65 years and over	690	30.0

Source: United States Census Bureau, 2016 Population Estimates

## Health Status Indicators 2012-2016

### Nativity

Percent of Low Birth Weight Infants	5.8
● Percent of Mothers Receiving Care in 1st Trimester	55.4
Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	11.6
Percent of Births Less Than 37 Wks. of Gestation	7.4
○ Average Age of Mother	28.7
Teenage Birth Rate <sup>2</sup>	LNE
Percent White Births	91.7
Percent American Indian Births	6.6
● Percent Unmarried	24.0
● Percent WIC births	22.3
Percent Breastfeeding at discharge	86.0
○ Percent Payment-Private Insurance	76.0
● Percent Payment-Medicaid	20.7
Percent C-Section	22.3

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

All Causes	747.7
Heart Disease	122.9
Malignant Neoplasms (Cancer)	137.7
Trachea, Bronchus, & Lung	51.4
Colon, Rectum, & Anus	LNE
Female Breast	LNE
Pancreas	11.2
Prostate	32.6
Leukemia	LNE
Chronic Lower Respiratory Diseases	47.5
Alzheimer's Disease	43.4
● Cerebrovascular Disease	20.5
○ Accidents	112.0
Motor Vehicle Accidents	35.3
Diabetes	33.7
Influenza and Pneumonia	24.8
Intentional Self-Harm (Suicide)	LNE
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	LNE

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	37
2. Malignant Neoplasms (Cancer)	36
3. Alzheimer's Disease	17
4. Accidents	15
5. Chronic Lower Respiratory Diseases	13
6. Diabetes	9
7. Cerebrovascular Disease	8
T8. Influenza and Pneumonia	7
T8. Septicemia	7
T10. Unspecified Dementia	5
T10. Essential (Primary) Hypertension and Hypertensive Renal Disease	5
Percent of Deaths due to tobacco use	15.6
Median age at death	85

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Roberts County

## Demographic Information



Roberts County is located in the extreme northeastern corner of the state and averages 9.2 persons per square mile.

### 2016 Population Information

Subject	Number	Percent
Total population	10,255	100.0
White	5,985	58.4
Black or African American	72	0.7
American Indian & Alaska Native	3,822	37.3
Asian	33	0.3
Native Hawaiian & Other Pacific Islander	2	0.0
Two or More Races	341	3.3
Under 5 years	902	8.8
Under 18 years	2,881	28.1
65 years and over	1,936	18.9

Source: United States Census Bureau, 2016 Population Estimates

## Health Status Indicators 2012-2016

### Natality

● Percent of Low Birth Weight Infants	4.3
● Percent of Mothers Receiving Care in 1st Trimester	57.8
○ Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	28.2
Percent of Births Less Than 37 Wks. of Gestation	9.7
● Average Age of Mother	26.0
○ Teenage Birth Rate <sup>2</sup>	31.8
Percent White Births	30.7
Percent American Indian Births	59.5
○ Percent Unmarried	64.2
○ Percent WIC births	58.5
● Percent Breastfeeding at discharge	68.9
● Percent Payment-Private Insurance	33.1
○ Percent Payment-Medicaid	57.6
Percent C-Section	24.7

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

○ All Causes	854.8
Heart Disease	172.1
Malignant Neoplasms (Cancer)	178.9
Trachea, Bronchus, & Lung	37.7
Colon, Rectum, & Anus	18.6
Female Breast	16.1
Pancreas	8.0
Prostate	20.2
Leukemia	7.3
Chronic Lower Respiratory Diseases	47.6
Alzheimer's Disease	27.0
Cerebrovascular Disease	48.6
○ Accidents	80.6
○ Motor Vehicle Accidents	41.7
Diabetes	29.7
Influenza and Pneumonia	18.4
Intentional Self-Harm (Suicide)	30.5
Chronic Liver Disease and Cirrhosis	26.8
Infant Mortality	5.54

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	131
2. Malignant Neoplasms (Cancer)	124
3. Accidents	45
4. Chronic Lower Respiratory Diseases	37
5. Cerebrovascular Disease	36
6. Alzheimer's Disease	22
7. Diabetes	20
8. Chronic Liver Disease and Cirrhosis	14
9. Intentional Self-Harm (Suicide)	13
10. Influenza and Pneumonia	12
Percent of Deaths due to tobacco use	17.8
Median age at death	77

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Sanborn County

## Demographic Information



Sanborn County is located in east central South Dakota and averages 4.1 persons per square mile.

## 2016 Population Information

Subject	Number	Percent
Total population	2,396	100.0
White	2,339	97.6
Black or African American	8	0.3
American Indian & Alaska Native	10	0.4
Asian	6	0.3
Native Hawaiian & Other Pacific Islander	0	0.0
Two or More Races	33	1.4
Under 5 years	196	8.2
Under 18 years	570	23.8
65 years and over	444	18.5

Source: United States Census Bureau, 2016 Population Estimates

## Health Status Indicators 2012-2016

### Nativity

Percent of Low Birth Weight Infants	3.8
Percent of Mothers Receiving Care in 1st Trimester	75.4
Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	16.3
Percent of Births Less Than 37 Wks. of Gestation	5.4
Average Age of Mother	28.1
Teenage Birth Rate <sup>2</sup>	LNE
Percent White Births	99.5
Percent American Indian Births	LNE
● Percent Unmarried	24.5
● Percent WIC births	25.3
Percent Breastfeeding at discharge	82.6
○ Percent Payment-Private Insurance	78.3
● Percent Payment-Medicaid	20.1
Percent C-Section	31.5

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

All Causes	770.4
Heart Disease	145.2
Malignant Neoplasms (Cancer)	169.0
Trachea, Bronchus, & Lung	52.6
Colon, Rectum, & Anus	LNE
Female Breast	LNE
Pancreas	14.1
Prostate	LNE
Leukemia	LNE
Chronic Lower Respiratory Diseases	48.2
○ Alzheimer's Disease	116.1
Cerebrovascular Disease	34.9
Accidents	36.8
Motor Vehicle Accidents	LNE
Diabetes	18.1
Influenza and Pneumonia	11.1
Intentional Self-Harm (Suicide)	LNE
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	LNE

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	31
2. Malignant Neoplasms (Cancer)	29
3. Alzheimer's Disease	27
4. Chronic Lower Respiratory Diseases	10
5. Cerebrovascular Disease	8
6. Accidents	6
Percent of Deaths due to tobacco use	13.6
Median age at death	83

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Spink County

## Demographic Information



Spink County is located in the center of eastern South Dakota and averages 4.3 persons per square mile.

### 2016 Population Information

Subject	Number	Percent
Total population	6,420	100.0
White	6,190	96.4
Black or African American	41	0.6
American Indian & Alaska Native	111	1.7
Asian	7	0.1
Native Hawaiian & Other Pacific Islander	0	0.0
Two or More Races	71	1.1
Under 5 years	371	5.8
Under 18 years	1,488	23.2
65 years and over	1,311	20.4

Source: United States Census Bureau, 2016 Population Estimates

## Health Status Indicators 2012-2016

### Nativity

Percent of Low Birth Weight Infants	4.6
Percent of Mothers Receiving Care in 1st Trimester	67.4
Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	15.9
Percent of Births Less Than 37 Wks. of Gestation	8.7
○ Average Age of Mother	28.5
● Teenage Birth Rate <sup>2</sup>	5.8
Percent White Births	96.7
Percent American Indian Births	LNE
● Percent Unmarried	27.6
● Percent WIC births	26.4
Percent Breastfeeding at discharge	81.8
○ Percent Payment-Private Insurance	75.6
● Percent Payment-Medicaid	19.5
Percent C-Section	29.1

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

All Causes	731.3
Heart Disease	158.9
Malignant Neoplasms (Cancer)	163.4
Trachea, Bronchus, & Lung	40.9
Colon, Rectum, & Anus	21.3
Female Breast	37.6
Pancreas	12.1
Prostate	LNE
Leukemia	LNE
Chronic Lower Respiratory Diseases	46.5
Alzheimer's Disease	52.7
● Cerebrovascular Disease	22.1
Accidents	69.1
Motor Vehicle Accidents	26.4
Diabetes	33.2
Influenza and Pneumonia	30.7
Intentional Self-Harm (Suicide)	LNE
Chronic Liver Disease and Cirrhosis	7.4
Infant Mortality	LNE

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	92
2. Malignant Neoplasms (Cancer)	80
3. Alzheimer's Disease	36
4. Accidents	27
5. Chronic Lower Respiratory Diseases	26
6. Influenza and Pneumonia	21
7. Diabetes	15
8. Cerebrovascular Disease	14
T9. Essential (Primary) Hypertension and Hypertensive Renal Disease	7
T9. Pneumonitis Due to Solids and Liquids	7

Percent of Deaths due to tobacco use	10.4
Median age at death	82

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Stanley County

## Demographic Information



Stanley County is located in the center of the state and averages 2.1 persons per square mile.

### 2016 Population Information

Subject	Number	Percent
Total population	2,993	100.0
White	2,676	89.4
Black or African American	26	0.9
American Indian & Alaska Native	203	6.8
Asian	10	0.3
Native Hawaiian & Other Pacific Islander	0	0.0
Two or More Races	78	2.6
Under 5 years	200	6.7
Under 18 years	734	24.5
65 years and over	564	18.8

Source: United States Census Bureau, 2016 Population Estimates

## Health Status Indicators 2012-2016

### Natality

Percent of Low Birth Weight Infants	4.6
● Percent of Mothers Receiving Care in 1st Trimester	57.5
Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	11.7
Percent of Births Less Than 37 Wks. of Gestation	8.1
○ Average Age of Mother	28.6
Teenage Birth Rate <sup>2</sup>	LNE
Percent White Births	86.1
Percent American Indian Births	10.3
Percent Unmarried	33.0
Percent WIC births	33.2
Percent Breastfeeding at discharge	81.1
Percent Payment-Private Insurance	67.3
Percent Payment-Medicaid	28.1
○ Percent C-Section	41.6

●Denotes a health status indicator which is significantly lower than the state average.  
○Denotes a health status indicator which is significantly higher than the state average.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

● All Causes	571.1
● Heart Disease	92.0
Malignant Neoplasms (Cancer)	167.4
Trachea, Bronchus, & Lung	48.5
Colon, Rectum, & Anus	23.9
Female Breast	LNE
Pancreas	16.3
Prostate	LNE
Leukemia	LNE
Chronic Lower Respiratory Diseases	82.1
Alzheimer's Disease	LNE
● Cerebrovascular Disease	15.1
● Accidents	19.6
Motor Vehicle Accidents	LNE
Diabetes	22.0
Influenza and Pneumonia	LNE
○ Intentional Self-Harm (Suicide)	57.9
Chronic Liver Disease and Cirrhosis	23.9
Infant Mortality	LNE

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Malignant Neoplasms (Cancer)	31
2. Heart Disease	16
3. Chronic Lower Respiratory Diseases	14
4. Intentional Self-Harm (Suicide)	9
Percent of Deaths due to tobacco use	28.0
Median age at death	68

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Sully County

## Demographic Information



Sully County is located in the central region of the state and averages 1.4 persons per square mile.

## 2016 Population Information

Subject	Number	Percent
Total population	1,421	100.0
White	1,348	94.9
Black or African American	6	0.4
American Indian & Alaska Native	36	2.5
Asian	1	0.1
Native Hawaiian & Other Pacific Islander	0	0.0
Two or More Races	30	2.1
Under 5 years	93	6.5
Under 18 years	315	22.2
65 years and over	292	20.5

Source: United States Census Bureau, 2016 Population Estimates

## Health Status Indicators 2012-2016

### Nativity

Percent of Low Birth Weight Infants	4.2
● Percent of Mothers Receiving Care in 1st Trimester	48.6
Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	12.5
Percent of Births Less Than 37 Wks. of Gestation	8.3
Average Age of Mother	27.5
Teenage Birth Rate <sup>2</sup>	LNE
Percent White Births	84.5
Percent American Indian Births	11.3
● Percent Unmarried	25.0
Percent WIC births	23.6
Percent Breastfeeding at discharge	76.4
Percent Payment-Private Insurance	69.4
Percent Payment-Medicaid	29.2
Percent C-Section	25.0

●Denotes a health status indicator which is significantly lower than the state average.  
○Denotes a health status indicator which is significantly higher than the state average.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

● All Causes	441.3
● Heart Disease	70.9
Malignant Neoplasms (Cancer)	161.4
Trachea, Bronchus, & Lung	LNE
Colon, Rectum, & Anus	LNE
Female Breast	LNE
Pancreas	31.5
Prostate	LNE
Leukemia	LNE
Chronic Lower Respiratory Diseases	27.2
Alzheimer's Disease	LNE
Cerebrovascular Disease	LNE
Accidents	28.8
Motor Vehicle Accidents	LNE
Diabetes	LNE
Influenza and Pneumonia	LNE
Intentional Self-Harm (Suicide)	LNE
Chronic Liver Disease and Cirrhosis	LNE
Infant Mortality	LNE

### Leading Causes of Death

1. Malignant Neoplasms (Cancer)
2. Heart Disease

	Total Deaths
Percent of Deaths due to tobacco use	20.8
Median age at death	75

●Denotes a health status indicator which is significantly lower than the state average.  
○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Todd County

## Demographic Information



Todd County is located in south central South Dakota, along the Nebraska border and averages 6.9 persons per square mile.

### 2015 Population Information

Subject	Number	Percent
Total population	10,155	100.0
White	930	9.2
Black or African American	46	0.5
American Indian & Alaska Native	8,913	87.8
Asian	17	0.2
Native Hawaiian & Other Pacific Islander	0	0.0
Two or More Races	249	2.5
Under 5 years	1,288	12.7
Under 18 years	4,142	40.8
65 years and over	795	7.8

Source: United States Census Bureau, 2018 Population Estimates

## Health Status Indicators 2012-2016

### Nativity

Percent of Low Birth Weight Infants	7.9
● Percent of Mothers Receiving Care in 1st Trimester	39.2
○ Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	22.5
○ Percent of Births Less Than 37 Wks. of Gestation	11.2
● Average Age of Mother	25.1
○ Teenage Birth Rate <sup>2</sup>	56.7
Percent White Births	4.3
Percent American Indian Births	91.8
○ Percent Unmarried	84.5
○ Percent WIC births	71.6
● Percent Breastfeeding at discharge	62.6
● Percent Payment-Private Insurance	5.1
○ Percent Payment-Medicaid	52.3
○ Percent C-Section	36.4

●Denotes a health status indicator which is significantly lower than the state average.  
○Denotes a health status indicator which is significantly higher than the state average.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

○ All Causes	1,310.9
○ Heart Disease	219.0
Malignant Neoplasms (Cancer)	201.5
Trachea, Bronchus, & Lung	39.8
Colon, Rectum, & Anus	33.0
Female Breast	LNE
Pancreas	LNE
Prostate	37.9
Leukemia	19.4
Chronic Lower Respiratory Diseases	66.1
Alzheimer's Disease	19.8
Cerebrovascular Disease	59.6
○ Accidents	148.9
○ Motor Vehicle Accidents	85.5
○ Diabetes	93.1
○ Influenza and Pneumonia	42.0
○ Intentional Self-Harm (Suicide)	72.2
○ Chronic Liver Disease and Cirrhosis	82.2
Infant Mortality	12.58

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	68
T2. Malignant Neoplasms (Cancer)	65
T2. Accidents	65
4. Intentional Self-Harm (Suicide)	33
5. Diabetes	32
6. Chronic Liver Disease and Cirrhosis	30
7. Chronic Lower Respiratory Diseases	21
8. Cerebrovascular Disease	18
9. Influenza and Pneumonia	13
10. Mental and Behavioral Disorders Due to Use of Alcohol	10
Percent of Deaths due to tobacco use	12.9
Median age at death	57

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Tripp County

## Demographic Information



Tripp County is located along the Nebraska border in south central South Dakota and averages 3.5 persons per square mile.

### 2016 Population Information

Subject	Number	Percent
Total population	5,492	100.0
White	4,517	82.2
Black or African American	20	0.4
American Indian & Alaska Native	785	14.3
Asian	33	0.6
Native Hawaiian & Other Pacific Islander	1	0.0
Two or More Races	136	2.5
Under 5 years	349	6.4
Under 18 years	1,255	22.9
65 years and over	1,204	21.9

Source: United States Census Bureau, 2016 Population Estimates

## Health Status Indicators 2012-2016

### Nativity

Percent of Low Birth Weight Infants	4.8
Percent of Mothers Receiving Care in 1st Trimester	69.5
Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	18.5
Percent of Births Less Than 37 Wks. of Gestation	9.7
Average Age of Mother	27.2
Teenage Birth Rate <sup>2</sup>	10.4
Percent White Births	65.8
Percent American Indian Births	30.2
Percent Unmarried	44.9
○ Percent WIC births	45.7
Percent Breastfeeding at discharge	71.2
● Percent Payment-Private Insurance	48.6
○ Percent Payment-Medicaid	48.3
○ Percent C-Section	38.1

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

All Causes	731.0
Heart Disease	180.0
Malignant Neoplasms (Cancer)	150.1
Trachea, Bronchus, & Lung	36.0
Colon, Rectum, & Anus	23.7
Female Breast	LNE
Pancreas	7.2
Prostate	LNE
Leukemia	LNE
Chronic Lower Respiratory Diseases	39.0
○ Alzheimer's Disease	58.3
Cerebrovascular Disease	32.1
Accidents	42.1
Motor Vehicle Accidents	11.1
● Diabetes	12.2
Influenza and Pneumonia	17.2
Intentional Self-Harm (Suicide)	LNE
Chronic Liver Disease and Cirrhosis	31.1
Infant Mortality	8.52

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	96
2. Malignant Neoplasms (Cancer)	73
3. Alzheimer's Disease	39
T4. Chronic Lower Respiratory Diseases	20
T4. Cerebrovascular Disease	20
6. Accidents	18
7. Influenza and Pneumonia	10
T8. Diabetes	7
T8. Chronic Liver Disease and Cirrhosis	7
10. Pneumonitis Due to Solids and Liquids	5
Percent of Deaths due to tobacco use	20.2
Median age at death	84

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Turner County

## Demographic Information



Turner County is located in southeastern South Dakota and averages 13.5 persons per square mile.

### 2016 Population Information

Subject	Number	Percent
Total population	8,317	100.0
White	8,101	97.4
Black or African American	33	0.4
American Indian & Alaska Native	80	1.0
Asian	17	0.2
Native Hawaiian & Other Pacific Islander	6	0.1
Two or More Races	80	1.0
Under 5 years	493	5.9
Under 18 years	1,954	23.5
65 years and over	1,731	20.8

Source: United States Census Bureau, 2016 Population Estimates

## Health Status Indicators 2012-2016

### Nativity

● Percent of Low Birth Weight Infants	3.7
○ Percent of Mothers Receiving Care in 1st Trimester	82.6
● Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	9.7
Percent of Births Less Than 37 Wks. of Gestation	6.4
○ Average Age of Mother	29.1
Teenage Birth Rate <sup>2</sup>	LNE
Percent White Births	98.8
Percent American Indian Births	0.7
● Percent Unmarried	21.4
● Percent WIC births	21.9
Percent Breastfeeding at discharge	82.5
○ Percent Payment-Private Insurance	74.9
● Percent Payment-Medicaid	19.8
Percent C-Section	26.2

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

All Causes	749.6
Heart Disease	164.5
Malignant Neoplasms (Cancer)	157.4
Trachea, Bronchus, & Lung	36.5
Colon, Rectum, & Anus	20.3
Female Breast	25.1
Pancreas	7.1
Prostate	15.8
Leukemia	11.0
○ Chronic Lower Respiratory Diseases	65.0
○ Alzheimer's Disease	73.3
● Cerebrovascular Disease	24.1
Accidents	50.1
Motor Vehicle Accidents	31.2
Diabetes	13.9
Influenza and Pneumonia	9.7
Intentional Self-Harm (Suicide)	11.8
Chronic Liver Disease and Cirrhosis	14.5
Infant Mortality	6.90

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	124
2. Malignant Neoplasms (Cancer)	102
3. Alzheimer's Disease	61
4. Chronic Lower Respiratory Diseases	44
5. Accidents	28
6. Cerebrovascular Disease	20
7. Diabetes	11
8. Influenza and Pneumonia	9
T9. Chronic Liver Disease and Cirrhosis	7
T9. Essential (Primary) Hypertension and Hypertensive Renal Disease	7
T9. Parkinson's Disease	7

Percent of Deaths due to tobacco use	13.6
Median age at death	82

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Union County

## Demographic Information



Union County is located in the southeastern corner of the state and averages 31.3 persons per square mile.

### 2016 Population Information

Subject	Number	Percent
Total population	14,934	100.0
White	14,218	95.2
Black or African American	150	1.0
American Indian & Alaska Native	130	0.9
Asian	187	1.3
Native Hawaiian & Other Pacific Islander	10	0.1
Two or More Races	239	1.6
Under 5 years	848	5.7
Under 18 years	3,608	24.2
65 years and over	2,483	16.6

Source: United States Census Bureau, 2016 Population Estimates

## Health Status Indicators 2012-2016

### Nativity

Percent of Low Birth Weight Infants	6.6
○ Percent of Mothers Receiving Care in 1st Trimester	84.9
● Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	9.3
Percent of Births Less Than 37 Wks. of Gestation	8.1
○ Average Age of Mother	28.8
● Teenage Birth Rate <sup>2</sup>	2.4
Percent White Births	94.8
Percent American Indian Births	1.4
● Percent Unmarried	23.3
● Percent WIC births	14.9
Percent Breastfeeding at discharge	82.2
○ Percent Payment-Private Insurance	76.3
● Percent Payment-Medicaid	20.6
○ Percent C-Section	32.3

●Denotes a health status indicator which is significantly lower than the state average.  
○Denotes a health status indicator which is significantly higher than the state average.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

● All Causes	635.2
Heart Disease	151.8
Malignant Neoplasms (Cancer)	151.1
Trachea, Bronchus, & Lung	44.7
Colon, Rectum, & Anus	13.7
Female Breast	19.4
Pancreas	10.2
Prostate	LNE
Leukemia	9.9
Chronic Lower Respiratory Diseases	41.8
Alzheimer's Disease	26.2
● Cerebrovascular Disease	24.8
● Accidents	28.7
● Motor Vehicle Accidents	7.0
Diabetes	14.6
Influenza and Pneumonia	20.7
Intentional Self-Harm (Suicide)	12.7
Chronic Liver Disease and Cirrhosis	11.6
Infant Mortality	6.25

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	146
2. Malignant Neoplasms (Cancer)	140
3. Chronic Lower Respiratory Diseases	38
4. Alzheimer's Disease	25
T5. Accidents	24
T5. Cerebrovascular Disease	24
7. Influenza and Pneumonia	19
8. Essential (Primary) Hypertension and Hypertensive Renal Disease	14
9. Diabetes	13
10. Unspecified Dementia	12
Percent of Deaths due to tobacco use	18.3
Median age at death	80

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Walworth County

## Demographic Information



Walworth County is located in north central South Dakota, near the North Dakota border and averages 7.7 persons per square mile.

### 2015 Population Information

Subject	Number	Percent
Total population	5,610	100.0
White	4,466	79.6
Black or African American	32	0.6
American Indian & Alaska Native	808	14.4
Asian	116	2.1
Native Hawaiian & Other Pacific Islander	1	0.0
Two or More Races	187	3.3
Under 5 years	387	6.9
Under 18 years	1,314	23.4
65 years and over	1,379	24.6

Source: United States Census Bureau, 2016 Population Estimates

## Health Status Indicators 2012-2016

### Nativity

Percent of Low Birth Weight Infants	5.7
● Percent of Mothers Receiving Care in 1st Trimester	62.3
Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	18.5
Percent of Births Less Than 37 Wks. of Gestation	9.6
● Average Age of Mother	26.9
Teenage Birth Rate <sup>2</sup>	LNE
Percent White Births	65.9
Percent American Indian Births	26.4
Percent Unmarried	41.8
○ Percent WIC births	43.0
Percent Breastfeeding at discharge	72.8
● Percent Payment-Private Insurance	47.8
○ Percent Payment-Medicaid	42.5
Percent C-Section	27.2

●Denotes a health status indicator which is significantly lower than the state average.  
○Denotes a health status indicator which is significantly higher than the state average.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

○ All Causes	824.8
Heart Disease	159.3
Malignant Neoplasms (Cancer)	153.5
Trachea, Bronchus, & Lung	46.2
Colon, Rectum, & Anus	10.4
Female Breast	25.8
Pancreas	LNE
Prostate	13.6
Leukemia	5.0
Chronic Lower Respiratory Diseases	50.5
○ Alzheimer's Disease	73.3
Cerebrovascular Disease	36.7
Accidents	68.7
Motor Vehicle Accidents	28.3
Diabetes	31.4
Influenza and Pneumonia	29.0
Intentional Self-Harm (Suicide)	32.2
Chronic Liver Disease and Cirrhosis	20.0
Infant Mortality	LNE

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	94
2. Malignant Neoplasms (Cancer)	75
3. Alzheimer's Disease	53
4. Chronic Lower Respiratory Diseases	27
5. Accidents	22
T6. Cerebrovascular Disease	19
T6. Diabetes	19
8. Influenza and Pneumonia	18
9. Unspecified Dementia	9
T10. Chronic Liver Disease and Cirrhosis	8
T10. Nephritis, Nephrotic Syndrome, and Nephrosis	8

Percent of Deaths due to tobacco use	16.4
Median age at death	83

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Yankton County

## Demographic Information



Yankton County is located in southeastern South Dakota on the Nebraska border and averages 43.1 persons per square mile.

### 2016 Population Information

Subject	Number	Percent
Total population	22,616	100.0
White	20,875	92.3
Black or African American	475	2.1
American Indian & Alaska Native	700	3.1
Asian	174	0.8
Native Hawaiian & Other Pacific Islander	9	0.0
Two or More Races	383	1.7
Under 5 years	1,309	5.8
Under 18 years	4,759	21.0
65 years and over	4,109	18.2

Source: United States Census Bureau, 2016 Population Estimates

## Health Status Indicators 2012-2016

### Natality

Percent of Low Birth Weight Infants	7.4
○ Percent of Mothers Receiving Care in 1st Trimester	84.6
○ Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	19.9
Percent of Births Less Than 37 Wks. of Gestation	7.4
● Average Age of Mother	27.3
Teenage Birth Rate <sup>2</sup>	11.3
Percent White Births	89.8
Percent American Indian Births	4.3
Percent Unmarried	40.3
Percent WIC births	36.3
Percent Breastfeeding at discharge	73.3
○ Percent Payment-Private Insurance	65.0
Percent Payment-Medicaid	33.2
○ Percent C-Section	29.6

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

All Causes	672.3
Heart Disease	159.0
Malignant Neoplasms (Cancer)	143.1
Trachea, Bronchus, & Lung	40.8
Colon, Rectum, & Anus	14.3
Female Breast	17.6
● Pancreas	4.6
Prostate	21.5
Leukemia	11.5
Chronic Lower Respiratory Diseases	53.7
Alzheimer's Disease	32.2
Cerebrovascular Disease	32.9
Accidents	49.2
Motor Vehicle Accidents	16.9
Diabetes	22.2
Influenza and Pneumonia	19.2
Intentional Self-Harm (Suicide)	14.4
● Chronic Liver Disease and Cirrhosis	5.8
Infant Mortality	8.61

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	280
2. Malignant Neoplasms (Cancer)	227
3. Chronic Lower Respiratory Diseases	90
4. Accidents	71
5. Alzheimer's Disease	68
6. Cerebrovascular Disease	63
7. Diabetes	41
8. Influenza and Pneumonia	36
9. Intentional Self-Harm (Suicide)	17
T10. Unspecified Dementia	13
T10. Essential (Primary) Hypertension	13
Hypertensive Renal Disease	
T10. Septicemia	13
Percent of Deaths due to tobacco use	21.2
Median age at death	81

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.


<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics

# Ziebach County

## Demographic Information



Ziebach County is located in north central South Dakota and averages 1.4 persons per square mile.

### 2016 Population Information

Subject	Number	Percent
Total population	2,801	100.0
White	680	24.3
Black or African American	13	0.5
American Indian & Alaska Native	1,999	71.4
Asian	3	0.1
Native Hawaiian & Other Pacific Islander	1	0.0
Two or More Races	105	3.7
Under 5 years	178	6.4
Under 18 years	915	32.7
65 years and over	210	7.5

Source: United States Census Bureau, 2016 Population Estimates

## Health Status Indicators 2012-2016

### Nativity

Percent of Low Birth Weight Infants	8.1
● Percent of Mothers Receiving Care in 1st Trimester	50.0
Percent of Mothers Who Used Tobacco While Pregnant <sup>1</sup>	19.7
Percent of Births Less Than 37 Wks. of Gestation	12.9
● Average Age of Mother	26.6
Teenage Birth Rate <sup>2</sup>	18.1
Percent White Births	13.3
Percent American Indian Births	80.9
○ Percent Unmarried	69.9
○ Percent WIC births	78.5
● Percent Breastfeeding at discharge	59.6
● Percent Payment-Private Insurance	17.8
○ Percent Payment-Medicaid	69.8
Percent C-Section	26.0

●Denotes a health status indicator which is significantly lower than the state average.  
○Denotes a health status indicator which is significantly higher than the state average.

<sup>1</sup>Data for mothers who used tobacco are self-reported.

<sup>2</sup>Teenage Birth rate is live births per 1,000 females age 15-17.

### Mortality<sup>3</sup>

All Causes	729.7
Heart Disease	180.4
● Malignant Neoplasms (Cancer)	45.5
Trachea, Bronchus, & Lung	LNE
Colon, Rectum, & Anus	LNE
Female Breast	LNE
Pancreas	LNE
Prostate	LNE
Leukemia	LNE
Chronic Lower Respiratory Diseases	76.6
Alzheimer's Disease	LNE
Cerebrovascular Disease	LNE
Accidents	94.7
Motor Vehicle Accidents	41.2
Diabetes	27.6
Influenza and Pneumonia	31.9
Intentional Self-Harm (Suicide)	17.1
Chronic Liver Disease and Cirrhosis	45.8
Infant Mortality	LNE

### Leading Causes of Death

Leading Causes of Death	Total Deaths
1. Heart Disease	15
2. Accidents	12
T3. Malignant Neoplasms (Cancer)	5
T3. Chronic Lower Respiratory Diseases	5
T3. Chronic Liver Disease and Cirrhosis	5

Percent of Deaths due to tobacco use	11.1
Median age at death	60

●Denotes a health status indicator which is significantly lower than the state average.

○Denotes a health status indicator which is significantly higher than the state average.

<sup>3</sup>All mortality rates except infant mortality are age-adjusted death rates per 100,000 population. Infant mortality is the number of infant (less than one year) deaths per 1,000 live births.

See technical notes for more information.

Source: South Dakota Department of Health, Office of Health Statistics